

Policymaker Summary of Working Group I (Scientific Assessment of Climate Change)

Executive Summary

1. *We are certain of the following:*

- there is a natural greenhouse effect which already keeps the Earth warmer than it would otherwise be;
- emissions resulting from human activities are substantially increasing the atmospheric concentrations of the greenhouse gases: carbon dioxide, methane, chlorofluorocarbons (CFCs) and nitrous oxide. These increases will enhance the greenhouse effect, resulting on average in an additional warming of the Earth's surface. The main greenhouse gas, water vapour, will increase in response to global warming and further enhance it.

2. *We calculate with confidence that:*

- some gases are potentially more effective than others at changing climate, and their relative effectiveness can be estimated. Carbon dioxide has been responsible for over half the enhanced greenhouse effect in the past, and is likely to remain so in the future;
- atmospheric concentrations of the long-lived gases (carbon dioxide, nitrous oxide and the CFCs) adjust only slowly to changes in emissions. Continued emissions of these gases at present rates would commit us to increased concentrations for centuries ahead. The longer emissions continue to increase at present day rates, the greater reductions would have to be for concentrations to stabilize at a given level;
- the long-lived gases would require immediate reductions in emissions from human activities of over 60% to stabilize their concentrations at today's levels; methane would require a 15-20% reduction.

3. *Based on current model results, we predict:*

- under the IPCC Business-as-Usual (Scenario A) emissions of greenhouse gases, a rate of increase of global-mean temperature during the next century of about 0.3°C per decade (with an uncertainty range of 0.2°C to 0.5°C per decade); this is greater than that seen over the past 10,000 years. This will result in a likely increase in global-mean temperature of about 1°C above the present value by 2025 and 3°C before the end of the next century. The rise will not be steady because of the influence of other factors;
- under the other IPCC emission scenarios which assume progressively increasing levels of controls, rates of

increase in global-mean temperature of about 0.2°C per decade (Scenario B), just above 0.1°C per decade (Scenario C) and about 0.1°C per decade (Scenario D);

- that land surfaces warm more rapidly than the ocean, and high northern latitudes warm more than the global-mean in winter;
- regional climate changes different from the global-mean, although our confidence in the prediction of the detail of regional changes is low. For example, temperature increases in Southern Europe and central North America are predicted to be higher than the global-mean, accompanied on average by reduced summer precipitation and soil moisture. There are less consistent predictions for the tropics and the southern hemisphere;
- under the IPCC Business-as-Usual emissions scenario, an average rate of global mean sea-level rise of about 6 cm per decade over the next century (with an uncertainty range of 3—10 cm per decade), mainly due to thermal expansion of the oceans and the melting of some land ice. The predicted rise is about 20 cm in global-mean sea level by 2030, and 65 cm by the end of the next century. There will be significant regional variations.

3.1 There are many uncertainties in our predictions particularly with regard to the timing, magnitude and regional patterns of climate change, due to our incomplete understanding of:

- sources and sinks of greenhouse gases, which affect predictions of future concentrations;
- clouds, which strongly influence the magnitude of climate change;
- oceans, which influence the timing and patterns of climate change;
- polar ice sheets which affect predictions of sea-level rise.

3.2 These processes are already partially understood, and we are confident that the uncertainties can be reduced by further research. However, the complexity of the system means that we cannot rule out surprises.

4. *Our judgement is that:*

- Global mean surface air temperature has increased by 0.3°C to 0.6°C over the last 100 years, with the five global average warmest years being in the 1980s. Over

Scientific Assessment of Climate Change

the same period global sea level has increased by 10-20 cm. These increases have not been smooth with time, nor uniform over the globe.

- The size of this warming is broadly consistent with predictions of climate models, but it is also of the same magnitude as natural climate variability. Thus the observed increase could be largely due to this natural variability; alternatively this variability and other human factors could have offset a still larger human-induced greenhouse warming. The unequivocal detection of the enhanced greenhouse effect from observations is not likely for a decade or more.
- There is no firm evidence that climate has become more variable over the last few decades. However, with an increase in the mean temperature, episodes of high temperatures will most likely become more frequent in the future, and cold episodes less frequent.
- Ecosystems affect climate, and will be affected by a changing climate and by increasing carbon dioxide concentrations. Rapid changes in climate will change the composition of ecosystems; some species will benefit while others will be unable to migrate or adapt fast enough and may become extinct. Enhanced levels of carbon dioxide may increase productivity and efficiency of water use of vegetation. The effect of warming on biological processes, although poorly understood, may increase the atmospheric concentrations of natural greenhouse gases.

5. ***To improve our predictive capability, we need:***

- to understand better the various climate-related processes, particularly those associated with clouds, oceans and the carbon cycle;
- to improve the systematic observation of climate-related variables on a global basis, and further investigate changes which took place in the past;
- to develop improved models of the Earth's climate system;
- to increase support for national and international climate research activities, especially in developing countries;
- to facilitate international exchange of climate data.

1. Introduction: what is the issue?

1.0.1 There is concern that human activities may be inadvertently changing the climate of the globe through the enhanced greenhouse effect, by past and continuing emissions of carbon dioxide and other gases which will cause the temperature of the Earth's surface to increase—popularly termed the “global warming”. If this occurs, consequent changes may have a significant impact on society.

1.0.2 The purpose of the Working Group I report, as determined by the first meeting of IPCC, is to provide a scientific assessment of:

- 1) the factors which may affect climate change during the next century especially those which are due to human activity;
- 2) the responses of the atmosphere-ocean-land-ice system;
- 3) current capabilities of modelling global and regional climate changes and their predictability;
- 4) the past climate record and presently observed climate anomalies.

1.0.3 On the basis of this assessment, the report presents current knowledge regarding predictions of climate change (including sea-level rise and the effects on ecosystems) over the next century, the timing of changes together with an assessment of the uncertainties associated with these predictions.

1.0.4 This Policymaker Summary aims to bring out those elements of the main report which have the greatest relevance to policy formulation, in answering the following questions:

- What factors determine global climate?
- What are the greenhouse gases, and how and why are they increasing?
- Which gases are the most important?
- How much do we expect the climate to change?
- How much confidence do we have in our predictions?
- Will the climate of the future be very different?
- Have human activities already begun to change global climate?
- How much will sea-level rise?
- What will be the effects on ecosystems?
- What should be done to reduce uncertainties, and how long will this take?

1.0.5 This report is intended to respond to the practical needs of the policymaker. It is neither an academic review, nor a plan for a new research programme. Uncertainties attach to almost every aspect of the issue, yet policymakers are looking for clear guidance from scientists; hence authors have been asked to provide their best estimates wherever possible, together with an assessment of the uncertainties.

1.0.6 This report is a summary of our understanding in 1990. Although continuing research will deepen this understanding and require the report to be updated at frequent intervals, basic conclusions concerning the reality of the enhanced greenhouse effect and its potential to alter global climate are unlikely to change significantly. Nevertheless, the complexity of the system may give rise to surprises.

2. What factors determine global climate ?

2.0.1 There are many factors, both natural and of human origin, that determine the climate of the earth. We look first at those which are natural, and then see how human activities might contribute.

2.1 What natural factors are important?

2.1.1 The driving energy for weather and climate comes from the sun. The Earth intercepts solar radiation (including that in the short-wave, visible, part of the spectrum); about a third of it is reflected, the rest is absorbed by the different components (atmosphere, ocean, ice, land and biota) of the climate system. The energy absorbed from solar radiation is balanced (in the long term) by outgoing radiation from the Earth and atmosphere; this terrestrial radiation takes the form of long-wave invisible infra-red energy, and its magnitude is determined by the temperature of the Earth-atmosphere system.

2.1.2 There are several natural factors which can change the balance between the energy absorbed by the Earth and that emitted by it in the form of longwave infra-red radiation; these factors cause the radiative forcing on climate. The most obvious of these is a change in the output of energy from the Sun. There is direct evidence of such variability over the 11-year solar cycle, and longer period changes may also occur. Slow variations in the Earth's orbit affect the seasonal and latitudinal distribution of solar radiation; these were probably responsible for initiating the ice ages.

2.1.3 One of the most important factors is the greenhouse effect; a simplified explanation of which is as follows. Short-wave solar radiation can pass through the clear atmosphere relatively unimpeded. But long-wave terrestrial radiation emitted by the warm surface of the Earth is partially absorbed and then re-emitted by a number of trace gases in the cooler atmosphere above. Since, on average, the outgoing long-wave radiation balances the incoming solar radiation, both the atmosphere and the




Figure 3 A simplified diagram illustrating the greenhouse effect

surface will be warmer than they would be without the greenhouse gases.

2.1.4 The main natural greenhouse gases are not the major constituents, nitrogen and oxygen, but water vapour (the biggest contributor), carbon dioxide, methane, nitrous oxide, and ozone in the troposphere (the lowest 10–15 km of the atmosphere) and stratosphere.

2.1.5 Aerosols (small particles) in the atmosphere can also affect climate because they can reflect and absorb radiation. The most important natural perturbations result from explosive volcanic eruptions which affect concentrations in the lower stratosphere. Lastly, the climate has its own natural variability on all timescales and changes occur without any external influence.

2.2 How do we know that the natural greenhouse effect is real?

2.2.1 The greenhouse effect is real; it is a well understood effect, based on established scientific principles. We know that the greenhouse effect works in practice, for several reasons.


2.2.2 Firstly, the mean temperature of the Earth's surface is already warmer by about 33°C (assuming the same reflectivity of the earth) than it would be if the natural greenhouse gases were not present. Satellite observations of the radiation emitted from the Earth's surface and through the atmosphere demonstrate the effect of the greenhouse gases.

2.2.3 Secondly, we know the composition of the atmospheres of Venus, Earth and Mars are very different, and their surface temperatures are in general agreement with greenhouse theory.

2.2.4 Thirdly, measurements from ice cores going back 160,000 years show that the Earth's temperature closely paralleled the amount of carbon dioxide and methane in the atmosphere. Although we do not know the details of cause and effect, calculations indicate that changes in these greenhouse gases were part, but not all, of the reason for the large (5–7°C) global temperature swings between ice ages and interglacial periods.

2.3 How might human activities change global climate?

2.3.1 Naturally occurring greenhouse gases keep the Earth warm enough to be habitable. By increasing their concentrations, and by adding new greenhouse gases like chlorofluorocarbons (CFCs), humankind is capable of raising the global-average annual-mean surface-air temperature (which, for simplicity, is referred to as the "global temperature"), although we are uncertain about the rate at which this will occur. Strictly, this is an enhanced greenhouse effect—above that occurring due to natural greenhouse gas concentrations; the word "enhanced" is usually omitted, but it should not be forgotten. Other changes in climate are expected to result, for example changes in precipitation, and a global warming will cause sea levels to rise; these are discussed in more detail later.



Analysis of air trapped in Antarctic ice cores shows that methane and carbon dioxide concentrations were closely correlated with the local temperature over the last 160 000 years. Present-day concentrations of carbon dioxide are indicated.

2.3.2 There are other human activities which have the potential to affect climate. A change in the albedo (reflectivity) of the land, brought about by desertification or deforestation affects the amount of solar energy absorbed at the Earth's surface. Human-made aerosols, from sulphur emitted largely in fossil fuel combustion, can modify clouds and this may act to lower temperatures. Lastly, changes in ozone in the stratosphere due to CFCs may also influence climate.

3. What are the greenhouse gases and why are they increasing?

3.0.1 We are certain that the concentrations of greenhouse gases in the atmosphere have changed naturally on ice-age time-scales, and have been increasing since pre-industrial times due to human activities. The table below summarizes the present and pre-industrial abundances, current rates of change and present atmospheric lifetimes of greenhouse gases influenced by human activities. Carbon dioxide, methane, and nitrous oxide all have significant natural and human sources, while the chlorofluorocarbons are only produced industrially.

3.0.2 Two important greenhouse gases, water vapour and ozone, are not included in the table overleaf. Water vapour has the largest greenhouse effect, but its concentration in the troposphere is determined internally within the climate system, and, on a global scale, is not affected by human sources and sinks. Water vapour will increase in response to global warming and further enhance it: this process is included in climate models. The concentration of ozone is changing both in the stratosphere and the troposphere due to human activities, but it is difficult to quantify the changes from present observations.

3.0.3 For a thousand years prior to the industrial revolution, abundances of the greenhouse gases were relatively constant. However, as the world's population increased, as the world became more industrialized and as agriculture developed, the abundances of the greenhouse gases increased markedly. The figures overleaf illustrate this for carbon dioxide, methane, nitrous oxide and CFC-11.

3.0.4 Since the industrial revolution the combustion of fossil fuels and deforestation have led to an increase of 26% in carbon dioxide concentration in the atmosphere. We know the magnitude of the present day fossil-fuel source, but the input from deforestation cannot be estimated accurately. In addition, although about half of the emitted carbon dioxide stays in the atmosphere, we do not know well how much of the remainder is absorbed by the oceans and how much by terrestrial biota. Emissions of chlorofluorocarbons, used as aerosol propellants, solvents, refrigerants and foam blowing agents, are also well known; they were not present in the atmosphere before their invention in the 1930s.

3.0.5 The sources of methane and nitrous oxide are less well known. Methane concentrations have more than doubled because of rice production, cattle rearing, biomass burning, coal mining and ventilation of natural gas; also, fossil fuel combustion may have also contributed through chemical reactions in the atmosphere which reduce the rate of removal of methane. Nitrous oxide has increased by about 8% since pre-industrial times, presumably due to human activities; we are unable to specify the sources, but it is likely that agriculture plays a part.

3.0.6 The effect of ozone on climate is strongest in the upper troposphere and lower stratosphere. Model calculations indicate that ozone in the upper troposphere should have increased due to human-made emissions of nitrogen oxides, hydrocarbons and carbon monoxide. While at ground level ozone has increased in the northern hemisphere in response to these emissions, observations are insufficient to confirm the expected increase in the upper troposphere. The lack of adequate observations prevents us from accurately quantifying the climatic effect of changes in tropospheric ozone.

3.0.7 In the lower stratosphere at high southern latitudes ozone has decreased considerably due to the effects of CFCs,

Scientific Assessment of Climate Change

	Carbon Dioxide	Methane	CFC-11	CFC-12	Nitrous Oxide
Atmospheric concentration	ppmv	ppmv	pptv	pptv	ppbv
Pre-industrial (1750-1800)	280	0.8	0	0	288
Present day (1990)	353	1.72	280	484	310
Current rate of change per year	1.8 (0.5%)	0.015 (0.9%)	9.5 (4%)	17 (4%)	0.8 (0.25%)
Atmospheric lifetime (years)	(50-200)†	10	65	130	150





ppmv = parts per million by volume:

ppbv = parts per billion (thousand million) by volume,

pptv = parts per trillion (million million) by volume

† The way in which CO₂ is absorbed by the oceans and biosphere is not simple and a single value cannot be given; refer to the main report for further discussion

Summary of key greenhouse gases affected by human activities



Concentrations of carbon dioxide and methane, after remaining relatively constant up to the 18th century, have risen sharply since then due to man's activities. Concentrations of nitrous oxide have increased since the mid-18th century, especially in the last few decades. CFCs were not present in the atmosphere until the 1930s.

and there are indications of a global-scale decrease which, while not understood, may also be due to CFCs. These observed decreases should act to cool the earth's surface, thus providing a small offset to the predicted warming produced by the other greenhouse gases. Further reductions in lower stratospheric ozone are possible during the next few decades as the atmospheric abundances of CFCs continue to increase.


3.1 Concentrations, lifetimes and stabilization of the gases

3.1.1 In order to calculate the atmospheric concentrations of carbon dioxide which will result from human-made emissions, we use computer models which incorporate details of the emissions and which include representations of the transfer of carbon dioxide between the atmosphere, oceans and terrestrial biosphere. For the other greenhouse gases, models which incorporate the effects of chemical reactions in the atmosphere are employed.

3.1.2 The atmospheric lifetimes of the gases are determined by their sources and sinks in the oceans, atmosphere and biosphere. Carbon dioxide, chloro-fluorocarbons and nitrous oxide are removed only slowly from the atmosphere and hence, following a change in emissions, their atmospheric concentrations take decades to centuries to adjust fully. Even if all human-made emissions of carbon dioxide were halted in the year 1990, about half of the increase in carbon dioxide concentration caused by human activities would still be evident by the year 2100.

3.1.3 In contrast, some of the CFC substitutes and methane have relatively short atmospheric lifetimes so that their atmospheric concentrations respond fully to emission changes within a few decades.

3.1.4 To illustrate the emission-concentration relationship clearly, the effect of hypothetical changes in carbon dioxide fossil fuel emissions is shown below:



Stabilization of atmospheric concentrations.

Reductions in the human-made emissions of greenhouse gases required to stabilize concentrations at present day levels:


Carbon Dioxide	>60%
Methane	15—20%
Nitrous Oxide	70—80%
CFC-11	70—75%
CFC-12	75—85%
HCFC-22	40—50%

Note that the stabilization of each of these gases would have different effects on climate, as explained in the next section.

(a) continuing global emissions at 1990 levels; (b) halving of emissions in 1990; (c) reductions in emissions of 2% per year (pa) from 1990 and (d) a 2% pa increase from 1990-2010 followed by a 2% pa decrease from 2010.

3.1.5 Continuation of present day emissions is committing us to increased future concentrations, and the longer emissions continue to increase, the greater would reductions have to be to stabilize at a given level. If there are critical concentration levels that should not be exceeded, then the earlier emission reductions are made the more effective they are.

3.1.6 The term "atmospheric stabilization" is often used to describe the limiting of the concentration of the greenhouse gases at a certain level. The amount by which human-made emissions of a greenhouse gas must be reduced in order to stabilize at present day concentrations, for example, is shown in the table above. For most gases the reductions would have to be substantial.



The relationship between hypothetical fossil fuel emissions of carbon dioxide and its concentration in the atmosphere is shown in the case where (a) emissions continue at 1990 levels, (b) emissions are reduced by 50% in 1990 and continue at that level, (c) emissions are reduced by 2% pa from 1990, and (d) emissions, after increasing by 2% pa until 2020, are then reduced by 2% pa thereafter.

3.2 How will greenhouse gas abundances change in the future?

3.2.1 We need to know future greenhouse gas concentrations in order to estimate future climate change. As already mentioned, these concentrations depend upon the magnitude of human-made emissions and on how changes in climate and other environmental conditions may influence the biospheric processes that control the exchange of natural greenhouse gases, including carbon dioxide and methane, between the atmosphere, oceans and terrestrial biosphere—the greenhouse gas “feedbacks”.

3.2.2 Four scenarios of future human-made emissions were developed by Working Group III. The first of these assumes that few or no steps are taken to limit greenhouse gas emissions, and this is therefore termed Business-as-Usual (BaU). (It should be noted that an aggregation of national forecasts of emissions of carbon dioxide and methane to the year 2025 undertaken by Working Group III resulted in global emissions 10-20% higher than in the BaU scenario.) The other three scenarios assume that progressively increasing levels of controls reduce the growth of emissions; these are referred to as scenarios B, C, and D. They are briefly described in the Annex. Future concentrations of some of the greenhouse gases which would arise from these emissions are shown opposite.




3.3 Greenhouse gas feedbacks

3.3.1 Some of the possible feedbacks which could significantly modify future greenhouse gas concentrations in a warmer world are discussed in the following paragraphs.

3.3.2 The net emissions of carbon dioxide from terrestrial ecosystems will be elevated if higher temperatures increase respiration at a faster rate than photosynthesis, or if plant populations, particularly large forests, cannot adjust rapidly enough to changes in climate.

3.3.3 A net flux of carbon dioxide to the atmosphere may be particularly evident in warmer conditions in tundra and boreal regions where there are large stores of carbon. The opposite is true if higher abundances of carbon dioxide in the atmosphere enhance the productivity of natural ecosystems, or if there is an increase in soil moisture which can be expected to stimulate plant growth in dry ecosystems and to increase the storage of carbon in tundra peat. The extent to which ecosystems can sequester increasing atmospheric carbon dioxide remains to be quantified.


3.3.4 If the oceans become warmer, their net uptake of carbon dioxide may decrease because of changes in (i) the chemistry of carbon dioxide in seawater (ii) biological activity in surface waters and (iii) the rate of exchange of carbon dioxide between the surface layers and the deep ocean. This last depends upon the rate of formation of deep water in the ocean which, in the North Atlantic for example, might decrease if the salinity decreases as a result of a change in climate.




Atmospheric concentrations of carbon dioxide, methane and CFC-11 resulting from the four IPCC emissions scenarios

3.3.5 Methane emissions from natural wetlands and rice paddies are particularly sensitive to temperature and soil moisture. Emissions are significantly larger at higher temperatures and with increased soil moisture; conversely, a decrease in soil moisture would result in smaller emissions. Higher temperatures could increase the emissions of methane at high northern latitudes from decomposable organic matter trapped in permafrost and methane hydrates.

3.3.6 As illustrated earlier, ice core records show that methane and carbon dioxide concentrations changed in a similar sense to temperature between ice ages and interglacials.



Increase in radiative forcing since the mid-18th century, and predicted to result from the four IPCC emissions scenarios, also expressed as equivalent carbon dioxide concentrations



The contribution from each of the human-made greenhouse gases to the change in radiative forcing from 1980 to 1990. The contribution from ozone may also be significant, but cannot be quantified at present

3.3.7 Although many of these feedback processes are poorly understood, it seems likely that, overall, they will act to increase, rather than decrease, greenhouse gas concentrations in a warmer world.

4. Which gases are the most important?

4.0.1 We are certain that increased greenhouse gas concentrations increase radiative forcing. We can calculate the forcing with much more confidence than the climate change that results because the former avoids the need to evaluate a number of poorly understood atmospheric responses. We then have a base from which to calculate the relative effect on climate of an increase in concentration of each gas in the present-day atmosphere, both in absolute terms and relative to carbon dioxide. These relative effects span a wide range; methane is about 21 times more effective, molecule-for-molecule, than carbon dioxide, and CFC-11 about 12,000 times more effective. On a kilogram-per-kilogram basis, the equivalent values are 58 for methane and about 4,000 for CFC-11, both relative to carbon dioxide. Values for other greenhouse gases are to be found in the full report.

4.0.2 The total radiative forcing at any time is the sum of those from the individual greenhouse gases. We show in the figure below how this quantity has changed in the past (based on observations of greenhouse gases) and how it might change in the future (based on the four IPCC emissions scenarios). For simplicity, we can express total forcing in terms of the amount of carbon dioxide which would give that forcing; this is termed the equivalent carbon dioxide concentration. Greenhouse gases have increased since pre-industrial times (the mid-18th century) by an amount that is radiatively equivalent to about a 50% increase

GLOBAL WARMING POTENTIALS

The warming effect of an emission of 1 kg of each gas relative to that of CO₂
 These figures are best estimates calculated on the basis of the present day atmospheric composition

	Time Horizon		
	20 yr	100 yr	500 yr
Carbon dioxide	1	1	1
Methane (including indirect)	63	21	9
Nitrous oxide	270	290	190
CFC-11	4500	3500	1500
CFC-12	7100	7300	4500
HCFC-22	4100	1500	510

Global Warming Potentials for a range of CFCs and potential replacements are given in the full text

in carbon dioxide, although carbon dioxide itself has risen by only 26%; other gases have made up the rest.

4.0.3 The contributions of the various gases to the total increase in climate forcing during the 1980s is shown above as a pie diagram; carbon dioxide is responsible for about half the decadal increase. (Ozone, the effects of which may be significant, is not included)

4.1 How can we evaluate the effect of different greenhouse gases?

4.1.1 To evaluate possible policy options, it is useful to know the relative radiative effect (and, hence, potential climate effect) of equal emissions of each of the greenhouse gases. The concept of relative Global Warming Potentials (GWP) has been developed to take into account the differing times that gases remain in the atmosphere.

4.1.2 This index defines the time-integrated warming effect due to an instantaneous release of unit mass (1 kg) of a given greenhouse gas in today's atmosphere, relative to that of carbon dioxide. The relative importances will change in the future as atmospheric composition changes because, although radiative forcing increases in direct proportion to the concentration of CFCs, changes in the other greenhouse gases (particularly carbon dioxide) have an effect on forcing which is much less than proportional.

4.1.3 The GWPs in the following table are shown for three time horizons, reflecting the need to consider the cumulative effects on climate over various time scales. The longer time horizon is appropriate for the cumulative effect; the shorter timescale will indicate the response to emission changes in the short term. There are a number of practical difficulties in devising and calculating the values of the GWPs, and the values given here should be considered as

preliminary. In addition to these direct effects, there are indirect effects of human-made emissions arising from chemical reactions between the various constituents. The indirect effects on stratospheric water vapour, carbon dioxide and tropospheric ozone have been included in these estimates.

4.1.4 The table indicates, for example, that the effectiveness of methane in influencing climate will be greater in the first few decades after release, whereas emission of the longer-lived nitrous oxide will affect climate for a much longer time. The lifetimes of the proposed CFC replacements range from 1 to 40 years; the longer lived replacements are still potentially effective as agents of climate change. One example of this, HCFC-22 (with a 15-year lifetime), has a similar effect (when released in the same amount) as CFC-11 on a 20-year timescale; but less over a 500-year timescale.

4.1.5 The table shows carbon dioxide to be the least effective greenhouse gas per kilogramme emitted, but its contribution to global warming, which depends on the product of the GWP and the amount emitted, is largest. In the example in the box below, the effects over 100 years of emissions of greenhouse gases in 1990 are shown relative to carbon dioxide. This is illustrative; to compare the effect of different emission projections we have to sum the effect of emissions made in future years.

4.1.6 There are other technical criteria which may help policymakers to decide, in the event of emissions reductions being deemed necessary, which gases should be considered. Does the gas contribute in a major way to current, and future, climate forcing? Does it have a long lifetime, so earlier reductions in emissions would be more effective than those made later? And are its sources and sinks well enough known to decide which could be controlled in practice? The table **opposite** illustrates these factors.

The Relative Cumulative Climate Effect of 1990 Man-Made Emissions

	GWP (100yr horizon)	1990 emissions (Tg)	Relative contribution over 100yr
Carbon dioxide	1	26000†	61%
Methane*	21	300	15%
Nitrous oxide	290	6	4%
CFCs	Various	0.9	11%
HCFC-22	1500	0.1	0.5%
Others*	Various		8.5%

* These values include the indirect effect of these emissions on other greenhouse gases via chemical reactions in the atmosphere. Such estimates are highly model dependent and should be considered preliminary and subject to change. The estimated effect of ozone is included under "others". The gases included under "others" are given in the full report.

† 26 000 Tg (teragrams) of carbon dioxide = 7 000 Tg (=7 Gt) of carbon

Characteristics of Greenhouse Gases

GAS	MAJOR CONTRIBUTOR?	LONG LIFETIME?	SOURCES KNOWN?
Carbon dioxide	yes	yes	yes
Methane	yes	no	semi-quantitatively
Nitrous oxide	not at present	yes	qualitatively
CFCs	yes	yes	yes
HCFCs, etc	not at present	mainly no	yes
Ozone	possibly	no	qualitatively

5. How much do we expect climate to change?

5.0.1 It is relatively easy to determine the direct effect of the increased radiative forcing due to increases in greenhouse gases. However, as climate begins to warm, various processes act to amplify (through positive feedbacks) or reduce (through negative feedbacks) the warming. The main feedbacks which have been identified are due to changes in water vapour, sea-ice, clouds and the oceans.

5.0.2 The best tools we have which take the above feedbacks into account (but do not include greenhouse gas feedbacks) are three-dimensional mathematical models of the climate system (atmosphere-ocean-ice-land), known as General Circulation Models (GCMs). They synthesize our knowledge of the physical and dynamical processes in the overall system and allow for the complex interactions between the various components. However, in their current state of development, the descriptions of many of the processes involved are comparatively crude. Because of this, considerable uncertainty is attached to these predictions of climate change, which is reflected in the range of values given; further details are given in a later section.

5.0.3 The estimates of climate change presented here are based on

- i) the “best estimate” of equilibrium climate sensitivity (i.e the equilibrium temperature change due to a doubling of carbon dioxide in the atmosphere) obtained from model simulations, feedback analyses and observational considerations (see later box: “What tools do we use?”)
- ii) a “box diffusion upwelling” ocean-atmosphere climate model which translates the greenhouse forcing into the evolution of the temperature response for the prescribed climate sensitivity. (This simple model has been calibrated against more complex atmosphere-

ocean coupled GCMs for situations where the more complex models have been run).

5.1 How quickly will global climate change?

a. If emissions follow a Business-as-Usual pattern


5.1.1 Under the IPCC Business-as-Usual (Scenario A) emissions of greenhouse gases, the average rate of increase of global-mean temperature during the next century is estimated to be about 0.3°C per decade (with an uncertainty range of 0.2°C to 0.5°C). This will result in a likely increase in global-mean temperature of about 1°C above the present value (about 2°C above that in the pre-industrial period) by 2025 and 3°C above today’s (about 4°C above pre-industrial) before the end of the next century.

5.1.2 The projected temperature rise out to the year 2100, with high, low and best-estimate climate responses, is shown in the diagram below. Because of other factors which influence climate, we would not expect the rise to be a steady one.

5.1.3 The temperature rises shown above are realized temperatures; at any time we would also be committed to a further temperature rise toward the equilibrium temperature (see box: “Equilibrium and Realized Climate Change”). For the BaU “best estimate” case in the year 2030, for example, a further 0.9°C rise would be expected, about 0.2°C of which would be realized by 2050 (in addition to changes due to further greenhouse gas increases); the rest would become apparent in decades or centuries.

5.1.4 Even if we were able to stabilize emissions of each of the greenhouse gases at present day levels from now on, the temperature is predicted to rise by about 0.2°C per decade for the first few decades.

5.1.5 The global warming will also lead to increased



Simulation of the increase in global mean temperature from 1850-1990 due to observed increases in greenhouse gases, and predictions of the rise between 1990 and 2100 resulting from the Business-as-usual scenario.

global average precipitation and evaporation of a few percent by 2030. Areas of sea-ice and snow are expected to diminish.

b. If emissions are subject to controls

5.1.6 Under the other IPCC emission scenarios which assume progressively increasing levels of controls, average rates of increase in global-mean temperature over the next century are estimated to be about 0.2°C per decade (Scenario B), just above 0.1°C per decade (Scenario C) and about 0.1°C per decade (Scenario D). The results are illustrated below with the Business-as-usual case for comparison. Only the best-estimate of the temperature rise is shown in each case.


5.1.7 The indicated range of uncertainty in global temperature rise given above reflects a subjective assessment of uncertainties in the calculation of climate response, but does not include those due to the transformation of emissions to concentrations, nor the effects of greenhouse gas feedbacks.

5.2 What will be the patterns of climate change by 2030?

5.2.1 Knowledge of the global-mean warming and change in precipitation is of limited use in determining the impacts of climate change, for instance on agriculture. For this we need to know changes, regionally and seasonally.

5.2.2 Models predict that surface air will warm faster over land than over oceans, and a minimum of warming will occur around Antarctica and in the northern North Atlantic region.

5.2.3 There are some continental-scale changes which are consistently predicted by the highest resolution models and for which we understand the physical reasons. The warming is predicted to be 50-100% greater than the global-mean in high northern latitudes in winter, and substantially smaller than the global-mean in regions of sea ice in summer. Precipitation is predicted to increase on average in middle and high latitude continents in winter (by some 5—10% over 35-55°N).



Simulations of the increase in global mean temperature from 1850-1990 due to observed increases in greenhouse gases, and predictions of the rise between 1990 and 2100 resulting from the IPCC Scenario B, C and D emissions, with the Business-as-usual case for comparison.

What tools do we use to predict future climate, and how do we use them?

The most highly developed tool which we have to predict future climate is known as a general circulation model or GCM. Such models are based on the laws of physics and use descriptions in simplified physical terms (called parameterisations) of the smaller-scale processes such as those due to clouds and deep mixing in the ocean. In a climate model an atmospheric component, essentially the same as a weather prediction model, is coupled to a model of the ocean, which can be equally complex.

Climate forecasts are derived in a different way from weather forecasts. A weather prediction model gives a description of the atmosphere's state up to 10 days or so ahead, starting from a detailed description of an initial state of the atmosphere at a given time. Such forecasts describe the movement and development of large weather systems, though they cannot represent very small scale phenomena; for example, individual shower clouds.

To make a climate forecast, the climate model is first run for a few (simulated) decades. The statistics of the model's output is a description of the model's simulated climate which, if the model is a good one, will bear a close resemblance to the climate of the real atmosphere and ocean. The above exercise is then repeated with increasing concentrations of the greenhouse gases in the model. The differences between the statistics of the two simulations (for example in mean temperature and interannual variability) provide an estimate of the accompanying climate change.

The long-term change in surface air temperature following a doubling of carbon dioxide (referred to as the climate sensitivity) is generally used as a benchmark to compare models. The range of results from model studies is 1.9 to 5.2°C. Most results are close to 4.0°C but recent studies using a more detailed but not necessarily more accurate representation of cloud processes give results in the lower half of this range. Hence the models results do not justify altering the previously accepted range of 1.5 to 4.5°C.

Although scientists are reluctant to give a single best estimate in this range, it is necessary for the presentation of climate predictions for a choice of best estimate to be made. Taking into account the model results, together with observational evidence over the last century which is suggestive of the climate sensitivity being in the lower half of the range, (see section: "Has man already begun to change global climate?") a value of climate sensitivity of 2.5°C has been chosen as the best estimate.

In this Assessment, we have also used much simpler models, which simulate the behaviour of GCMs, to make predictions of the evolution with time of global temperature from a number of emission scenarios. These so-called box-diffusion models contain highly simplified physics but give similar results to GCMs when globally averaged.

A completely different, and potentially useful, way of predicting patterns of future climate is to search for periods in the past when the global mean temperatures were similar to those we expect in future, and then use the past spatial patterns as analogues of those which will arise in the future. For a good analogue, it is also necessary for the forcing factors (for example, greenhouse gases, orbital variations) and other conditions (for example, ice cover, topography, etc.) to be similar: direct comparisons with climate situations for which these conditions do not apply cannot be easily interpreted. Analogues of future greenhouse-gas-changed climates have not been found.

We cannot therefore advocate the use of palaeo-climates as predictions of regional climate change due to future increases in greenhouse gases. However, palaeo-climatological information can provide useful insights into climate processes, and can assist in the validation of climate models.

5.2.4 Five regions, each a few million square kilometres in area and representative of different climatological regimes, were selected by IPCC for particular study (see map below). In the box below are given the changes in temperature, precipitation and soil moisture, which are predicted to occur by 2030 on the Business-as-Usual scenario, as an average over each of the five regions. There may be considerable variations within the regions. In general, confidence in these regional estimates is low, especially for the changes in precipitation and soil moisture, but they are examples of our best estimates. We cannot yet

give reliable regional predictions at the smaller scales demanded for impacts assessments.

5.3 How will climate extremes and extreme events change?


5.3.1 Changes in the variability of weather and the frequency of extremes will generally have more impact than changes in the mean climate at a particular location. With the possible exception of an increase in the number of intense showers, there is no clear evidence that weather

Equilibrium and realized climate change

When the radiative forcing on the earth-atmosphere system is changed, for example by increasing greenhouse gas concentrations, the atmosphere will try to respond (by warming) immediately. But the atmosphere is closely coupled to the oceans, so in order for the air to be warmed by the greenhouse effect, the oceans also have to be warmed; because of their thermal capacity this takes decades or centuries. This exchange of heat between atmosphere and ocean will act to slow down the temperature rise forced by the greenhouse effect.

In a hypothetical example where the concentration of greenhouse gases in the atmosphere, following a period of constancy, rises suddenly to a new level and remains there, the radiative forcing would also rise rapidly to a new level. This increased radiative forcing would cause the atmosphere and oceans to warm, and eventually come to a new, stable, temperature. A commitment to this equilibrium temperature rise is incurred as soon as the greenhouse gas concentration changes. But at any time before equilibrium is reached, the actual temperature will have risen by only part of the equilibrium temperature change, known as the realized temperature change.

Models predict that, for the present day case of an increase in radiative forcing which is approximately steady, the realized temperature rise at any time is about 50% of the committed temperature rise if the climate sensitivity (the response to a doubling of carbon dioxide) is 4.5°C and about 80% if the climate sensitivity is 1.5°C. If the forcing were then held constant, temperatures would continue to rise slowly, but it is not certain whether it would take decades or centuries for most of the remaining rise to equilibrium to occur.



variability will change in the future. In the case of temperatures, assuming no change in variability, but with a modest increase in the mean, the number of days with temperatures above a given value at the high end of the distribution will increase substantially. On the same assumptions, there will be a decrease in days with temperatures at the low end of the distribution. So the number of very hot days or frosty nights can be substantially changed without any change in the variability of the weather. The number of days with a minimum threshold amount of soil moisture (for viability of a certain crop, for example) would be even more sensitive to changes in average precipitation and evaporation.

5.3.2 If the large scale weather regimes, for instance depression tracks or anticyclones, shift their position, this would effect the variability and extremes of weather at a particular location, and could have a major effect. However, we do not know if, or in what way, this will happen.

5.4 Will storms increase in a warmer world?

5.4.1 Storms can have a major impact on society. Will their frequency, intensity or location increase in a warmer world?

5.4.2 Tropical storms, such as typhoons and hurricanes, only develop at present over seas that are warmer than about

Estimates for changes by 2030

(IPCC Business-as-Usual scenario; changes from pre-industrial)

The numbers given below are based on high resolution models, scaled to be consistent with our best estimate of global mean warming of 1.8°C by 2030. For values consistent with other estimates of global temperature rise, the numbers below should be reduced by 30% for the low estimate or increased by 50% for the high estimate. Precipitation estimates are also scaled in a similar way.

Confidence in these regional estimates is low

Central North America (35°—50°N 85°—105°W)

The warming varies from 2 to 4°C in winter and 2 to 3°C in summer. Precipitation increases range from 0 to 15% in winter whereas there are decreases of 5 to 10% in summer. Soil moisture decreases in summer by 15 to 20%.

Southern Asia (5°—30°N 70°—105°E)

The warming varies from 1 to 2°C throughout the year. Precipitation changes little in winter and generally increases throughout the region by 5 to 15% in summer. Summer soil moisture increases by 5 to 10%.

Sahel (10°—20°N 20°W—40°E)


The warming ranges from 1 to 3°C. Area mean precipitation increases and area mean soil moisture decreases marginally in summer. However, throughout the region, there are areas of both increase and decrease in both parameters.

Southern Europe (35°—50°N 10°W—45°E)

The warming is about 2°C in winter and varies from 2 to 3°C in summer. There is some indication of increased precipitation in winter, but summer precipitation decreases by 5 to 15%, and summer soil moisture by 15 to 25%.

Australia (12°—45°S 110°—115°E)

The warming ranges from 1 to 2°C in summer and is about 2°C in winter. Summer precipitation increases by around 10%, but the models do not produce consistent estimates of the changes in soil moisture. The area averages hide large variations at the sub-continental level.



Map showing the locations and extents of the five areas selected by IPCC

26°C. Although the area of sea having temperatures over this critical value will increase as the globe warms, the critical temperature itself may increase in a warmer world. Although the theoretical maximum intensity is expected to increase with temperature, climate models give no consistent indication whether tropical storms will increase or decrease in frequency or intensity as climate changes; neither is there any evidence that this has occurred over the past few decades.

5.4.3 Mid-latitude storms, such as those which track across the North Atlantic and North Pacific, are driven by the equator-to-pole temperature contrast. As this contrast will probably be weakened in a warmer world (at least in the northern hemisphere), it might be argued that mid-latitude storms will also weaken or change their tracks, and there is some indication of a general reduction in day-to-day variability in the mid-latitude storm tracks in winter in model simulations, though the pattern of changes vary from model to model. Present models do not resolve smaller-scale disturbances, so it will not be possible to assess changes in storminess until results from higher resolution models become available in the next few years.

5.5 Climate change in the longer term

5.5.1 The foregoing calculations have focussed on the period up to the year 2100; it is clearly more difficult to make calculations for years beyond 2100. However, while the timing of a predicted increase in global temperatures has substantial uncertainties, the prediction that an increase will eventually occur is more certain. Furthermore, some model calculations that have been extended beyond 100 years suggest that, with continued increases in greenhouse climate forcing, there could be significant changes in the ocean circulation, including a decrease in North Atlantic deep water formation.

5.6 Other factors which could influence future climate

5.6.1 Variations in the output of solar energy may also affect climate. On a decadal timescale solar variability and changes in greenhouse gas concentration could give changes of similar magnitudes. However the variation in solar intensity changes sign so that over longer timescales the increases in greenhouse gases are likely to be more important. Aerosols as a result of volcanic eruptions can lead to a cooling at the surface which may oppose the greenhouse warming for a few years following an eruption. Again, over longer periods the greenhouse warming is likely to dominate.

5.6.2 Human activity is leading to an increase in aerosols in the lower atmosphere, mainly from sulphur emissions. These have two effects, both of which are difficult to quantify but which may be significant particularly at the regional level. The first is the direct effect of the aerosols on the radiation scattered and absorbed by the atmosphere. The

second is an indirect effect whereby the aerosols affect the microphysics of clouds leading to an increased cloud reflectivity. Both these effects might lead to a significant regional cooling; a decrease in emissions of sulphur might be expected to increase global temperatures.

5.6.3 Because of long-period couplings between different components of the climate system, for example between ocean and atmosphere, the earth's climate would still vary without being perturbed by any external influences. This natural variability could act to add to, or subtract from, any human-made warming; on a century timescale this would be less than changes expected from greenhouse gas increases.

6. How much confidence do we have in our predictions?

6.0.1 Uncertainties in the above climate predictions arise from our imperfect knowledge of:

- future rates of human-made emissions
- how these will change the atmospheric concentrations of greenhouse gases
- the response of climate to these changed concentrations.

6.0.2 Firstly, it is obvious that the extent to which climate will change depends on the rate at which greenhouse gases (and other gases which affect their concentrations) are emitted. This in turn will be determined by various complex economic and sociological factors. Scenarios of future emissions were generated within IPCC WG III and are described in the annex.

6.0.3 Secondly, because we do not fully understand the sources and sinks of the greenhouse gases, there are uncertainties in our calculations of future concentrations arising from a given emissions scenario. We have used a number of models to calculate concentrations and chosen a best estimate for each gas. In the case of carbon dioxide, for example, the concentration increase between 1990 and 2070 due to the Business-as-Usual emissions scenario spanned almost a factor of two between the highest and lowest model result (corresponding to a range in radiative forcing change of about 50%)

6.0.4 Furthermore, because natural sources and sinks of greenhouse gases are sensitive to a change in climate, they may substantially modify future concentrations (see earlier section: "Greenhouse gas feedbacks"). It appears that, as climate warms, these feedbacks will lead to an overall increase, rather than decrease, in natural greenhouse gas abundances. For this reason, climate change is likely to be greater than the estimates we have given.

Confidence in predictions from climate models

What confidence can we have that climate change due to increasing greenhouse gases will look anything like the model predictions? Weather forecasts can be compared with the actual weather the next day and their skill assessed; we cannot do that with climate predictions. However, there are several indicators that give us some confidence in the predictions from climate models.

When the latest atmospheric models are run with the present atmospheric concentrations of greenhouse gases and observed boundary conditions their simulation of present climate is generally realistic on large scales, capturing the major features such as the wet tropical convergence zones and mid-latitude depression belts, as well as the contrasts between summer and winter circulations. The models also simulate the observed variability: for example, the large day-to-day pressure variations in the middle latitude depression belts and the maxima in interannual variability responsible for the very different character of one winter from another both being represented. However, on regional scales (2,000 km or less), there are significant errors in all models.

Overall confidence is increased by atmospheric models' generally satisfactory portrayal of aspects of variability of the atmosphere, for instance those associated with variations in sea surface temperature. There has been some success in simulating the general circulation of the ocean, including the patterns (though not always the intensities) of the principal currents, and the distributions of tracers added to the ocean.

Atmospheric models have been coupled with simple models of the ocean to predict the equilibrium response to greenhouse gases, under the assumption that the model errors are the same in a changed climate. The ability of such models to simulate important aspects of the climate of the last ice age generates confidence in their usefulness. Atmospheric models have also been coupled with multilayer ocean models (to give coupled ocean-atmosphere GCMs) which predict the gradual response to increasing greenhouse gases. Although the models so far are of relatively coarse resolution, the large scale structures of the ocean and the atmosphere can be simulated with some skill. However, the coupling of ocean and atmosphere models reveals a strong sensitivity to small scale errors which leads to a drift away from the observed climate. As yet, these errors must be removed by adjustments to the exchange of heat between ocean and atmosphere. There are similarities between results from the coupled models using simple representations of the ocean and those using more sophisticated descriptions, and our understanding of such differences as do occur gives us some confidence in the results.

6.0.5 Thirdly, climate models are only as good as our understanding of the processes which they describe, and this is far from perfect. The ranges in the climate predictions given above reflect the uncertainties due to model imperfections: the largest of these is cloud feedback (those factors affecting the cloud amount and distribution and the interaction of clouds with solar and terrestrial radiation), which leads to a factor of two uncertainty in the size of the warming. Others arise from the transfer of energy between the atmosphere and ocean, the atmosphere and land surfaces, and between the upper and deep layers of the ocean. The treatment of sea-ice and convection in the models is also crude. Nevertheless, for reasons given in the box below, we have substantial confidence that models can predict at least the broad-scale features of climate change.

6.0.6 Furthermore, we must recognize that our imperfect understanding of climate processes (and corresponding ability to model them) could make us vulnerable to surprises: just as the human-made ozone hole over Antarctica was entirely unpredicted. In particular, the ocean circulation, changes in which are thought to have led to periods of comparatively rapid climate change at the end of the last ice age, is not well observed, understood or modelled.

7. Will the climate of the future be very different?

7.0.1 When considering future climate change, it is clearly essential to look at the record of climate variation in the past. From it we can learn about the range of natural climate variability, to see how it compares with what we expect in the future, and also look for evidence of recent climate change due to man's activities.

7.0.2 Climate varies naturally on all time scales from hundreds of millions of years down to the year to year. Prominent in the Earth's history have been the 100,000 year glacial-interglacial cycles when climate was mostly cooler than at present. Global surface temperatures have typically varied by 5-7°C through these cycles, with large changes in ice volume and sea level, and temperature changes as great as 10-15°C in some middle and high latitude regions of the northern hemisphere. Since the end of the last ice age, about 10,000 years ago, global surface temperatures have probably fluctuated by little more than 1°C. Some fluctuations have lasted several centuries, including the Little Ice Age which ended in the nineteenth century and which appears to have been global in extent.

Scientific Assessment of Climate Change

7.0.3 The changes predicted to occur by about the middle of the next century due to increases in greenhouse gas concentrations from the Business-as-Usual emissions will make global-mean temperatures higher than they have been in the last 150,000 years.

7.0.4 The rate of change of global temperatures predicted for Business-as-Usual emissions will be greater than those which have occurred naturally on earth over the last 10,000 years, and the rise in sea level will be about three to six times faster than that seen over the last 100 years or so.

8. Has man already begun to change the global climate?

8.0.1 The instrumental record of surface temperature is fragmentary until the mid-nineteenth century, after which it slowly improves. Because of different methods of measurement, historical records have to be harmonized with modern observations, introducing some uncertainty. Despite these problems we believe that a real warming of the globe of 0.3°C–0.6°C has taken place over the last century; any bias due to urbanization is likely to be less than 0.05°C.

8.0.2 Moreover since 1900 similar temperature increases are seen in three independent data sets: one collected over land and two over the oceans. The figure below shows current estimates of smoothed global mean surface temperature over land and ocean since 1860. Confidence in the records has been increased by their similarity to recent satellite measurements of mid-tropospheric temperatures.


8.0.3 Although the overall temperature rise has been broadly similar in both hemispheres, it has not been steady, and differences in their rates of warming have sometimes persisted for decades. Much of the warming since 1900 has been concentrated in two periods, the first between about 1910 and 1940 and the other since 1975: the five warmest

years on record have all been in the 1980s. The northern hemisphere cooled between the 1940s and the early 1970s when southern hemisphere temperatures stayed nearly constant. The pattern of global warming since 1975 has been uneven with some regions, mainly in the northern hemisphere, continuing to cool until recently. This regional diversity indicates that future regional temperature changes are likely to differ considerably from a global average.

8.0.4 The conclusion that global temperature has been rising is strongly supported by the retreat of most mountain glaciers of the world since the end of the nineteenth century and the fact that global sea level has risen over the same period by an average of 1 to 2 mm per year. Estimates of thermal expansion of the oceans, and of increased melting of mountain glaciers and the ice margin in West Greenland over the last century, show that the major part of the sea-level rise appears to be related to the observed global warming. This apparent connection between observed sea-level rise and global warming provides grounds for believing that future warming will lead to an acceleration in sea-level rise.

8.0.5 The size of the warming over the last century is broadly consistent with the predictions of climate models, but is also of the same magnitude as natural climate variability. If the sole cause of the observed warming were the human-made greenhouse effect, then the implied climate sensitivity would be near the lower end of the range inferred from the models. The observed increase could be largely due to natural variability; alternatively this variability and other man-made factors could have offset a still larger man-made greenhouse warming. The unequivocal detection of the enhanced greenhouse effect from observations is not likely for a decade or more, when the commitment to future climate change will then be considerably larger than it is today.

8.0.6 Global mean temperature alone is an inadequate indicator of greenhouse-gas-induced climatic change.



Global mean combined land and sea-surface temperatures, 1861–1989, relative to the average for 1951–1980


Identifying the causes of any global mean temperature change requires examination of other aspects of the changing climate, particularly its spatial and temporal characteristics—the man-made climate change “signal”. Patterns of climate change from models such as the northern hemisphere warming faster than the southern hemisphere, and surface air warming faster over land than over oceans, are not apparent in observations to date. However, we do not yet know what the detailed “signal” looks like because we have limited confidence in our predictions of climate change patterns. Furthermore, any changes to date could be masked by natural variability and other (possibly man-made) factors, and we do not have a clear picture of these.

9. How much will sea level rise ?


9.0.1 Simple models were used to calculate the rise in sea level to the year 2100; the results are illustrated below. The calculations necessarily ignore any long-term changes, unrelated to greenhouse forcing, that may be occurring but cannot be detected from the present data on land ice and the ocean. The sea-level rise expected from 1990-2100 under the IPCC Business-as-Usual emissions scenario is shown below. An average rate of global mean sea-level rise of about 6 cm per decade over the next century (with an uncertainty range of 3—10 cm per decade). The predicted rise is about 20 cm in global mean sea level by 2030, and 65 cm by the end of the next century. There will be significant regional variations.

9.0.2 The best estimate in each case is made up mainly of positive contributions from thermal expansion of the oceans and the melting of glaciers. Although, over the next 100 years, the effect of the Antarctic and Greenland ice sheets is expected to be small, they make a major contribution to the uncertainty in predictions.


9.0.3 Even if greenhouse forcing increased no further, there would still be a commitment to a continuing sea-level



Commitment to sea level rise in the year 2030. The curve shows the sea level rise due to Business-As-Usual emissions to 2030, with the additional rise that would occur in the remainder of the century even if climate forcing was stabilised in 2030.



Model estimates of sea-level rise from 1990—2100 due to all four emissions scenarios.



Sea level rise predicted to result from Business-as-Usual emissions, showing the best-estimate and range

rise for many decades and even centuries, due to delays in climate, ocean and ice mass responses. As an illustration, if the increases in greenhouse gas concentrations were to suddenly stop in 2030, sea level would go on rising from 2030 to 2100, by as much again as from 1990-2030, as shown in the diagram on p.35.

9.0.4 Predicted sea level rises due to the other three emissions scenarios are also shown, with the Business-as-Usual case for comparison: only best-estimate calculations are shown.

9.0.5 The West Antarctic Ice Sheet is of special concern. A large portion of it, containing an amount of ice equivalent to about 5 m of global sea level, is grounded far below sea level. There have been suggestions that a sudden outflow of ice might result from global warming and raise sea level quickly and substantially. Recent studies have shown that individual ice streams are changing rapidly on a decade-to-century timescale; however this is not necessarily related to climate change. Within the next century, it is not likely that there will be a major outflow of ice from West Antarctica due directly to global warming.

9.0.6 Any rise in sea level is not expected to be uniform over the globe. Thermal expansion, changes in ocean circulation, and surface air pressure will vary from region to region as the world warms, but in an as yet unknown way. Such regional details await further development of more realistic coupled ocean-atmosphere models. In addition,

vertical land movements can be as large or even larger than changes in global mean sea level; these movements have to be taken into account when predicting local change in sea level relative to land.

9.0.7 The most severe effects of sea-level rise are likely to result from extreme events (for example, storm surges) the incidence of which may be affected by climatic change.

10. What will be the effect of climate change on ecosystems?

10.0.1 Ecosystem processes such as photosynthesis and respiration are dependent on climatic factors and carbon dioxide concentration in the short term. In the longer term, climate and carbon dioxide are among the factors which control ecosystem structure, i.e., species composition, either directly by increasing mortality in poorly adapted species, or indirectly by mediating the competition between species. Ecosystems will respond to local changes in temperature (including its rate of change), precipitation, soil moisture and extreme events. Current models are unable to make reliable estimates of changes in these parameters on the required local scales.

10.0.2 Photosynthesis captures atmospheric carbon dioxide, water and solar energy and stores them in organic compounds which are then used for subsequent plant growth, the growth of animals or the growth of microbes in

Deforestation and reforestation

Man has been deforesting the Earth for millennia. Until the early part of the century, this was mainly in temperate regions, more recently it has been concentrated in the tropics. Deforestation has several potential impacts on climate: through the carbon and nitrogen cycles (where it can lead to changes in atmospheric carbon dioxide concentrations), through the change in reflectivity of terrain when forests are cleared, through its effect on the hydrological cycle (precipitation, evaporation and runoff) and surface roughness and thus atmospheric circulation which can produce remote effects on climate.

It is estimated that each year about 2 Gt of carbon (GtC) is released to the atmosphere due to tropical deforestation. The rate of forest clearing is difficult to estimate; probably until the mid-20th century, temperate deforestation and the loss of organic matter from soils was a more important contributor to atmospheric carbon dioxide than was the burning of fossil fuels. Since then, fossil fuels have become dominant: one estimate is that around 1980, 1.6 GtC was being released annually from the clearing of tropical forests, compared with about 5 GtC from the burning of fossil fuels. If all the tropical forests were removed, the input is variously estimated at from 150 to 240 GtC: this would increase atmospheric carbon dioxide by 35 to 60 ppmv.

To analyse the effect of reforestation we assume that 10 million hectares of forests are planted each year for a period of 40 years, i.e., 4 million km² would then have been planted by 2030, at which time 1 GtC would be absorbed annually until these forests reach maturity. This would happen in 40—100 years for most forests. The above scenario implies an accumulated uptake of about 20GtC by the year 2030 and up to 80GtC after 100 years. This accumulation of carbon in forests is equivalent to some 5-10% of the emission due to fossil fuel burning in the Business-as-Usual scenario.

Deforestation can also alter climate directly by increasing reflectivity and decreasing evapotranspiration. Experiments with climate models predict that replacing all the forests of the Amazon Basin by grassland would reduce the rainfall over the basin by about 20%, and increase mean temperature by several degrees.

the soil. All of these organisms release carbon dioxide via respiration into the atmosphere. Most land plants have a system of photosynthesis which will respond positively to increased atmospheric carbon dioxide ("the carbon dioxide fertilization effect") but the response varies with species. The effect may decrease with time when restricted by other ecological limitations, for example, nutrient availability. It should be emphasized that the carbon content of the terrestrial biosphere will increase only if the forest ecosystems in a state of maturity will be able to store more carbon in a warmer climate and at higher concentrations of carbon dioxide. We do not yet know if this is the case.

10.0.3 The response to increased carbon dioxide results in greater efficiencies of water, light and nitrogen use. These increased efficiencies may be particularly important during drought and in arid/semi-arid and infertile areas.

10.0.4 Because species respond differently to climatic change, some will increase in abundance and/or range while others will decrease. Ecosystems will therefore change in structure and composition. Some species may be displaced to higher latitudes and altitudes, and may be more prone to local, and possibly even global, extinction; other species may thrive.

10.0.5 As stated above, ecosystem structure and species distribution are particularly sensitive to the rate of change of climate. We can deduce something about how quickly global temperature has changed in the past from paleoclimatological records. As an example, at the end of the last glaciation, within about a century, temperature increased by up to 5°C in the North Atlantic region, mainly in Western Europe. Although during the increase from the glacial to the current interglacial temperature simple tundra ecosystems responded positively, a similar rapid temperature increase applied to more developed ecosystems could result in their instability.

11. What should be done to reduce uncertainties, and how long will this take?

11.0.1 Although we can say that some climate change is unavoidable, much uncertainty exists in the prediction of global climate properties such as the temperature and rainfall. Even greater uncertainty exists in predictions of regional climate change, and the subsequent consequences for sea level and ecosystems. The key areas of scientific uncertainty are:

- *clouds*: primarily cloud formation, dissipation, and radiative properties, which influence the response of the atmosphere to greenhouse forcing;
- *oceans*: the exchange of energy between the ocean and the atmosphere, between the upper layers of the

ocean and the deep ocean, and transport within the ocean, all of which control the rate of global climate change and the patterns of regional change;

- *greenhouse gases*: quantification of the uptake and release of the greenhouse gases, their chemical reactions in the atmosphere, and how these may be influenced by climate change.
- *polar ice sheets*: which affect predictions of sea level rise

11.0.2 Studies of land surface hydrology, and of impact on ecosystems, are also important.

11.0.3 To reduce the current scientific uncertainties in each of these areas will require internationally coordinated research, the goal of which is to improve our capability to observe, model and understand the global climate system. Such a programme of research will reduce the scientific uncertainties and assist in the formulation of sound national and international response strategies.

11.0.4 Systematic long-term observations of the system are of vital importance for understanding the natural variability of the Earth's climate system, detecting whether man's activities are changing it, parametrizing key processes for models, and verifying model simulations. Increased accuracy and coverage in many observations are required. Associated with expanded observations is the need to develop appropriate comprehensive global information bases for the rapid and efficient dissemination and utilization of data. The main observational requirements are:

- i) the maintenance and improvement of observations (such as those from satellites) provided by the World Weather Watch Programme of WMO;
- ii) the maintenance and enhancement of a programme of monitoring, both from satellite-based and surface-based instruments, of key climate elements for which accurate observations on a continuous basis are required, such as the distribution of important atmospheric constituents, clouds, the Earth's radiation budget, precipitation, winds, sea surface temperatures and terrestrial ecosystem extent, type and productivity;
- iii) the establishment of a global ocean observing system to measure changes in such variables as ocean surface topography, circulation, transport of heat and chemicals, and sea-ice extent and thickness;
- iv) the development of major new systems to obtain data on the oceans, atmosphere and terrestrial ecosystems using both satellite-based instruments and instruments based on the surface, on automated instrumented vehicles in the ocean, on floating and deep sea buoys, and on aircraft and balloons;

Scientific Assessment of Climate Change

v) the use of paleoclimatological and historical instrumental records to document natural variability and changes in the climate system, and subsequent environmental response.

11.0.5 The modelling of climate change requires the development of global models which couple together atmosphere, land, ocean and ice models and which incorporate more realistic formulations of the relevant processes and the interactions between the different components. Processes in the biosphere (both on land and in the ocean) also need to be included. Higher spatial resolution than is currently generally used is required if regional patterns are to be predicted. These models will require the largest computers which are planned to be available during the next decades.

11.0.6 Understanding of the climate system will be developed from analyses of observations and of the results from model simulations. In addition, detailed studies of particular processes will be required through targeted observational campaigns. Examples of such field campaigns include combined observational and small scale modelling studies for different regions, of the formation, dissipation, radiative, dynamical and microphysical properties of clouds, and ground-based (ocean and land) and aircraft measurements of the fluxes of greenhouse gases from specific ecosystems. In particular, emphasis must be placed on field experiments that will assist in the development and improvement of sub-grid-scale parametrizations for models.

11.0.7 The required programme of research will require unprecedented international cooperation, with the World Climate Research Programme (WCRP) of the World Meteorological Organization and International Council of Scientific Unions (ICSU), and the International Geosphere-Biosphere Programme (IGBP) of ICSU both playing vital roles. These are large and complex endeavours that will require the involvement of all nations, particularly the developing countries. Implementation of existing and planned projects will require increased financial and human resources; the latter requirement has immediate implications at all levels of education, and the international community of scientists needs to be widened to include more members from developing countries.

11.0.8 The WCRP and IGBP have a number of ongoing or planned research programs, that address each of the three key areas of scientific uncertainty. Examples include:

- **clouds:**
International Satellite Cloud Climatology Project (ISCCP);
Global Energy and Water Cycle Experiment (GEWEX).
- **oceans:**
World Ocean Circulation Experiment (WOCE);
Tropical Oceans and Global Atmosphere (TOGA).

- **trace gases:**
Joint Global Ocean Flux Study (JGOFS);
International Global Atmospheric Chemistry (IGAC)
Past Global Changes (PAGES).

11.0.9 As research advances, increased understanding and improved observations will lead to progressively more reliable climate predictions. However considering the complex nature of the problem and the scale of the scientific programmes to be undertaken we know that rapid results cannot be expected. Indeed further scientific advances may expose unforeseen problems and areas of ignorance.

11.0.10 Timescales for narrowing the uncertainties will be dictated by progress over the next 10—15 years in two main areas:

- Use of the fastest possible computers, to take into account coupling of the atmosphere and the oceans in models, and to provide sufficient resolution for regional predictions.
- Development of improved representation of small scale processes within climate models, as a result of the analysis of data from observational programmes to be conducted on a continuing basis well into the next century.

Annex

EMISSIONS SCENARIOS FROM WORKING GROUP III OF THE INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE

The Steering Group of the Response Strategies Working Group requested the USA and the Netherlands to develop emissions scenarios for evaluation by the IPCC Working Group I. The scenarios cover the emissions of carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), chlorofluorocarbons (CFCs), carbon monoxide (CO) and nitrogen oxides (NO_x) from the present up to the year 2100. Growth of the economy and population was taken common for all scenarios. Population was assumed to approach 10.5 billion in the second half of the next century. Economic growth was assumed to be 2—3% annually in the coming decade in the OECD countries and 3—5 % in the Eastern European and developing countries. The economic growth levels were assumed to decrease thereafter. In order to reach the required targets, levels of technological development and environmental controls were varied.



In the *Business-as-Usual scenario* (Scenario A) the energy supply is coal intensive and on the demand side only modest efficiency increases are achieved. Carbon monoxide controls are modest, deforestation continues until the tropical forests are depleted and agricultural emissions of methane and nitrous oxide are uncontrolled. For CFCs the Montreal Protocol is implemented albeit with only partial participation. Note that the aggregation of national projections by IPCC Working Group III gives higher emissions (10—20%) of carbon dioxide and methane by 2025.

In *Scenario B* the energy supply mix shifts towards lower carbon fuels, notably natural gas. Large efficiency increases

are achieved. Carbon monoxide controls are stringent, deforestation is reversed and the Montreal Protocol implemented with full participation.

In *Scenario C* a shift towards renewables and nuclear energy takes place in the second half of next century. CFCs are now phased out and agricultural emissions limited.

For *Scenario D* a shift to renewables and nuclear in the first half of the next century reduces the emissions of carbon dioxide, initially more or less stabilizing emissions in the industrialized countries. The scenario shows that stringent controls in industrialized countries combined with moderated growth of emissions in developing countries could stabilize atmospheric concentrations. Carbon dioxide emissions are reduced to 50% of 1985 levels by the middle of the next century.



Emissions of carbon dioxide and methane (as examples) to the year 2100, in the four scenarios developed by IPCC Working Group III.

