

Chapter 5. Coastal Systems and Low-Lying Areas**Coordinating Lead Authors**

Iñigo J. Losada (Spain), Poh Poh Wong (Singapore)

Lead Authors

Jean-Pierre Gattuso (France), Jochen Hinkel (Germany), Abdellatif Khattabi (Morocco), Kathleen McInnes (Australia), Yoshiki Saito (Japan), Asbury Sallenger (USA), Anond Snidvongs (Thailand)

Contributing Authors

So-Min Cheong (Republic of Korea), Kirstin Dow (USA), Carlos Duarte (Spain), Kris Ebi (USA), Jack Middelburg (The Netherlands), Susanne Moser (USA), Marcel Stive (The Netherlands), Richard Tol (The Netherlands), Athanasios Vafeidis (Greece)

Review Editors

Robert Nicholls (UK), Filipe Santos (Portugal)

Volunteer Chapter Scientist

Sara Amez (Spain)

Contents

Executive Summary

5.1. Introduction

5.2. Coastal Systems

5.2.1. Definitions

5.2.2. Climatic and Non-Climatic Drivers and Variability

5.2.2.1. Climatic Drivers

5.2.2.2. Non-Climate Drivers

5.3. Observed Impacts

5.3.1. Impacts on Coastal Habitats and Ecosystems

5.3.1.1. Rocky Shores

5.3.1.2. Beaches and Sand Dunes

5.3.1.3. Estuaries, Tidal Flats, and Lagoons

5.3.1.4. Deltas

5.3.1.5. Mangroves and Salt Marshes

5.3.1.6. Coral Reefs

5.3.1.7. Submerged Vegetation

5.3.2. Impacts on Human Systems

5.3.2.1. Human Settlements

5.3.2.2. Industry, Transport, and Infrastructures

5.3.2.3. Fisheries, Aquaculture, and Agriculture

5.3.2.4. Coastal Tourism and Recreation

5.3.2.5. Water Resources

5.3.2.6. Health

5.4. Projected Impacts

5.4.1. Impacts on Habitats and Ecosystems

5.4.1.1. Rocky Shores

5.4.1.2. Beaches and Sand Dunes

- 1 5.4.1.3. Estuaries, Tidal Flats, and Lagoons
- 2 5.4.1.4. Deltas
- 3 5.4.1.5. Mangroves and Salt Marshes
- 4 5.4.1.6. Coral Reefs
- 5 5.4.1.7. Seagrasses and Algae
- 6 5.4.2. Impacts on Human Systems
- 7 5.4.2.1. Human Settlements
- 8 5.4.2.2. Industry, Transport, and Infrastructures
- 9 5.4.2.3. Fisheries, Aquaculture, and Agriculture
- 10 5.4.2.4. Coastal Tourism and Recreation
- 11 5.4.2.5. Water Resources
- 12 5.4.2.6. Health
- 13
- 14 5.5. Assessing Vulnerabilities, Risks, and Costs
- 15 5.5.1. Approaches
- 16 5.5.2. Coastal Systems
- 17 5.5.2.1. Rocky Shores
- 18 5.5.2.2. Beaches and Sand Dunes
- 19 5.5.2.3. Estuaries
- 20 5.5.2.4. Temperate Lagoons
- 21 5.5.2.5. Salt Marshes
- 22 5.5.2.6. Mangroves
- 23 5.5.2.7. Seagrass Meadows
- 24 5.5.3. Human Activities
- 25 5.5.4. Costs
- 26 5.5.5. Uncertainties and the Long-Term Commitment to Sea-Level Rise
- 27
- 28 5.6. Adaptation and Managing Risks
- 29 5.6.1. Approaches
- 30 5.6.2. Practices
- 31 5.6.3. Adaptation Costs
- 32 5.6.4. Constraints
- 33 5.6.5. Links between Adaptation and Mitigation
- 34
- 35 5.7. Uncertainties and Data Gaps
- 36
- 37 5.8. Conclusion

38
39 Frequently Asked Questions

40
41 References

42 43 44 **Executive Summary**

45
46 **Coasts are increasingly exposed to varying extreme weather and climate events and impacts from more**
47 **gradual climate change and increased sea-level rise.** The main climate drivers include changing storm regimes,
48 temperature increases, precipitation changes, changes in runoff and sediment transport from watersheds into coastal
49 waters, and increased salinization. It is very likely the mean sea-level rise will contribute to upward trends in
50 extreme coastal high water levels. Locations currently experiencing coastal erosion and inundation will continue to
51 do so due to increasing sea level, in the absence of changes in other contributing factors (*very high confidence*).

52 [5.2.2.1]

53

1 **Vulnerability and exposure to climate change at the coast is exacerbated by population growth, socio-**
2 **economic growth and urbanization.** More than 200 million people are already exposed to flooding by extreme
3 water levels worldwide and this population could be increased by a factor of 4 due to rising population and
4 coastward migration, especially in Asia. Assuming a sea-level rise of 0.5 to 2.0 m and no upgrade in coastal
5 defences, 72 to 187 million people could be displaced due to submergence and erosion by 2100 and about 70% of
6 these affected are from East, Southeast and South Asia (*very high confidence*). [5.3.2, 5.5.3]

7
8 **Impacts of climate change vary globally with different burdens for both developed and developing countries**
9 More assets of developed countries are increasingly affected. In developing countries, the poorer sectors are most
10 vulnerable (5.4.2.1). Developing countries and small island states within the tropics relying on coastal tourism, are
11 impacted not only by weather and climate extremes, future sea-level rise but also the added impacts of coral
12 bleaching and ocean acidification and reduction in tourist flows from mid-latitudes (*very high confidence*). [5.3.2.4]

13
14 **Various approaches in coastal management have made possible for coastal regions to achieve their diverse**
15 **goals in their adaptation to climate change.** Overall, these approaches achieve greater integration, smoother
16 governance, improved social, ecological and economic outcomes, the minimization of risks and impacts from
17 coastal hazards, economic development and use of coastal resources, and protection of coastal environmental
18 resources, natural assets, and ecosystems (*very high confidence*). [Table 5-6, 5.6.1, 5.6.2]

19
20 **While cost of adaptation to sea-level rise is high, the costs of inaction are larger than the sum of adaptation**
21 **and residual damage costs for the 21st century and the global scale.** Even with mean sea-level rise of 2 m by
22 2100, protection is considered economically rational for most countries. Under medium socio-economic
23 development assumption, the expected direct cost of coastal flooding may reach US\$300 billion per year in 2100
24 without adaptation and US\$90 billion per year with adaptation under a 1.26 m sea-level rise scenario (*high*
25 *confidence*). [5.5.4]

26
27 **An extensive set of information is available on global and regional costs of adaptation.** New studies have
28 emerged using a wider range of scenarios, expanded on the impacts considered, and integrating other adaptation
29 options. A wide range of adaptation measures is available, and specific on protection measures and beach
30 nourishment. With additional accessible information available for assessment, e.g. LIDAR (Light Detection And
31 Ranging) data, and knowledge sharing platforms, policymakers should be in a better position to assess local areas
32 for adaptation to climate change (*high confidence*). [5.5.1, 5.5.4, 5.7]

33 34 35 **5.1. Introduction**

36
37 This chapter presents an updated picture of the impacts of climate change and sea-level rise on the coasts. Unlike the
38 coastal chapter in the previous assessment (AR4), some materials pertinent to the oceans are not covered in this
39 chapter but in two new ocean chapters (Chapters 6 and 30).

40
41 The topics to be covered in this chapter are developed along the outline for sectoral chapters approved by the IPCC.
42 Preceding the various sections is an Executive Summary summarizing the key messages with a line of sight to the
43 various sections in the chapter.

44
45 This chapter is organized around eight sections with this first section dealing with the scope, summary and
46 conclusion of the AR4 and key issues. Section 2 provides the necessary definitions that include the coastal systems
47 and climate and non-climate drivers. The coastal systems include both coastal ecosystems and human systems and
48 this division is generally followed for the rest of the sections in the chapter. The observed impacts of climate change
49 on coastal systems and human systems are assessed in section 3 followed by the projected impacts on both systems
50 in section 4. Section 5 assesses the vulnerabilities, risks and costs. Section 6 deals with adaptation and managing
51 risks. There are four case studies distributed within the chapter. Uncertainties and data gaps are assessed in section 7
52 followed by the conclusion in section 8.

53
54 The coasts chapter in AR4 assessed the impact of climate change and global sea-level rise of 0.6 m up to 2100.

1
2 The coastal ecosystems are affected mainly by higher sea level, increasing temperature, changes in precipitation,
3 increased extreme events and reduction in ocean pH. Human activities continued to increase their pressure on the
4 coasts with rapid urbanization in coastal areas and growth of megacities with consequences on the coastal resources.
5 Regionwise, South, Southeast and East Asia, Africa and small islands are most vulnerable. The AR4 chapter offers a
6 range of adaptation measures, many under the ICZM framework that can be carried out in the developed and
7 developing countries, recognizing that the latter would face more challenges. Various issues on increasing the
8 adaptive capacity or increasing the resilience of coastal communities were discussed.
9

10 A number of key issues related to the coasts have arisen since the AR4.

11
12 Coastal systems and their functions and services and how they can be affected by climate change are now better
13 understood. Their linkages landward to the watersheds and seaward to the seas and oceans have to be considered for
14 an integrated assessment of climate change impacts. Semi-empirical models on sea level project higher estimates of
15 sea-level rise by 2100 than that reported in the AR4 but there are still uncertainties (Rahmstorf, 2010; Lowe and
16 Gregory, 2010). This may have serious implications for coastal cities, deltas and low-lying states. While erosion
17 from a higher sea-level rise is expected in future, its relationships with coastal systems such as beaches, barriers,
18 mangroves and coral reefs have to be better established at regional and even local scales and not just at global scale.
19

20 Another concern is ocean acidification and implications of reduced calcification in shellfish impacting worldwide
21 commercial aquaculture (Barton *et al.*, 2012). It also causes coral reefs to lose their structural stability with negative
22 implications for reef communities and shore protection (Kapos *et al.*, 2009; Manzello *et al.*, 2008; see also Box 5-3).
23 An important amount of new findings regarding the impacts of climate change on human settlements, key coastal
24 habitats and ecosystems such as rocky shores, beaches, estuaries deltas, mangroves, coral reefs or submerged
25 vegetation is currently available and reviewed. Unfortunately, it will be shown that uncertainties regarding
26 projections of potential impacts on coastal systems are still high and that further work is required.
27

28 This chapter also provides a more updated assessment of vulnerability, risks and costs to the coasts since the AR4.
29 Assessments of vulnerability have progressed beyond assessment of potential impacts in that they include
30 information on adaptation. A larger number of studies now include estimates of inaction and adaptation.
31

32 The human drivers continue to put heavy pressure on the coasts resulting in increased degradation. Adaptation has
33 been accepted and a wider range of approaches and frameworks such as integrated coastal management, ecosystem-
34 based adaptation, community-based adaptation and disaster risk reduction and management are being used.
35 However, the relative costs of adaptation have to be worked out as well as more information on the constraints and
36 limitations and where to apply. Future land-use in the coastal areas will be dominated by climate change effects and
37 these would be quite profound over the next 50 years (Hadley, 2009).
38

39 On future coasts of developed countries the major effects of climate change will interact with a variety of human
40 activities and drivers of change. For example over the next 50 years on SE England, climate change impacts arising
41 from weather and climate extremes and sea-level rise would occurred with demand for housing and recreational
42 facilities and construction of renewable energy infrastructure at the coast (Hadley, 2009). On coasts of developing
43 countries, the weather and climate extremes put an additional risk to many of the fastest-growing coastal urban
44 areas, such as in Bangladesh and China (McGranahan *et al.*, 2007; Smith 2011)
45

46 5.2. Coastal Systems

47 5.2.1. Definitions

48
49
50
51 Coastal systems include estuaries, coastal plains dominated by mangrove forests and salt marshes, coastal seas and
52 human-built systems. Located at the coastal zone, an interface between purely terrestrial systems and purely marine
53 ones, coastal systems are subject to very large environmental gradients, which, combined with numerous types of
54 geomorphological features, leads to a generally high spatial heterogeneity and high number of habitats. The coastal

1 zone is home to a large variety of important ecosystems whose functions provide goods and services that satisfy
2 human needs, directly or indirectly (De Groot *et al.*, 2002). Ecosystem functions and services can be affected by the
3 variability or long-term change of climatic drivers as well as by non-climatic drivers.

4
5 _____ START BOX 5-1 HERE _____
6

7 **Box 5-1. Definitions Central for this Chapter**

8
9 *Coastal systems:* Include estuaries, coastal plains dominated by mangrove forests and salt marshes, coastal seas and
10 human-built systems.

11
12 *Coastal zone:* Area between purely terrestrial systems and purely marine ones. It is subject to very large
13 environmental gradients, which, combined with numerous types of geomorphological features, leads to a generally
14 high spatial heterogeneity and high number of habitats. Hence, the coastal zone is characterized by strong physical,
15 chemical, biological and biogeochemical interactions and hosts a large variety of ecosystems (Crossland *et al.*,
16 2005). It is also one of the most perturbed areas in the world where non-climate-related drivers are generally greatly
17 affected by human activities and combine with changes in climate-related drivers to affect natural systems and in
18 turn human activities. For the purpose of this assessment, coastal systems and low-lying areas include estuaries,
19 coastal plains dominated by mangrove forests and salt marshes, and coastal seas. Its boundary towards the open
20 ocean is at the continental shelf break, which lies between 110 and 146 m depth (Shepard, 1939 in Sverdrup *et al.*
21 1942), making the marine part of the coastal zone a narrow band with an average width of 34 km (Smith, 2005).
22

23 [INSERT FIGURE 5-1 HERE

24 Figure 5-1: Coastal zone.]
25

26 *Coasts:* Used for convenience to refer to coastal systems and low-lying areas.
27

28 *Ecosystem:* an assemblage of organisms of different types (species, life forms) together with their abiotic
29 environment in space and time (Jax, 2006). The main coastal environments are beaches and intertidal flats, rocky
30 shores, coral reefs, coastal lagoon and lakes, ice shelf (Whitfield and Elliott, 2012).
31

32 *Ecosystem functions:* capacity of natural processes and components to provide goods and services that satisfy human
33 needs, directly or indirectly (De Groot *et al.*, 2002). They are grouped in four categories:

- 34 – *Regulation functions:* relate to the capacity of natural and semi-natural ecosystems to regulate essential
35 ecological processes and life support systems through biogeochemical cycles and other biospheric processes.
36 They provide services that have direct and indirect benefits to humans (e.g., clean air, water and soil, and
37 biological control services).
- 38 – *Habitat functions:* natural ecosystems provide refuge and reproduction habitat to wild plants and animals and
39 thereby contribute to the conservation of biological and genetic diversity and evolutionary processes.
- 40 – *Production functions:* photosynthesis and nutrient uptake by autotrophs converts energy, carbon dioxide,
41 water and nutrients into organic matter which is then used by secondary producers to create an even larger
42 variety of living biomass. This broad diversity in organic matter provides ecosystem goods for human
43 consumption, ranging from food and raw materials to energy resources and genetic material.
- 44 – *Information functions:* Because most of human evolution took place within the context of undomesticated
45 habitat, natural ecosystems provide an essential ‘reference function’ and contribute to the maintenance of
46 human health by providing opportunities for reflection, spiritual enrichment, cognitive development,
47 recreation and aesthetic experience.
48

49 *Ecosystem services:* the benefits, in the form of goods and services, people obtain from ecosystems (Millennium
50 Ecosystem Assessment, 2005). They include goods obtained from ecosystems such as food, fiber, fuel, fresh water
51 and genetic resources, regulating services such as air quality maintenance, climate regulation and water regulation,
52 as well as non-material cultural services such as spiritual enrichment, recreation, and aesthetic experiences (Groot *et*
53 *al.*, 2002). Ecosystem services are provided by ecosystem functions (see ‘Ecosystem functions’).
54

1 *Habitats*: Physical environment in which a species, or assemblage of species, lives.

2
3 *Low-lying areas*: Area or range where coastal and marine processes operate in addition to climate change-related
4 drivers.

5
6 *Drivers*: Any environmental or biotic factor that exceeds natural levels of variation (Breitburg *et al.*, 1999). Climate-
7 related drivers exhibit a wide range of variation at all spatial and temporal scales. This range includes the global or
8 regional annual mean and extreme values, such as sea-level and temperature increases and changes in storm events
9 projected for the next decades. As a result of their location at the interface between atmosphere, land and ocean,
10 coastal systems are subject to a large range of climate-related and non-climate-related drivers.

11 _____ END BOX 5-1 HERE _____
12
13
14

15 5.2.2. *Climatic and Non-Climatic Drivers and Variability*

16 5.2.2.1. *Climatic Drivers*

17
18 Any environmental or biotic factor that exceeds natural levels of variation (Breitburg *et al.*, 1999) is defined as a
19 driver. Climate-related drivers exhibit a wide range of variation at all spatial and temporal scales. This range
20 sometimes includes the global or regional annual mean values projected for the next decades. As a result of their
21 location at the interface between atmosphere, land and ocean, coastal systems are subject to large range of climate-
22 related and non-climate-related drivers.
23

24
25 Climate indices or modes of variability combine complex temporal and spatial changes in several drivers including
26 some considered in the present section, into a simple metric. Since climate does not affect organisms and
27 communities through a single driver but through a blend of multiple drivers, climate indices such as the North
28 Atlantic Oscillation (NAO) and the El Niño Southern Oscillation (ENSO) have been useful to investigate the
29 ecological effects of climate change (Stenseth *et al.*, 2002).
30

31 5.2.2.1.1. *Sea-level including extremes*

32
33 Since human-induced global warming emerged, sea-level rise has been pointed out as a major threat to coastal
34 systems and low-lying areas around the globe (Nicholls, 2010). There is also major concern about higher extreme
35 sea levels due to more intense storms surges and waves superimposed on these mean rises.
36

37
38 It is virtually certain that global mean sea level (MSL) has been rising since 1900 at a rate of 1.7 mm yr⁻¹ and 3.2
39 mm yr⁻¹ since 1993 (AR5, Chap 13). Current observations show large regional variability around the global mean
40 trend on interdecadal periods.
41

42 It is considered likely that extreme sea levels have increased at most locations around the world, largely due to the
43 change in Mean Sea Level (Menendez and Woodworth, 2010) and very likely that mean sea level rise will
44 contribute to upward trends in extreme coastal high water levels Seneviratne *et al.* (2012).
45

46 Consequently, there is *very high confidence* that locations currently experiencing coastal erosion and inundation will
47 continue to do so due to increasing sea level, in the absence of changes in other contributing factors.
48

49 5.2.2.1.2. *Wind, tropical and extratropical storms*

50
51 Extreme wind speeds pose a direct threat to coastal population, the integrity of offshore and coastal infrastructures
52 or to navigation. They also contribute to storm surge and associated flooding events (McInnes *et al.*, 2011). Longer-
53

1 term changes in prevailing winds can cause changes in the stability of sand dunes, or in the wave climate mean
2 energy flux resulting in changes in coastline stability (Reguero *et al.*, 2012).

3
4 A number of recent studies report trends in mean and extreme wind speeds in different areas of the world based on
5 the analysis of instrumental wind observations and numerical reanalysis. Most of them report declining or increasing
6 trends but mostly on continental areas. Only limited studies consider wind stress fields in the ocean, essentially due
7 to the limited long-term, high quality wind measurements in the marine environment.

8
9 Based on reanalysis information, Yang *et al.* (2007) and Xue *et al.* (2010) reported increasing evidence for
10 strengthening of the zonal wind stress field in the Southern Ocean. Using a 23-year database satellite altimeter
11 measurements global changes in oceanic wind speed have shown that the mean and 90th percentile, wind speeds
12 over the majority of the world's oceans have increased by at least 0.25 to 0.5% per year (a 5 to 10% net increase
13 over the past 20 years). The trend is stronger in the Southern Hemisphere than in the Northern Hemisphere. The only
14 significant exception to this positive trend is the central north Pacific, where there are smaller localized increases in
15 wind speed of approximately 0.25%. Extreme wind speeds show a more positive trend increasing over the majority
16 of the world's oceans by at least 0.75% per year (Young *et al.* 2011).

17
18 Due to the shortcomings associated to the length and quality of the observations we have low confidence in wind
19 trends and their causes at this stage (McInnes, et al., 2011; Seneviratne, *et al.*, 2012).

20
21 Consideration of extreme winds requires the analysis of extreme phenomena such as tropical and extratropical
22 cyclones. Tropical cyclones pose a significant threat to coastal population, mostly not due to extreme winds but for
23 the associated storm surge most often combined with fresh water flooding due to extreme rainfall (Rappaport, 2000).
24 There is low confidence that any observed long-term (i.e. 40 years or more) increases in tropical cyclone activity are
25 robust, after accounting for past changes in observing capability (Seneviratne, *et al.*, 2012). Still, since around the
26 mid-seventies, each year, about 90 tropical cyclones occur globally, resulting in a major threat for coastal systems.

27
28 Extratropical cyclones exist throughout the mid latitudes in both hemispheres mainly developing over the oceanic
29 basins. From recent studies it can be concluded that it is likely that there has been a poleward shift in the main
30 northern and southern storm tracks during the last 50 years. There is low confidence in the amplitude, and in some
31 regions, in the sign of regional changes in extratropical storms, (Seneviratne, *et al.*, 2012). Thus the role of changes
32 in the intensity and shift in the geographical location of extratropical cyclones on ocean waves and storm surges
33 requires further studies.

34
35 _____ START BOX 5-2 HERE _____

36 37 **Box 5-2. Case Study – Tropical Cyclones**

38
39 Tropical cyclones, called also typhoons and hurricanes, cause powerful strong winds, torrential rains and high waves
40 and storm surge, all of which have major impacts on people, human systems and ecosystems. Though the strongest
41 storms (Categories 3, 4, and 5) are comparatively rare, they are generally responsible for the majority of damage.
42 For example, Bangladesh and India account for 86% of mortality from tropical cyclones (Murray *et al.*, 2012).
43 Coastal systems and low-lying areas suffer from these impacts.

44
45 Densely populated deltas, particularly in Asia, are recognized as one of the most vulnerable areas to tropical
46 cyclones (Nicholls *et al.*, 2007). The estimated population density is 1,000 people/km² for nine megadeltas in Asia
47 in 2015 (Woodroffe *et al.*, 2006) compared to the average population density of 500 people/km² for 40 deltas
48 globally (Ericson *et al.*, 2006), which is ten times larger than the global average population density.

49
50 Extreme cyclones in Asia since 1970 caused over 0.5 million fatalities (Murray *et al.*, 2012) e.g., cyclones Bhola in
51 1970, Gorky in 1991, Thelma in 1998, Gujarat in 1998, Orissa in 1999, Sidr in 2007, and Nargis in 2008. Tropical
52 cyclone Nargis hit Myanmar on 2 May 2008 and caused over 138,000 fatalities. Several-meter high storm surges
53 widely flooded densely populated coastal areas of the Ayeyarwady Delta and surrounding areas (Revenga *et al.*,

1 2003; Brakenridge *et al.*, 2012). The flooded areas were captured by a NASA MODIS image on 5 May 2008 (Figure
2 5-2).

3
4 [INSERT FIGURE 5-2 HERE

5 Figure 5-2: The intersection of inland and storm surge flooding. Red shows May 5, 2008 MODIS mapping of the
6 tropical cyclone Nargis storm surge along the Ayeyarwady Delta and to the east, Myanmar. The blue areas to the
7 north were flooded by the river in prior years. (From Brakenridge *et al.*, 2012, submitted in 2011).]

8
9 Murray *et al.* (2012) compared the response to Indian Ocean cyclones in Bangladesh (Sidr in 2007) and in Myanmar
10 (Nargis in 2008) in the context of the developments in preparedness and response in Bangladesh through the
11 experiences with previous cyclones Bhola and Gorky and other events. They demonstrated that climate change
12 adaptation efforts could be effective in limiting the impacts from extreme tropical cyclone events by the use of
13 disaster risk reduction methods. These include the construction of multi-storied cyclone shelters, improvement of
14 forecasting and warning capacity, establishing coastal volunteer network, and coastal reforestation of mangroves.
15 Although cyclone Sidr was both slightly more powerful and affected more people than cyclone Nargis, it caused
16 much fewer human losses as a result of the combined disaster risk reduction methods.

17
18 _____ END BOX 5-2 HERE _____
19
20

21 5.2.2.1.3. *Wave climate*

22
23 Since AR4 a series of new studies with detailed quantification and regionalization have confirmed the connection
24 between significant wave height (SWH) and extreme wave heights with climate variability. It is likely that SWH has
25 been increasing over much of the North Pacific since 1900 and the North Atlantic from the 1950s (AR5, chap 3,
26 page 3.17). Several authors reported rates of increase in the SWH, from the 1950s varying from 8 cm per decade to
27 20 cm in winter months in the North Atlantic with smaller values in the North Pacific (Gulev and Gregorieva, 2006;
28 Wang *et al.* 2009; Sendo *et al.* 2010). Dodet *et al.*, 2010 and Wang *et al.*, 2009 report also evidence of increasing
29 peak period during 1953-2009 in the Northeast Atlantic of up to 0.1 s per decade.

30
31 Changes in extreme wave heights have been detected in several areas around the globe. In particular, an increase in
32 the frequency and intensity of the most severe storms has been found in the northeast Pacific (Menendez *et al.*,
33 2008) or North American Atlantic Coast (Komar and Allan, 2008). Significant wave height data sets from 26
34 buoys over the period 1985–2007 reveals significant positive long-term trends in extreme wave height between 30–
35 45°N near the western coast of the US averaging 2.35 cm yr⁻¹ (Izaguirre *et al.*, 2011). These changes at high latitudes
36 in extreme conditions have been confirmed by Young *et al.* (2011) based on a 23-year database of satellite altimeter
37 measurements. More neutral conditions are found in equatorial regions and no clear statistically significant trends
38 for mean monthly values. Conclusions regarding long-term trends on extreme waves should be taken with care
39 considering the relatively short length of available data sets.

40
41 Very limited information of reliable long-term trends in SWH is available in the Southern Hemisphere. Hemer
42 (2010) and Hemer *et al.* (2010) reported the existence of areas with statistically significant increases in SWH.
43 Dragani *et al.* (2010) reported a 7% increase in SWH during 1990 and early 2000s in the area of the South American
44 shelf.

45
46 The mean annual significant wave height is an indicator of how wave climate is evolving under mean conditions,
47 influencing port activities among others. H_{s12}, the significant wave height exceeded on average every 12 hours every
48 year, is intimately linked to the depth of closure of the beach profile (Birkemeier, 1985) and so to potential erosion,
49 as well as the mean energy flow direction, which is related to the transport of sediments and pocket beach planform
50 rotation (González and Medina, 2001).

51
52 Reguero *et al.* (2012), considering numerical reanalysis over the 1948-2010 period, indicated that the largest trends
53 in SWH variation can be found at the Pacific coast of Mexico with about 0.6 cm yr⁻¹ representing almost a 30%
54 increase over 6 decades and also in southern Chile, reaching 1 cm⁻¹. In the range of high percentiles of wave heights,

1 Hs₁₂ seems also to have increased at a rate of about 2 cm yr⁻¹ on the Pacific coast of Mexico and 3 cm yr⁻¹ on the
2 south-eastern margin of the continent, implying about a 45% and 35-45 % of change respectively for the last 6
3 decades. Mean energy flux direction shows sustained changes in the eastern coast from 0.2 (clockwise) to 0.4 deg yr⁻¹
4 (counterclockwise), with lower changes in the western coast of the continent.
5

6 In order to evaluate impacts in coastal areas, it has to be said that trends in SWH in deep water are not necessarily
7 the same as those affecting the beaches and coastal infrastructures, because wave propagation involves several
8 processes, which modify the characteristics of the waves nearshore, (Reguero *et al.*, 2012.)
9

10 5.2.2.1.4. *Temperature changes*

11 More than 70% of the world's coastlines have significantly warmed during the past 30 years, with rates of change
12 highly heterogeneous both spatially and seasonally (Lima and Wetthey, 2012). The average rate is $0.18 \pm 0.16^\circ\text{C}$ per
13 decade and the average change in seasonal timing was -3.3 ± 4.4 days per decade. These values are significantly
14 larger than in the global ocean where the average of change is about 0.1°C per decade in the upper 75 m during the
15 period 1970-2009 (Rhein *et al.*, WGI AR5) and the seasonal shift -2.3 days per decade (Lima and Whethey, 2012).
16 During the period 1985-2005, the annual, night-time, warming of coastal waters along the coasts of the Iberian
17 Peninsula and France exhibited a north-south gradient from 0.12 to 0.35°C per decade (Gómez *et al.*, 2008).
18 Importantly with respect to impacts, the warming also differs seasonally. Gómez *et al.* (2008) have shown that most
19 of the warming occurred in spring and summer, with values as high as 0.5°C per decade. Temperature controls the
20 rate of fundamental biochemical processes such as enzyme reactions and membrane transport (Hochachka and
21 Somero, 2002) with wide-ranging consequences on life history traits (e.g., development rate and survival),
22 population growth and biogeochemical processes in coastal organisms and ecosystems (Hoegh-Guldberg and Bruno,
23 2010; see Table 5-1).
24
25

26 [INSERT TABLE 5-1 HERE

27 Table 5-1: Survey of literature on the impacts of anthropogenic climate change on marine ecosystems (adapted and
28 updated from Hoegh-Guldberg and Bruno, 2010). [To be revised and confined to coastal ecosystems.]
29
30

31 5.2.2.1.5. *Ocean acidification*

32 The oceans absorb about 25% of anthropogenic CO₂ emissions, leading to changes in the carbonate chemistry of
33 seawater, including an increase in the concentration of inorganic carbon and ocean acidity (decreased pH) and a
34 decrease in the concentration of carbonate ion (Box 3.2 in Rhein *et al.*, WGI report). These changes are collectively
35 referred to as anthropogenic ocean acidification and are detectable. The decrease of surface ocean pH ranges
36 between -0.0010 and -0.0018 pH unit yr⁻¹. In contrast with the open ocean where changes in the carbonate chemistry
37 are generally moderate at timescales shorter than 1 year, coastal waters exhibit much larger changes due to changes
38 in upwelling intensity (Feely *et al.*, 2008), deposition of atmospheric nitrogen and sulphur (Doney *et al.*, 2007),
39 carbonate chemistry of the freshwater supply (Salisbury *et al.*, 2008), as well as inputs of nutrients and organic
40 matter (Borges, 2011; Cai *et al.*, 2011) which control primary production (counteracting ocean acidification) and
41 respiration (promoting ocean acidification).
42
43

44 Short-term (hours to weeks) changes of up to 0.5 pH unit are not unusual in coastal ecosystems (Hofmann *et al.*,
45 2011). There are few time series with a timespan of more than 5 years in the coastal ocean (Wootton *et al.*, 2008;
46 Provoost *et al.*, 2010; Waldbusser *et al.*, 2010). Some exhibit considerable differences with the open ocean stations.
47 For example, the surface pH of a southern North Sea station increased as a result of increased availability of
48 nutrients from 1976 to 1987 (Figure 5-3; Provoost *et al.*, 2010). A phosphorus removal policy has limited primary
49 production and led to a decrease in pH much larger than would be expected from the invasion of atmospheric CO₂
50 alone (-0.016 pH unit yr⁻¹). The spatial and vertical variability is also considerably larger than in the open ocean. For
51 example, pH_{NBS} ranges from 6 to 9 in 24 estuaries (Borges and Abril, 2012).
52
53
54

1
2 [INSERT FIGURE 5-3 HERE

3 Figure 5-3: Time series of modelled gross primary production (A; Gypens *et al.*, 2009) and measured pH_T (B) at a
4 fixed station in the southern North Sea (Borges, 2011). pH is expressed on the total scale. Shown is the regression
5 before and after 1987 (solid lines) and the change in pH expected from increased atmospheric CO_2 alone (broken
6 line).]

7
8 Lerman *et al.* (2011) projected that, under the IS92a scenario, pH_T will decrease from about 8.16 in the year 1850 to
9 7.83 in 2100 but the considerable temporal and spatial variability of coastal pH illustrate the fact that ocean
10 acidification generated by the uptake of anthropogenic CO_2 can be greatly lessened or enhanced by coastal
11 biogeochemical processes (Borges and Gypens, 2010; Feely *et al.*, 2010). Cai *et al.* (2011) have shown that
12 atmospheric CO_2 invasion is a significant but minor (24%) component of the 0.45 unit decline in pH in the Northern
13 Gulf of Mexico since pre-industrial time. Using the IS92a CO_2 emission scenario, they also projected that the overall
14 decline in pH by 2100 will reach 0.74, a value that is much greater than that projected in the open ocean.

15 16 17 5.2.2.1.6. *Coastal upwelling*

18
19 The hypothesis that the intensity of coastal upwelling has increased because stronger warming on land compared to
20 the sea leads to the enhancement of upwelling-favourable winds (Bakun, 1990) has recently gained support
21 (Narayan *et al.*, 2010). Upwelled waters are rich in CO_2 and nutrients; they are also cold, leading to a decrease in
22 temperature of 0.3 to 0.4°C per decade since the mid 20th century off Peru (Gutiérrez *et al.*, 2011).

23 24 25 5.2.2.1.7. *Changes in freshwater input*

26
27 Land-use change and climate change have modified river runoff and thus freshwater, sediment and nutrient delivery
28 to coastal systems (Piao *et al.*, 2007). Clearing of land for agricultural use increases erosion, sediment yield and
29 runoff. Although clearing of land for agriculture has started thousands to hundreds years ago depending on the
30 continent (Ruddiman, 2007; Stinchcomb *et al.*, 2011), land-use change has intensified due to human population
31 growth and has increased global runoff on average 0.08 mm y⁻¹ over the last century (Piao *et al.*, 2007). River
32 runoff is generally higher and more variable because of lowered retention due to land clearing (link to other
33 chapters).

34
35 The hydrological cycle is intensified with global warming (Huntington, 2006; link to other chapters), because
36 specific humidity increases approximately exponentially with temperature. Global warming via changes in
37 hydrological cycling is thought to account for about 50% of runoff increase (Piao *et al.*, 2007; cross link required).
38 However, changes are regionally variable. For instance, a detailed 500-yr reconstruction for the Baltic Sea revealed
39 enhanced runoff in the northern Baltic and reduced runoff in the southern Baltic (Hansson *et al.*, 2011). A thorough
40 attribution study revealed that the frequencies of floods have increased significantly in UK and Wales due to
41 increasing greenhouse gas concentrations (Pall *et al.*, 2011).

42
43 Changes in river runoff have multiple effects on coastal systems. Relevant are not only changes in the quantity and
44 quality of runoff but also in the temporal distribution. Freshets or sudden overflow and other pulsed discharges of
45 freshwater into marine systems may impact coastal communities not able to deal with low-salinity water and has
46 consequence for the efficiency of estuaries to retain or filter material delivered by the rivers. Freshwater pulses may
47 cause delivery of riverine nutrients to open sea systems that would otherwise have been processed during transit.

48 49 50 5.2.2.2. *Non-Climate Drivers*

51
52 Coastal systems are subject to a wide range of non climate-related drivers (e.g., Crain *et al.*, 2009) the impacts of
53 which can interact with those climate-related drivers. Some of the major drivers are briefly reviewed below.

1
2 5.2.2.2.1. *Hypoxia*
3

4 The excessive input of nutrients generates coastal eutrophication and the subsequent decomposition of organic
5 matter leads to a decrease in the oxygen concentration (hypoxia) that is reinforced by ocean warming which
6 decreases the solubility of oxygen in seawater. Upwelling of low oxygen waters (e.g., Grantham *et al.*, 2004) and
7 ocean warming (Shaffer *et al.*, 2009) are secondary drivers. Cultural eutrophication induced hypoxia interacts with
8 climate-change induced de-oxygenation. Attribution of low oxygen conditions to natural variability, climate change
9 and cultural eutrophication is therefore difficult (Zhang *et al.*, 2010). Hypoxia poses a serious threat to marine life,
10 which is exacerbated when combined with elevated temperature (Vaquer-Sunyer and Duarte, 2010). The number of
11 so-called “dead zones” has approximately doubled each decade since 1960 (Diaz and Rosenberg, 2008). Fisheries
12 catches from these areas are generally lower than what would be predicted from nutrient loading (Breitburg *et al.*,
13 2009).
14

15
16 5.2.2.2.2. *Water diversion in watersheds*
17

18 Human engineering can affect the runoff of individual river basins to the coastal ocean much more than climate
19 change (Wisser *et al.*, 2010). The main drivers are expansion of irrigation and the construction of structures for
20 water diversion, flood control, power generation and recreation that retains 15% of the global water discharge, hence
21 altering the delivery of sediment and nutrients to coastal systems. An estimated 25% of the world’s river basins run
22 dry before reaching the oceans, due to use of freshwater resources in the basins (Molden *et al.* 2007).
23

24 However, the direct human influence on annual stream flow is small compared with climatic changes during 1948–
25 2004 for most of the world’s major rivers (Dai *et al.*, 2009) and at the global scale (Wisser *et al.*, 2010).
26

27
28 5.2.2.2.3. *Sediment delivery*
29

30 The reduction in sediment delivery to the coast due to trapping behind dams, irrigation (water diversion), sand and
31 gravel mining in river channels causes the sinking of world river deltas, shoreline erosion, threatened mangroves
32 swamps and wetlands, and increased salinization of cultivated land and ground water (Syvitski, 2008). Riverine
33 sediment discharge globally is estimated to be ~20 Gt yr⁻¹ before dam construction in 1950s (e.g., Syvitski *et al.*,
34 2005; Milliman and Farnsworth, 2011), however present sediment discharge has decreased down to 12–13 Gt yr⁻¹
35 (Syvitski and Kettner, 2011). On the other hand soil erosion due to land–use change causes the increase in sediment
36 discharge (e.g., Restropo and Syvitski, 2006), and also impacts coastal ecosystem, particularly coral reefs
37 (McCulloch *et al.*, 2003).
38

39
40 5.2.2.2.4. *Subsidence: relative sea-level rise*
41

42 Subsidence is a common feature of coastal plains and leads to amplified hazards from relative sea-level rise and
43 flooding in the coastal cities built on plains (Mazzotti *et al.*, 2009). Accelerated compaction associated with
44 subsurface resources extraction (gas, petroleum and groundwater) can exceed rates of natural subsidence and global
45 sea-level rise by an order of magnitude (Syvitski, 2008). Increased sediment consolidation due to artificial loads and
46 buildings can lead to significant augmentation of subsidence and relative sea-level rise (Mazzottii *et al.*, 2009).
47

48 5.2.2.2.5. *Habitat loss*
49

50 The conversion of wetlands, intertidal and shallow subtidal habitats to make way for coastal development including
51 land reclamation, harbors or ponds for fish farming is a major factor leading to loss of coastal habitats such as salt
52 marshes, seagrass beds, mangrove forests, beaches and mudflats (Crain *et al.*, 2009).
53
54
55

5.3. Observed Impacts

5.3.1 Impacts on Coastal Habitats and Ecosystems

Coastal habitats and ecosystems have been changed with impacts from both climatic and non-climatic drivers. Halpern *et al.* (2008) have shown that coastal ecosystems, which are subject to both land- and ocean-based anthropogenic stressors, are those experiencing the greater cumulative impact of human activities. Anthropogenic drivers associated with global climate change are distributed widely and are an important component of global cumulative impacts. There are few coral reefs, seagrass beds, mangroves, rocky reefs and shelves with limited impact, and it is hard to discriminate climatic and non-climatic drivers from all ecosystems. However some ecosystems indicate the changes impacted by climate drivers. Coral bleaching impacted by climate change shows detection (D) with very high confidence and attribution (A) with high confidence. Decreased calcification is regarded as D with medium confidence and A with low confidence, and polar-ward expansion of coral species over recent decades is D with high confidence and A with low confidence. However overexploitation and habitat destruction have been responsible for most of the historical changes that occurred in coastal systems (Lotze *et al.*, 2006).

[INSERT FIGURE 5-4 HERE

Figure 5-4: Confidence in Detection and Attribution of observed impacts for coastal systems. Values will be inserted at right positions post FOD, and iterated across chapters to ensure consistency. [Combined one for all coasts still to be developed.]

The following sections assess the impacts of climatic and non-climatic drivers observed in coastal habitats and ecosystems. The summary of these impacts leads to the recognition of key vulnerable systems and hotspots.

5.3.1.1. Rocky Shores

Rocky shores occur at the margins of the oceans throughout the world and can be natural or man-made (e.g., docks, dykes, breakwaters). They are characterized by very steep environmental gradients, especially in the intertidal area where environmental challenges are posed by both aquatic and aerial climatic regimes (e.g., temperature and desiccation). Changes in abundance and distribution of rocky shore species have long been recognized, for example as early as the late 1940s in the North East Atlantic (Hawkins *et al.*, 2008) and perturbation experiments provided information of environmental limits, acclimation and adaptation, particularly to changes in temperature (e.g., Peck *et al.*, 2009a). The challenge is to distinguish the response to changes from climatic drivers, hydrology or from natural temporal and spatial fluctuations.

Species can be eliminated from intertidal habitats by increases in air and water temperature, changes in upwelling regimes, and changes in oxygen levels that can lead to lethal and sublethal effects, which in turn affect population size, species interactions and species persistence. Antarctic ectotherms have very poor abilities to acclimate to elevated temperature and are at least as sensitive as tropical marine ectotherms (Peck *et al.*, 2009a) and an Antarctic brittle star is incapable of acclimating to a temperature of 2°C, only 0.5°C above currently experienced summer maximum temperatures (Peck *et al.*, 2009b). Helmuth *et al.* (2006) reported shifts of range limits of many intertidal species of up to 50 km per decade, much faster than most recorded shifts of terrestrial species. However, the geographical distribution of some species did not change in the past decades. The lack of ranges shifts could be due to weak local warming (Rivadeneira and Fernández, 2005) or overriding effects of variables such as timing of low tide, hydrographic features, lack of suitable bottom types, larval dispersal, food supply, predation and competition (Poloczanska *et al.*, 2011; Helmuth *et al.*, 2002, 2006).

Changes in current patterns and increased storminess can dislodge benthic invertebrates affect the distribution of propagules and recruitment. For example, changes in hurricane activity could subject mussels to more frequent and more severe disturbances compared to those that occurred during 1971-1994 (Carrington, 2002).

1 Rocky shores are one of the few ecosystems for which field evidence of effects of ocean acidification is available.
2 Wootton *et al.* (2008) provided observational and modeling analysis of rocky shore community dynamics in relation
3 to pH and associated physical factors over nine years (2000–2008). The community structure shifted from a mussel
4 to an algal-barnacle-dominated community but attribution to a specific driver or set of drivers is difficult.
5 Observations near natural CO₂ vents in the Mediterranean Sea showed profound changes of the community structure
6 of shallow rocky shores at an average pH value around the one expected in 2100. Subtidal calcifiers are absent
7 below mean pHT 7.8 (Hall-Spencer *et al.*, 2008) while calcareous and turf algae are significantly reduced and other
8 macroalgae are tolerant (Porzio *et al.*, 2011). The negative effects of ocean acidification on several Mediterranean
9 rocky shore invertebrates are mostly due to increased calcium carbonate dissolution in organisms that have no
10 organic protective layer or have lost it, and are exacerbated at higher temperatures (Rodolfo-Metalpa *et al.*, 2011).
11 Similar tolerance to low pH is found in calcifying invertebrates of the Baltic Sea (Thomsen *et al.*, 2010), probably at
12 the cost of increased energy expenditure (Thomsen and Melzner, 2010). Increased acidity increases the rate of
13 dissolution of the calcium carbonate framework.
14
15

16 5.3.1.2. Beaches and Sand Dunes

17

18 Throughout the world, beaches and dunes, as well as bluffs and cliffs, have in general been undergoing net erosion
19 over the past century or longer (e.g. Bird, 2000, estimated at least 70% of the world's sandy beaches were eroding).
20 This erosion is due to a variety of processes only some of which are climate related, like rising mean sea levels
21 (Ranasignhe and Stive, 2009); changes in the frequency and severity of transient storm associated erosion events
22 (Tebaldi *et al.*, 2012); wave propagation caused by sea-level changes realigning shorelines (Tamura *et al.* 2010);
23 sustained changes in the direction of mean wave energy (Reguero *et al.* 2012); changes in the loss of natural
24 protective structures such as coral reefs (Grevelle and Mimura, 2008) or mangrove forests due to increased ocean
25 temperatures or ocean acidification (Bongaerts *et al.* 2010); permafrost degradations and sea ice retreat, which
26 exposes soft shores to waves and storms (Manson and Solomon, 2007). These processes may act at different time
27 scales. For example, there is high confidence that erosion on the beaches of Southeast Australia is intimately related
28 to interannual changes in swell direction (Harley *et al.* 2010).
29

30 Non-climate related processes include reductions of sand supply, through trapping by river dams or coastal
31 protection structures. Reasonably accurate maps have been available since about the mid-19th century to compare to
32 more recent maps and imagery to quantify combined climate and non-climate changes. For example, the long-term
33 rate of erosion along the U.S. Mid-Atlantic and New England coasts is 0.5 ± 0.09 m yr⁻¹ (Hapke *et al.*, 2011). This is
34 based on over 21,000 measurement locations equally spaced along the more than 1,000 km of coast, 68% of which
35 indicated net erosion. Arctic coasts have experienced some of the greatest magnitudes of erosion in the world, for
36 example in parts of Alaska as much as 0.9 km of retreat in 50 years (Mars and Houseknecht, 2007) and in parts of
37 Siberia a range of 434 m of retreat to 92 m of deposition in 56 years (Lantuit *et al.*, 2011). However, in a survey of
38 27 atoll islands in the central Pacific, 43% of islands were found to have remained stable and 43% were found to
39 have increased in area (Webb and Kench, 2010). Where an eroding shoreline approaches hard, immobile, structures
40 like seawalls or resistant natural cliffs, the beaches will narrow due to coastal squeeze that removes the sands and
41 associated habitats, and steepens the beach slope, impacting the survivability of a variety of organisms (e.g. in
42 northern Scotland, see Jackson and McIlvenny, 2011). With coastal squeeze, sand dunes will ultimately be removed
43 as the beach erodes and narrows. Extreme storms can erode and completely remove dunes, degrading land
44 elevations and exposing them to inundation and further change (e.g. Plant *et al.*, 2010).
45

46 Unfortunately, due to the scarcity and fragmentary nature of the information available and to the multiple natural
47 and anthropogenic drivers contributing to coastal erosion, attributing shoreline changes to climate change is still
48 difficult. In the absence of adaptation measures there is a high confidence that beaches and sand dunes currently
49 affected by erosion, will continue to do so under increasing sea levels or changing wave heights and mean energy
50 flux direction.
51
52
53

5.3.1.3. Estuaries, Tidal Flats, and Lagoons

Estuaries connect rivers with adjacent coastal systems and are the primary conduit for water, nutrients and particulates from the continent to the sea. Depending on the hydrology, oceanographic, climatic and geological settings estuaries can be well-mixed or stratified, shallow or deep, river or tidally dominated, but a common characteristic is the presence of fresh and marine water within the system and consequently salinity gradients (Heip *et al.*, 1995; Breitburg *et al.*, 2009).

Riverine transport of particles and delivery of suspended matter from the sea supports high rates of sediment deposition in estuarine systems. Sediment accumulation in estuaries is heterogeneous and habitat specific (generally little in the main channels and more accumulates in marginal systems such as marshes) and affected directly by dredging activities for shipping and indirectly via habitat loss, sea-level, storminess and land-use changes related changes in sediment supply by rivers (Syvitski *et al.*, 2005). Climate and non-climate induced changes in estuarine sediment budgets have consequences for carbon, nutrients and contaminants budgets.

Estuarine systems, with low tidal ranges, are strongly affected by run-off since the water residence time is primarily governed by runoff. Water residence time is a key predictor and governing factor for many ecosystem and biogeochemical processes including nutrient processing, the metabolic balance, carbon dioxide exchange rates and hypoxia (Howarth *et al.*, 2009). Floods, freshets and other runoff events may diminish estuarine communities and in that way the processing of organic matter and nutrients in these systems.

Estuaries are known to be sites with high-intensity water-air and sediment-air carbon dioxide exchange. Most estuaries are a source of carbon dioxide to the atmosphere (Borges, 2005), the global carbon dioxide emission rate is about 0.25 Pg y⁻¹ (Cai, 2011; Laruelle *et al.*, 2010). Although most researchers agree that estuaries emit carbon dioxide there is debate on whether the carbon dioxide originates from riverine carbon, i.e. input of carbon dioxide rich rivers and respiration fuelled by riverine particulate and dissolved organic matter, or from within mangroves and tidal marshes within the estuary (Borges, 2005; Hofmann *et al.*, 2008; Cai, 2011).

Increasing atmospheric carbon dioxide levels would theoretically impede these effluxes (lower gradient from water to air), but this is difficult to detect because of the high heterogeneity and large temporal variability of estuarine carbon dioxide pressures (Borges, 2005; Chen and Borges, 2009). Increasing atmospheric carbon dioxide may also lead to acidification of estuarine waters and if waters become undersaturated with respect to calcium carbonate, this may have major consequences for some calcifiers, including ecological key species such as ecosystem engineers and commercially important species (e.g., oysters, mussels, Gazeau *et al.*, 2007). However, acidification of estuarine waters is not only due to atmospheric carbon dioxide uptake as in the open ocean and on the continental shelf, but also due to mixing of fresh and marine waters, input of riverine waters rich in carbon dioxide and nitrification supported by high ammonium concentrations (Salisbury *et al.*, 2008; Hofmann *et al.* 2009). Changes in eutrophication and the balance between production and respiration have been identified to overrule atmospheric carbon dioxide induced acidification processes in coastal waters (Borges and Gypens, 2010). A detailed analysis of long-term time series for estuarine waters in the Dutch coastal zone revealed large changes, both increases and decreases (Provoost *et al.*, 2010).

Riverine delivery of nutrients has increased significantly the last century and is projected to increase further (Bouwman *et al.*, 2011). The elevated nutrient loadings to estuaries have resulted in major changes in biogeochemical processes, community structure and ecosystem functions (Howarth *et al.* 2009). Eutrophication has modified food-web structure, has led to more intense and longer lasting hypoxia and to more frequent occurrence of harmful algal blooms (Breitburg *et al.*, 2009; Howarth *et al.*, 2009). These nutrient-induced environmental issues have affected estuarine fishery yield and sustainance.

Coastal lagoons are shallow bodies of seawater or brackish water separated from the ocean by a barrier, connected at least intermittently to the ocean. Coral reef lagoons are considered elsewhere in this chapter. Temperate coastal lagoons are formed and maintained through sediment transport and are therefore highly susceptible to alterations of sediment input from land and erosional processes driven by changes in sea level, precipitation, and storminess.

1
2 Temperate coastal lagoons often host salt marshes, seagrasses and macroalgae (see sections 5.3.1.5, 5.3.1.7) and
3 aquaculture. Due to their restricted exchange with the adjacent ocean, they are particularly vulnerable to
4 eutrophication.

5
6 On average, the fisheries yield is higher in coastal lagoons than in other ecosystems (Kapesky, 1984 in Pauly and
7 Yáñez-Arancibia, 1994) and there appears to be an empirical correlation between primary production and fisheries
8 yields (Nixon, 1982). Hence, any change in primary production generated by climatic or non-climatic drivers could
9 impact fisheries. For example, it has been shown that changes in water temperature and reduction in plankton
10 productivity caused by the modification of seasonal precipitation patterns will negatively affect clam aquaculture in
11 the lagoon of Venice (Canu *et al.*, 2010). Small changes in salinity were also shown to generate major changes in
12 food webs (Jeppesen *et al.*, 2007) but the global impact on lagoon fisheries remains uncertain.

13 14 15 5.3.1.4. Deltas

16
17 Coastal zones receive substantial amounts of nutrient and sediment from rivers, and deltas are formed at river-mouth
18 areas influenced by a combination of river, tide and wave processes. Deltas are one of the most important
19 ecosystems and habitats, and also areas for socio-economic human activities. Deltas consist of a compound coastal
20 system of natural systems, e.g., tidal flats, salt marshes, mangroves, beaches, estuaries, low-lying wetlands, and also
21 human systems, e.g., houses, agriculture, aquaculture, industry, transport, with a high population density of more
22 than 10 times the world average (Ericson *et al.*, 2006; Foufoula-Georgiou *et al.*, 2011). However deltas are among
23 the highly vulnerable and stressed coastal systems and are subjected to human and climate impacts from both
24 drainage basins of rivers and oceans, and also within delta itself: e.g., changes in runoff, sea-level rise, as well as
25 human activities as land-use changes, dam construction, irrigation, mining, extraction of subsurface resources,
26 urbanization. In particular, Asian megadeltas and low-lying coastal urban areas including megacities are identified
27 as one of vulnerable areas (Nicholls *et al.*, 2007).

28
29 Decreased sediment discharge due to construction of dams and irrigation makes imbalance of sediment in coastal
30 zones, resulting in loss of beaches, mangroves and tidal flats by coastal erosion (e.g., Nile and Ebro, Sanchez-Arcilla
31 *et al.*, 1998; Po, Simeoni and Corbau, 2009; Krishna-Godavari, Nageswara Rao *et al.*, 2010; Changjiang, Yang and
32 Milliman, 2011; Huanghe, Chu *et al.*, 1996). In particular, 25% of the wetlands of the Mississippi Delta have been
33 lost over the last few centuries (Blum and Roberts, 2009). Fluctuations in sea-level rise rate and sediment supply
34 mainly dominate the formation and evolution of coastal wetlands. The wetlands with high sediment input, mainly
35 riverine are only ones for surviving accelerated sea-level rise, based on a comparative study of the wetlands of
36 Mediterranean deltas and lagoons (Day *et al.*, 2011). High sediment input and high capture efficiency of sediments
37 including inundation frequency by tides are necessary for sediment accumulation (Day *et al.*, 2011; Andersen *et al.*,
38 2011). The dominant species in the marsh, together with nutrient availability, also control the rate of organic peat
39 production. The highest rates of marsh vertical accretion are found in fluvially dominated systems due to high
40 inorganic sediment influx (FitzGerald *et al.*, 2008). On the other hand, land-use change from natural delta plains to
41 agricultural and industrial use is a major cause of wetlands loss more than natural causes. Globally, 42 deltas show
42 wetland loss with an average annual rate of 26,000 km² for the last 14 years (Coleman *et al.*, 2008).

43
44 Major sea-level rise impacts are coastal wetland change, increased coastal flooding, increased coastal erosion, and
45 saltwater intrusion into estuaries and deltas (McLeod *et al.*, 2010b). Subsidence due to natural and anthropogenic
46 compaction of underlying sediments is a common feature of river deltas and leads to amplified hazards from relative
47 sea-level rise in the coastal cities built on deltaic plains (Day and Giosan, 2008; Mazzotti *et al.*, 2011). The most
48 dramatic subsidence effects have been caused by drainage and groundwater fluid withdrawal. Over the 20th century,
49 coasts have subsided by up to 5 m in Tokyo, 3 m in Shanghai, and 2 m in Bangkok. To avoid submergence and/or
50 frequent flooding, these cities now all depend on a substantial flood defence and water management infrastructure
51 (Nicholls *et al.*, 2010). In Thailand the extreme Chao Phraya flood of 2011 in the delta plain caused a loss of human
52 life and impacted the global economy. Increased sediment consolidation due to artificial loads can also lead to
53 significant augmentation of subsidence and relative sea-level rise. For the Fraser River delta, areas with recent large
54 structures may undergo relative sea-level rise of as much as ~1–2 m (Mazzotti *et al.*, 2011). Relative sea-level rise

1 also impacts the decrease in sediment delivery from rivers to the coastal zone by deposition in river channels in
2 deltas in the Po river delta (Syvitski *et al.*, 2005).

3
4 Thirty-three deltas in the world show that 85% of the deltas experienced severe flooding in the past decade, resulting
5 in the temporary submergence of 260,000 km² (Syvitski *et al.*, 2009). Deltas have received substantial impacts by
6 river floods and from oceans by storm surges and tsunamis. Tropical storms also have impacted ecosystems and
7 human systems in deltas, e.g., the city of New Orleans and the Mississippi Delta by Hurricane Katrina in 2005
8 (Dixon *et al.*, 2006), Ayeyarwady (Irrawaddy) Delta by Cyclone Nargis in 2008, Ganges-Brahmaputra Delta by
9 Cyclone Gorky in 1991 and Cyclone Sidr in 2007 (Murray *et al.*, 2012). The tropical cyclones in the North Indian
10 Ocean account for 86% of the world mortalities (ISDR, 2009), which occur mostly in highly populated deltas in this
11 region.

12 13 14 5.3.1.5. *Mangroves and Salt Marshes*

15
16 Mangrove forests occur along the coast of more than hundred countries. These ecosystems provide many functions
17 including coastal defence, nursery grounds for fishes and carbon storage (Bouillon *et al.*, 2008, Feller *et al.*, 2010).
18 Mangrove trees are found in the intertidal zone along subtropical and tropical coasts. These forests are essential in
19 protecting shorelines (Gedan *et al.*, 2011). They stabilize sediments and enhance settling and retention of fine-
20 grained sedimentary materials. Mangrove forests act as sediment sinks and as consequence of this also as organic
21 carbon sinks (Duarte *et al.*, 2005).

22
23 The area of mangrove forests has declined by 30 to 50% during the last 50 years due to coastal development, over-
24 harvesting and increasing use for aquaculture (Duarte *et al.*, 2005; Donato *et al.*, 2011; Irving *et al.*, 2011). Clear-
25 felling to generate space for commercial pond aquaculture for fish and crustacean is in particular important. Annual
26 rate of areal decrease for the period 1970 to 2000 were about 2% yr⁻¹ (Duarte *et al.*, 2005; Irving *et al.*, 2011),
27 implying that without further protection they will disappear in as little as 100 years. This will have consequences for
28 coastal protection and carbon burial. Mangrove forests are the most carbon dense forest on earth with about 1 Gg
29 carbon stored per ha, primarily below ground (Donato *et al.*, 2011). Reclamation of mangrove forest results in 112
30 to 392 Mg C loss per ha, depending on the depth to which soil carbon is oxidized. This represents 0.02-0.12 Pg yr⁻¹,
31 as much as around 10% of emissions from deforestation globally, despite mangroves accounting for just 0.7% of
32 tropical forest area (Donato *et al.*, 2011). This carbon loss should be combined with the loss of long-term carbon
33 sequestration because of loss of organic carbon burial that would otherwise occur (Duarte *et al.*, 2005; Irving *et al.*,
34 2011).

35
36 Coastal wetlands are prominent features and important habitats along the coastline. Mangroves dominate subtropical
37 and tropical coastlines while tidal marshes (saline, brackish and fresh-water tidal) dominate temperate systems.
38 Saltmarshes provide many ecosystem functions and services including coastal defence against storms and waves,
39 nutrient removal and transformation, nursery for fish and shrimp, fishing, carbon burial and tourism (Bromberg
40 Gedan *et al.*, 2009; Irving *et al.*, 2011). Coastal marshes play a major role in protecting shorelines via multiple
41 mechanisms including wave attenuation and shoreline stabilization (Bromberg Gedan *et al.* 2011). Saltmarshes have
42 been used and shaped by humans since Medieval Times. Human impacts include use as pasturelands for livestock,
43 use of marsh plants for construction, conversion of marshes into agricultural, urban and industrial use (Bromberg
44 Gedan *et al.* 2009). Moreover, deliberate introduction of species and invasive species have modified marsh
45 communities and functioning (Neira *et al.*, 2006). Intertidal *Spartina* and phragmites have been introduced
46 deliberately for coastal protection or were favoured by nutrient enrichments. Changes in marsh hydrology due to
47 ditching or tidal restriction have significantly affected coastal marsh distribution patterns and functioning (Bromberg
48 Gedan *et al.*, 2009; 2011).

49
50 Saltmarshes represent a major sink for sediment and thus organic carbon (Duarte *et al.*, 2005). Any loss of saltmarsh
51 area (climate change, habitat destruction) thus lowers natural CO₂ sequestration potential (Irving *et al.*, 2011).
52 Decline in saltmarsh area, therefore, exacerbates climate change and also implies that shorelines become more
53 vulnerable to erosion due increased sea level rise and increased wave action.

1 The distribution of tidal marshes is closely linked with sea level and thus sea level rise. Historical records show that
2 saltmarshes have generally adapted accretion rates to match sea-level rise (Redfield, 1972). The response of
3 saltmarsh to sea-level rise involves landward migration of salt marsh vegetation zones and submergence at lower
4 elevations and drowning of interior marshes. Marsh can increase accretion rates by either accumulating more
5 external mineral particles or by accumulation of peat, the relative importance of these two modes of accretion
6 depending on geological setting and ecosystem production (Allen, 1995; Middelburg *et al.*, 1997)

7 8 9 5.3.1.6. Coral Reefs

10
11 Coral reefs are shallow-water structures made of calcium carbonate mostly secreted by scleractinian corals and
12 algae. They harbour a biodiversity that is disproportionately high compared to their surface area and are sources of
13 key services to humans. Coral reefs are susceptible to several climatic (sea-level rise, warming and ocean
14 acidification) and non climatic (e.g., coastal development, pollution, nutrient over-enrichment and overfishing).
15 Most human-induced disturbances until the early 1980s were local but climate-related disturbances have become
16 more obvious in recent decades.

17
18 Coral reef growth is intimately linked to sea level. Within the uncertainties of both the estimates of past sea level
19 rise and coral reef growth, most reef ecosystems seem to have kept pace with sea-level rise during the past 100 years
20 (Buddemeier and Smith, 1988).

21
22 Increased temperature triggers bleaching of corals, which are key reef ecosystem engineers (Wild *et al.*, 2011).
23 Bleaching involves the loss of endosymbiotic algae, which live in the coral tissues and play a key role in their
24 physiology, especially nutrition (Baker *et al.*, 2008) (see chapter 6 for physiological details and chapter 30 for a
25 regional analysis). Mass coral bleaching has occurred in association with episodes of elevated sea temperatures over
26 the past 30 years (e.g., Hoegh-Guldberg, 1999; Kleypas *et al.*, 2008). For example, the level of thermal stress at
27 most of the 47 reef sites where bleaching occurred during 1997-98 was unmatched in the period 1903 to 1999
28 (Lough, 2000). The intensity of bleaching events is very variable on yearly timescale: the percentage of reef cells
29 which exhibited at least one bleaching event was 7% in 1985-1994 and 38% in the subsequent decade due to intense
30 El Niño events. Bleaching is not always fatal for coral colonies; recovery depends on (1) the magnitude and duration
31 of the elevated temperature event, (2) the species that have been lost, (3) the acclimation potential of the species
32 remaining, and (4) the interaction with other drivers. Reef recovery from the 1998 global bleaching event was
33 significant but slow in the Indian Ocean (median rate of recovery of about 1% yr⁻¹, absent in the western Atlantic
34 and locally variable elsewhere (Baker *et al.*, 2008). It has also been limited in the southern Arabian Gulf although
35 the community is among the most tolerant to environmental extremes (Burt *et al.*, 2011).

36
37 The increase in temperature is also suspected to have caused a poleward range expansion of corals (e.g., Precht and
38 Aronson, 2004). The northward speed along the coasts of Japan is to up to 14 km y⁻¹ since the year 1930, with no
39 evidence of southward range shrinkage or local extinction (Yamano *et al.*, 2011). Some of the new sightings could
40 be due to a recent increase in monitoring and the introduction of larvae or adults by humans can contribute to range
41 expansion. Nevertheless, increased temperature favours range shifts.

42
43 The decline of seawater pH has become a recent source of concern for the future of coral reefs. The geological
44 record indicates that four of five global metazoan reef crises in the last 500 Myr were probably at least partially
45 governed by ocean acidification and rapid increase in temperature (Kiessling and Simpson, 2011). Experimental
46 evidence shows that lower pH decreases the rate of calcification of most, but not all, reef-building corals and
47 coralline algae (reviewed in Andersson *et al.*, 2011; Pandolfi *et al.*, 2011) and enhances the competitiveness of
48 seaweeds over corals (Diaz-Pulido *et al.*, 2011).

49
50 Retrospective studies have provided clear outcomes but attribution to drivers has proven difficult. Although
51 perturbation experiments suggest that coral calcification may have decreased since the beginning of the industrial
52 revolution, clear evidence is not available in all field samples. Most (e.g., De'ath *et al.*, 2009, Manzello, 2010), but
53 not all (Bessat and Buigues, 2001; Helmle *et al.*, 2011) retrospective studies show decreasing trends in calcification
54 for the past several decades but whether the decrease is due to ocean acidification, other environmental drivers (e.g.,

1 ocean warming), or a combination of drivers remains unclear. Despite a few shortcomings (Riebesell, 2008),
2 observations near CO₂ vents (Fabricius *et al.*, 2011) have shown that ocean acidification has dramatic impacts on the
3 biodiversity of natural communities even though reef-building corals are not completely eliminated at lower pH
4 (pH_T 7.73 to 8.00) compared to control (pH_T 7.97–8.14) and the rate of calcification of the one of the resistant
5 species exhibits small changes relative to pH. At lower pH, the community composition tips in favour of seagrasses
6 and fleshy non-calcareous macroalgae and against hard corals.

7
8 Published evidence supports the hypothesis that coral infectious diseases are emerging in response to drivers such as
9 ocean warming, altered rainfall, increased storm frequency, sea level rise, altered circulation, and ocean acidification
10 (Sokolow, 2009). Based on population reduction in the recent past, it is estimated that one third of all coral species
11 exhibit a high risk of extinction (Carpenter *et al.*, 2008). Although less well documented, non-coral benthic
12 invertebrates are also at risk (Przeslawski *et al.*, 2008). Reef fish are also vulnerable, less from climatic drivers than
13 from overfishing (Graham *et al.*, 2011).

14
15 Almost half of all coral reefs experience medium high to very high impact of human activities (30-50% to 50-70%
16 degraded; Halpern *et al.*, 2008). Many coral reefs have been subject to widespread degradation since the 1970. In the
17 Indo-Pacific, the estimated yearly coral cover loss was approximately 1% over the last twenty years (Bruno and
18 Selig, 2007). For example, in Jamaica, coral cover has declined from more than 50% in the late 1970s to less than
19 5% in the early 1990s. A dramatic phase shift has occurred producing a system dominated by fleshy macroalgae
20 (more than 90% cover).

21 _____ START BOX 5-3 HERE _____

24 **Box 5-3. Case Study – Coral Reefs**

25 [cross chapter (5, 6, 25, and 30) box]

26
27 Coral reef ecosystems are mostly distributed in the tropics and play multiple key roles despite their relatively small
28 surface area. Almost 500 million people live within 100 kilometers of a coral reef (Burke *et al.*, 2011), deriving
29 benefits such as provisioning functions (e.g. food, construction material, medicine), regulating functions (e.g.
30 shoreline protection, maintenance of good water quality), cultural functions (e.g., tourism) and supporting functions
31 (e.g. oxygen supply) (Hoegh-Guldberg, 2011). A wide range of climatic and non-climatic drivers affect coral reefs
32 and negative impacts are already observed (section 5.3.1.6). Coral bleaching, largely triggered by positive
33 temperature anomalies, is the most widespread and conspicuous impact (Figures 5-4 and 5-5) (Chapter 6, section
34 6.2.2.4; Chapter 25, section 25.6.3; Chapter 30, section 30.5.3-6 and 30.10.2). Increased seawater acidity limits the
35 calcification rate of many coral reef builders, increases reef dissolution, and reduces biodiversity (Figure 5-5c,d).

36
37 [INSERT FIGURE 5-5 HERE

38 Figure 5-5: The same coral community before and after a bleaching event in February 2002 at 5 m depth, Halfway
39 Island, Great Barrier Reef. Coral cover at the time of bleaching was 95% almost all of which was severely bleached,
40 resulting in mortality of 20.9% (Elvidge *et al.* 2004). C and D: three CO₂ seeps in Milne Bay Province, Papua New
41 Guinea show that prolonged exposure to high CO₂ is related to fundamental changes in coral reef structures
42 (Fabricius *et al.*, 2011). Coral communities at three high CO₂ (median pH_T 7.7, 7.7 and 8.0), compared with three
43 control sites (median pH_T 8.02), are characterised by significantly reduced coral diversity (-39%), severely reduced
44 structural complexity (-67%), low densities of young corals (-66%) and few crustose coralline algae (-85%). Reef
45 development ceases at pH values below 7.7. Photo credit: R. Berkelmans (A and B) and K. Fabricius (C and D).]

46
47 There is robust evidence and high agreement that coral reefs are one of the most vulnerable marine ecosystems
48 (Chapters 5, 6, 25 and 30). Globally, more than half of the world's reefs are under medium or high risk (Burke *et al.*,
49 2011) even in the absence of climatic factors. Future impacts of climate drivers (warming, sea level rise and
50 increased acidity) will considerably exacerbate the impacts of non-climatic drivers (high agreement, robust
51 evidence).

52
53 Damages to coral reefs have implications for several key regional services:

- 1 • Resources: Coral reefs produce 10-12% of the fish caught in tropical countries, and 20-25% of the fish
2 caught by developing nations (Garcia and Moreno, 2003). Over half (55%) of the 49 island countries
3 considered by Newton *et al.* (2012) are already exploiting their coral reef fisheries in an unsustainable way.
- 4 • Tourism: More than 100 countries benefit from the recreational value provided by their coral reefs (Burke
5 *et al.*, 2011). For example, the Great Barrier Reef Marine Park attracts about 1.9 million visits each year
6 and generates A\$ 5.4 billion to the Australian economy and 54,000 jobs (90% in the tourism sector; Biggs,
7 2011).
- 8 • Coastal protection: Coral reefs protect the shoreline from the destructive action of storm surges and
9 cyclones, providing the only habitable land for several island nations and habitats suitable for the
10 establishment and maintenance of productive mangroves and wetlands, as well as areas for recreational
11 activities. This role is threatened by sea-level rise as well as the decrease in coral cover, lower rate of
12 calcification, and higher rate of dissolution of the reef framework due to ocean warming and acidification.
13

14 Marine protected areas (MPAs) may be useful to increase ecosystem resilience and moderate the impacts of climate
15 change (McLeod *et al.*, 2009). Although MPAs are a key conservation and management tool, they have generally no
16 effect on the resilience of coral reefs to thermal stress (Selig *et al.*, 2012) and hence alternative strategies need
17 consideration (Rau *et al.*, 2012). Controlling the input of nutrients and sediment from land is an important
18 complementary management strategy. However, there is *high confidence* that, in the long term, limiting the amount
19 of warming and acidity are the most important tools to maintain viability of current coral reef systems and
20 communities.

21 _____ END BOX 5-3 HERE _____
22
23
24

25 5.3.1.7. Submerged Vegetation

26
27 Seagrass meadows are ecosystems composed by marine angiosperms, a group of about 60 species of clonal
28 angiosperms, distributed in shallow coastal areas of all continents, except Antarctica (Hemminga and Duarte, 2000).
29 Seagrass meadows rank amongst the most valuable ecosystems, in terms of the services and benefits they support, in
30 the biosphere, but are also highly vulnerable and about one third of the area they occupied has been lost since World
31 War II, declining globally at rates of 7% y⁻¹ since 1990 (Orth *et al.*, 2006; Waycott *et al.*, 2009). Whereas
32 eutrophication is recognised as the primary force accounting for the global seagrass decline (Duarte 2000, Orth *et al.*
33 2006; Waycott *et al.*, 2009), seagrass meadows are vulnerable to climate change (Short and Neckles, 1999, Duarte,
34 2000). Climate change affects seagrass meadows in multiple ways, as seagrass meadows are affected by warming,
35 sea-level rise, and changes in wave energy and storminess (Short and Neckles, 1999; Duarte, 2000).
36

37 Seagrass meadows are particularly vulnerable to temperature extremes, as many seagrass meadows occur in areas
38 where maximum temperatures are close to their physiological maxima. In these situations increased maximum
39 temperature by a few degrees Celsius triggers seagrass mortality (e.g. Massa *et al.*, 2009; Marbà and Duarte, 2010).
40 Evidence for negative effects of high temperature on seagrass biomass has been reported for seagrass meadows in
41 the Atlantic Ocean (Reusch *et al.*, 2005), Mediterranean Sea (Marbà and Duarte, 2010) and Australia (Rasheed and
42 Unsworth, 2011). Heat waves lead to widespread seagrass mortality as documented for *Zostera* species, the
43 dominant seagrass genus in the Atlantic (Reusch *et al.*, 2005), and *Posidonia oceanica*, the dominant species in the
44 Mediterranean Sea (Marbà and Duarte, 2010). In particular, Marbà and Duarte (2010) demonstrated that *P. oceanica*
45 meadows are highly vulnerable to warming, as demonstrated by a direct functional relationship between maximum
46 seawater temperature and mortality rates of *P. oceanica* shoots, with shoot mortality rates increasing by 0.022 yr⁻¹
47 for each additional degree of annual maximum temperature. Warming also triggers flowering of *P. oceanica* (Díaz-
48 Almela *et al.*, 2007), but the increased recruitment rate is insufficient to compensate for the losses resulting from
49 elevated temperature (Díaz-Almela *et al.*, 2009). Current observations indicate that seagrass meadows are already
50 under stress due to realised climate (e.g. Marbà and Duarte, 2010, Rasheed and Unsworth, 2011).
51

52 Seagrasses, particularly those in shallow waters, are often carbon-limited (Hemminga and Duarte, 2000), and may
53 benefit from increased CO₂. Due to the realised increased in CO₂ concentration in surface waters (Duarte, 2002),

1 seagrass photosynthetic rates may have already increased by 20% (Hemminga and Duarte, 2000; Hendriks *et al.*,
2 2010).

3
4 Sea-level rise may result in the upslope migration of seagrass meadows, with both their shallow and depth limit
5 migrating upwards to maintain their depth range (Duarte, 2002). However, sea-level rise often results in submarine
6 erosion and the loss of seagrass meadows, particularly where shorelines have been occupied by infrastructure
7 (Marbá and Duarte, 1997; Duarte, 2002). Extreme events, such as droughts, can also impact on estuarine seagrasses.
8 Cardoso *et al.* (2008) concluded that extreme weather events contributed to the overall degradation of seagrass
9 meadows in a Portuguese estuary.

10
11 Loss of seagrass meadows with climate change erodes natural CO₂ sequestration potential, as seagrass meadows act
12 as CO₂ sinks, ranking among the most intense CO₂ sinks in the biosphere (Duarte *et al.*, 2010; Kennedy *et al.* 2010).
13 Loss of seagrass meadows, therefore, aggravates climate change and also renders shorelines more vulnerable to
14 erosion due increased sea level rise and increased wave action.

15
16 Macroalgal beds grow in shallow coastal areas worldwide, including rocky and sandy shores, and form highly
17 productive communities with rapid turnover. Temperature affects growth and biogeographic ranges of macroalgae,
18 especially in polar and cold-temperate regions. Macroalgae are also affected by increased CO₂, which is expected to
19 lead to enhanced photosynthetic rates (Wu *et al.*, 2008).

20
21 Contrasting response of macroalgae and corals to climate change has lead to the prediction of a tendency for phase
22 shifts from corals to macroalgae. However, a recent global assessment concluded that coral reef ecosystems appear
23 to be more resistant to macroalgal blooms than assumed (Bruno *et al.*, 2009).

24 25 26 **5.3.2. Impacts on Human Systems**

27
28 Coasts are complex, linked social-ecological systems where anthropogenic alteration has modified the natural
29 processes to the extent that the system dynamics are difficult to separate in terms of human effects and natural
30 processes (Kittinger and Ayers, 2010). The following sections assess understanding of observed weather and climate
31 impacts on human systems on the coasts, with the severity depending on their exposure and vulnerability.

32 33 34 **5.3.2.1. Human Settlements**

35
36 Aspects of weather and climate, such as storms, heat waves, riverine and flash flooding, storm surge inundation and
37 erosion pose a range of hazards for coastal settlements (Handmer *et al.*, 2012). Settlements can amplify the impacts
38 of weather and climate extremes. The urban heat island effect caused by the large amounts of heat absorbing
39 materials used in settlements, and heat emitted from energy and vehicle usage increases daytime maximum and
40 reduces nighttime minimum temperatures. Heat waves can also reduce air quality and increase the number of days in
41 which pollution levels are elevated (Wilby, 2007; Handmer *et al.*, 2012). Most significant for coastal settlements is
42 the potential from flooding from extreme sea levels, which in recent decades have increased at rates similar to mean
43 sea levels rise (Menendez and Woodworth, 2010). Excessive hard paving associated with settlements increases
44 runoff concentration while increased occupation of flood plains for settlements reduces floodplain storage. These
45 can amplify the impacts of flooding as well as increase the risks for those settled in vulnerable areas. Problems of
46 coastal flooding are further enhanced in areas experiencing subsidence due to compaction and groundwater
47 extraction. Settled areas adjacent to the shore are also vulnerable to loss of land through erosion.

48
49 Globally, the Low Elevation Coastal zone (LECZ) of less than 10 m above sea level constitutes 2% of world's land
50 area but contains 10% of world's population (600 million) and 13% of world's urban population (360 million) based
51 on 2000 estimates. About 65% of the worlds cities with populations of over 5 million are in this zone including a
52 disproportionate number of small island states and densely populated megadeltas (McGranahan *et al.*, 2007). Urban
53 poor in informal settlements, of whom there are about 1 billion worldwide, are particularly vulnerable to weather

1 and climate impacts (Handmer *et al.*, 2012). Of the top ten nations classified by population and proportion of
2 population in coastal low-lying areas the majority are developing countries (Table 5-2).
3

4 [INSERT TABLE 5-2 HERE

5 Table 5-2: Top ten nations with the largest populations and the highest proportions of population in low-lying
6 coastal areas (Countries with fewer than 100,000 inhabitants are not included; 15 SIDs (Small Island Developing
7 States) with total of 423,000 inhabitants are also excluded). Source: Bollman *et al.*, 2010.]
8

9 Since the AR4, a large number of coastal assessments have been undertaken that consider exposure to submergence
10 from increasing sea levels or flooding from extreme sea level events ranging from global to local scale (see Table 5-
11 4). In general these studies find current exposure to be largest in the populous Asian cities. Population growth,
12 socio-economic growth and urbanization are the most important drivers of increased exposure of these cities,
13 although climate change and subsidence exacerbate the effect in Asia, particularly in the megadeltas. In Bangkok
14 subsidence has trebled flood damage costs. The Pearl River and Mekong deltas are particularly vulnerable to
15 subsidence as result of land compaction or extraction of groundwater. As urban population represents increasing
16 proportion of world populations, urban floods will account for an increasing percentage of total flood impact as seen
17 in Pakistan, Australia and Brazil (Jha *et al.*, 2011).
18

19 To summarise, there is *very high confidence* that coastal settlements currently exhibit high exposure to weather and
20 climate particularly through sea level extremes (high agreement, robust evidence). The increasing trend of
21 urbanisation over recent years is contributing to this exposure.
22
23

24 5.3.2.2. *Industry, Transport, and Infrastructures*

25
26 Coastal industries, their supporting infrastructure including transport (ports, roads, rail, airports), power and water
27 supply, storm water and sewerage are sensitive to a range of extreme weather and climate events, many of which are
28 projected to change over coming decades (Seneviratne *et al.*, 2012). Severe storms with associated winds, rain,
29 lightning and storm surges are particularly disruptive to transport and power and water supplies. Heat extremes
30 affect asphalt road surfaces and cause movement of materials used in infrastructure (e.g., buckling of rail tracks).
31 Droughts affects water quality and supply as does increasing salinisation of coastal aquifers due to rising sea levels
32 or extreme sea level events (Handmer *et al.*, 2012).
33

34 Sea transport accounts for more than 80% of global goods trade (by volume) and so disruption to port activities in
35 one location can disrupt supply chains, which can have far reaching consequences (Handmer *et al.*, 2012). Increased
36 sediment mobility and changes in sedimentation/erosion patterns restrict operations of harbours and access channels
37 (UNCTAD, 2008). Coastal industries in high latitudes are affected by permafrost thaw causing ground stability and
38 erosion thereby affecting transport safety and the industries that rely on such travel in these regions (e.g., Pearce *et al.*, 2010).
39
40

41 For certain coastal environments such as barriers, climate-related modification of roads brings significant changes
42 including coastal squeeze. To summarise, climate impacts on coastal industries vary considerably depending on
43 geographical location, associated weather and climate and specific composition of industries within particular
44 coastal regions.
45
46

47 5.3.2.3. *Fisheries, Aquaculture, and Agriculture*

48
49 Fisheries constitute one of the most important economic sectors in the coastal region. Fisheries and aquaculture
50 industries are estimated to employ 43.5 million people globally with the majority in developing countries (Daw *et al.*, 2009). Nearly 1.5 billion people worldwide rely on fish for more than 20% of their dietary animal protein
51 (Badjeck *et al.*, 2010) and many small island states rely on fisheries and aquaculture for 50% of their animal protein
52 (Barange and Perry, 2009). The coastal zone also supports significant agricultural food production. For example,
53 rice production is a major crop grown in the low-lying deltaic regions of Asia; 54% and 17% of Vietnam's rice
54

1 production occurs in the Mekong and Red River deltas respectively; 68% and 34% of Myanmar's and Bangladesh's
2 rice production occurs in the Irrawaddy and Ganges-Brahmaputra deltas (Wassman *et al.*, 2009).

3
4 Recent studies have investigated the changes in the regional abundance of fish species. In the North Sea, Hiddink
5 and Hofstede (2008) found that ocean warming over the period 1977-2002 led to 8 times more fish species
6 exhibiting increased distribution than decreased distribution. The species showing increase were of southerly origin
7 while those showing decrease were the larger northerly species. In southeastern Australia, Last *et al.* (2011) also
8 found an increasing abundance of 45 fish species of warm temperate origin, which they linked to the observed
9 strengthening of the East Australian Current (EAC) bringing warm waters further south (Ridgeway, 2007). Long
10 term declines in abundance were found in only 9 species, which were attributed to fishing pressure. They noted
11 however the potential for longer term detrimental effects for cool temperate species endemic to the region whose
12 southward migration would be constrained by the southern limits of the continental shelf.

13
14 Warming ocean temperatures can also lead to the migration of more pest species. For example, in southeastern
15 Australia, the sea urchin, problematic because it can overgraze seaweed beds, has also migrated south under the
16 influence of the warmer EAC (Ling *et al.*, 2009). In coastal Louisiana, saltwater intrusion was found to reduce the
17 population size of the freshwater western mosquitofish (*Gambusia affinis*) although populations from more brackish
18 sites were less affected suggesting some degree of localised adaptation (Purcell *et al.*, 2010). Eutrophication and
19 hypoxia give rise to harmful algal blooms (HABs) and episodes of HABs have increased in frequency and intensity,
20 harming fisheries and human health (MEA, 2005).

21
22 In coastal Bangladesh salinity has become a major problem for traditional agriculture, displacing production of
23 vegetables, fruit and animals and increasing shrimp farming. Once established, shrimp farming further increases the
24 salinity of the local environment rendering it unsuitable for a return to traditional farming and necessitating the
25 cultivation of high yielding, salt-tolerant rice varieties (Rahman *et al.*, 2011). Saltwater inundation events and
26 droughts have reduced the yield or destroyed crops such as taro and breadfruit, particularly in low-lying island
27 nations (e.g. Keener *et al.*, 2012).

28
29 In summary, there is *medium confidence* that changes have occurred to the distribution of fish species with evidence
30 of poleward expansion of temperate species (high agreement, limited evidence). There is limited evidence to suggest
31 that tropical and subtropical aquaculture has not been adversely affected by rising ocean temperatures to date. There
32 is also *medium confidence* that coastal agriculture has experienced negative impacts due mainly to increased
33 frequency of saltwater inundation (high agreement, limited evidence).

34 35 36 5.3.2.4. Coastal Tourism and Recreation

37
38 Coastal tourism is considered the most important and fastest growing tourism sector (UNEP, 2009). Yet, coastal
39 zones and islands as tourism destinations are among the most vulnerable to climate change. For example, the island
40 nations of the Caribbean, Mediterranean, Indian Ocean and Pacific Ocean and Australia/New Zealand comprise five
41 of eleven most at-risk tourism destinations (WTO, 2007). Significant impacts are occurring from changes in extreme
42 events (e.g. flooding, tropical storms, storm surges, heat waves) and climate variability (e.g. drought and prevailing
43 winds accelerating coastal erosion) (Scott *et al.*, 2008).

44
45 Impacts are particularly relevant for tropical island coasts that depend on coral reefs for tourism where coral
46 bleaching events observed since 1980 due to SST increase are a major concern. Globally, tourism and recreation
47 accounted for \$9.6 billion net benefits of coral reefs. In the Caribbean alone, the net benefits from dive tourism are
48 estimated to be \$2.1 billion. Australia's Great Barrier Reefs has a recreational use value of \$700 million to \$1.6
49 billion per year (Conservation International, 2008).

50
51 In summary, coastal tourism is highly vulnerable to weather and climate extremes and rising sea-levels (*high*
52 *confidence*) with the additional sensitivity to ocean temperature and chemistry changes for the sectors that rely on
53 reef tourism.

5.3.2.5. *Water Resources*

Coastal aquifers are sensitive to a range of climate-related factors such as temperature and evaporation rise, rainfall decline and/or changes in rainfall intensity that affect aquifer recharge, and rising sea levels and storm surges that cause salt water intrusion into aquifers. Combined with this excessive groundwater extraction for coastal settlements and agriculture causes intrusion of saline water laterally from the ocean or from deeper, more saline layers below the aquifer (the pathway depends on the particular hydrogeological characteristics). Intrusion can also occur as a result of poorly constructed or abandoned wells, open boreholes and dredged channels (Barlow and Reichard, 2011). However, Werner (2010) noted that there are fundamental knowledge gaps that impede assessment of changes in seawater intrusion due to land and water resource management changes. This will impede the assessment of climate change impacts that may arise from changes in ground water recharge and sea level rise.

Many parts of the world where groundwater is used for human consumption or agriculture are experiencing changes in water quality such as salinisation (e.g., Custodio, 2010; Werner, 2010; Steyl and Dennis, 2010; Bocanegra *et al.*, 2010; Barlow and Reichard, 2010; Essink *et al.*, 2010). For example, coastal aquifers on the east and west coasts of the US have experienced increased levels of salinity largely due to excessive anthropogenic water extraction (Barlow and Reichard, 2011). In the semi-arid Chaouia Coast region of Morocco, about half of the wells that commenced pumping in the 1960s have ceased to be operational because of water shortage (84%) or salinity (16%). Modelling indicates that despite rainfall declines since 1977, the reduction in water resources is less sensitive to the drought conditions than it is to the intensive and uncontrolled water pumping for agriculture over this time (Moustadraf *et al.*, 2008). In rainfall intensive Taiwan, about 34% of annual water is sourced from groundwater, but over exploitation has led to a lowering of groundwater levels, seawater intrusion, salinization of soil and reduced well yields (Hsu *et al.*, 2007).

Around 8,000 inhabited small tropical islands in the Pacific, Indian and Atlantic Oceans face water supply problems. Many rely on extraction of groundwater from thin freshwater lenses in highly permeable aquifers (Terry and Falkland, 2010). Droughts, storm surges and sea level variations, which are strongly linked to natural modes of variability such as ENSO strongly influence the groundwater supply. For example, saline intrusion into freshwater aquifers following cyclone-induced overwashes can take many months to resolve (Terry and Falkland, 2010). The combination of natural with anthropogenic drivers of over-extraction, pollution from human, animal and industrial sources, mining of sand and gravel for building and erosion due to shoreline works compound groundwater supply problems in these locations (White *et al.*, 2007; White and Falkland, 2010).

In summary, there is *very high confidence* that increased usage of groundwater resources for agriculture and coastal settlements globally has led to a reduction in groundwater quality, including increased salinization (high agreement, much evidence). There is *high confidence* that climate change through changing patterns of precipitation and sea level rise will exacerbate these problems.

5.3.2.6. *Health*

In coastal regions, climate may impact human health directly through the occurrence of events such as floods, droughts, storm surges and extreme temperatures and indirectly through changes in the transmission of vector, food and water borne infectious diseases. Diseases associated with air pollutants and aeroallergens can also be affected by climate (Ebi *et al.*, 2008). Understanding the relationship between climate and health is often confounded by socio-economic factors that influence coastal settlement patterns and the level and organisation of the response of authorities to health related issues (Baulcomb, 2011).

Flood mortality risk has fallen since 1980 in all regions apart from South Asia and cyclone mortality risk has fallen in all regions since 2000 and is now lower than in 1980. Since exposure has increased over the same period, this result is significant and reflects how development has reduced vulnerability and strengthened countries capacity to respond to disasters. However, mortality is still rising in the countries with the weakest risk governance capacities (UNISDR, 2011). Heat extremes can also directly affect health outcomes in coastal cities, even those of tropical

1 countries that are acclimatised to high temperatures (McMichael *et al.*, 2008). Health impacts affect low income
2 groups and countries more severely than high income countries (Handmer *et al.*, 2012)

3
4 Vector-borne diseases such as malaria, dengue and leishmaniasis are sensitive to increasing temperature, humidity
5 and rainfall, which can increase suitable mosquito habitats, shorten the breeding cycles and reduce the time to
6 infection (Stratten *et al.*, 2008; van Kleef *et al.*, 2010). For example, dengue risk in Hawaii contracts under El Nino-
7 induced droughts and increases with increased precipitation during La Nina (Kolivras, 2010). Although dengue has
8 been reported a few new locations, its distribution generally contracted over the past century, which probably relates
9 to changes in housing conditions and methods of water supply (Van Kleef *et al.*, 2010). On the other hand, despite
10 efforts to eradicate malaria from the 1950's to 1980's the death rate from malaria is higher today than 40 years ago
11 (Stratten *et al.*, 2008). Jackson *et al.* (2010) found minimal correlation reported malaria rates and climate in west
12 Africa and suggested that climate linkages may be more complex than those they considered. In Bangladesh, more
13 visceral leishmaniasis cases are found near flood control embankments, implying that building more embankments
14 in response to sea-level rise may increase the vectors and incidence of this disease (Shahid, 2009).

15
16 Food and water borne infectious diseases include viral, bacterial or parasitic pathogens that are transmitted to
17 humans through ingestion or inhalation of contaminated food and water. A link between infectious diseases and
18 temperature has been reported in several recent studies including diarrhea in Taiwan (Chou *et al.*, 2011), non-
19 cholera diarrhoea in Bangladesh (Hashizume *et al.*, 2007), salmonella infection in tropical and temperate coastal
20 cities in Australia (Zhang *et al.*, 2010), bacillary dysentery in China (Zhang *et al.*, 2007), infectious gastrointestinal
21 disease in Japan (Onozuka *et al.*, 2010) and rotovirus in Bangladesh (Hashizume *et al.*, 2008). Precipitation,
22 humidity and, in the rotovirus case, river levels also affected incidence with relationships that differed across
23 studies. Schistosomiasis, a parasitic disease in which free swimming larvae of a parasitic worm penetrate human
24 flesh for reproduction, represents a significant health burden to the 76 countries in which it is endemic (Yang, 2006).
25 Mangal *et al.* (2008) found that human infections increase linearly with temperature to 30°C but beyond 35°C, the
26 infection rate drops off because of increased mortality in the snail intermediate host.

27
28 Climate variations potentially provide a strong link to outbreaks of disease due to aquatic bacteria such as cholera
29 (Costello *et al.*, 2010) and toxins generated by Harmful Algal Blooms (HABs) such as ciguatera. An increase in both
30 the extent and the frequency of HABs has been reported in the eastern North Sea due to increases in sea surface
31 temperature and decreased salinity (Jaykus *et al.*, 2008). An indirect link through increased sea levels, increased
32 precipitation and flash floods may also contribute to the occurrence of HABs by increasing nutrient rich water
33 release to coastal and marine waters (Jaykus *et al.*, 2008; Erdner *et al.*, 2011). Nontoxic blooms that grow to high
34 biomass also can have detrimental effects on biodiversity through oxygen depletion and physical shading of the
35 benthos (Erdner *et al.*, 2011) with flow on effects to ecosystem health, human nutrition and health.

36
37 In summary, there is clear evidence of relationships between climate and many diseases that affect human health
38 mainly through increases in temperature, rainfall and humidity. However, the relationships are complex and vary
39 between diseases and even regionally for the same disease (high agreement, robust evidence).

40 41 42 **5.4. Projected Impacts**

43 44 **5.4.1. Impacts on Habitats and Ecosystems**

45
46 Coastal ecosystems are subject to both land- and ocean-based anthropogenic drivers and are experiencing some of
47 the greatest cumulative impact of human activities (Halpern *et al.*, 2008). Anthropogenic drivers associated with
48 global climate change are distributed widely and are an important component of global cumulative impacts. There
49 are few coral reefs, seagrass beds, mangroves, rocky reefs and shelves with limited impact. Lotze *et al.* (2006)
50 argued that overexploitation and habitat destruction have been responsible for most of the historical changes that
51 occurred in coastal systems and that eutrophication, although severe in the last phase of estuarine history, largely
52 followed rather than drove observed declines in diversity, structure, and functioning. Further, extreme climate events
53 produce simultaneous changes to the mean and to the variance of climatic variables over ecological time scales.
54 Here we discuss projected physical and biological impacts to specific coastal habitats and ecosystems.

5.4.1.1. Rocky Shores

Rocky shores composed of both soft (readily-erodible) and hard (relatively resistant) material will likely erode more rapidly with rising sea level (Trenhaile, 2011). For shores composed of resistant materials, the rates will be relatively low. Beaches that lie seaward of rocky shores will be squeezed, particularly on resistant rocky coasts, and lost, leaving the rocky habitats exposed to ocean forces. In a feedback, eroding soft rocks in bluffs composed of loosely-consolidated sand deposits will release sand that may naturally nourish the ocean fronting beach and help maintain the beach and protect the cliff.

The abundance and distribution of rocky-shore species have long been recognized to vary, and will continue to do so in a warming world. The number of species, notably calcifiers, found in rocky shore stations located near natural CO₂ vents at Ischia (Mediterranean Sea) is 30% lower at pH levels close to those expected in 2100 under the IS92a CO₂ emissions scenario (Barry *et al.*, 2011). However, the temporal variability in pH may have contributed to the pronounced biodiversity shifts observed, as these stations experienced short periods of pH_T as low as 7.4-7.5. Model projections that assume a linear increase in temperature and ocean acidity, and include the interactive effects of temperature and pH suggests that a local population of the barnacle *S. balanoides* will become extinct in the Channel ten years earlier than would occur if there was only global warming and no concomitant decrease in pH (Findlay *et al.*, 2010).

5.4.1.2. Beaches and Sand Dunes

Climate change has undoubtedly contributed to the observed erosion impacts found along the world's beaches and dunes (See 5.3), but there are other processes, unrelated to climate change, that contribute as well, such as dams capturing fluvial sand (e.g., in Morocco, Chaibi and Sedrati, 2009). Definitively linking sea-level rise to observed magnitudes of beach erosion has been challenging (e.g. see Sallenger *et al.*, 2000), although recent results using a Bayesian network has been successful in hindcasting sea-level rise induced shoreline change (Gutierrez *et al.*, 2011). With projected sea-level rise approaching 1 m or more over the next century, the resulting inundation and erosion should become readily detectable and progressively important. The impacts will be first apparent by sea-level rise adding to storm surge, making wave runup higher, and more frequent on beaches and dunes. For example, Tebaldi *et al.* (2012) showed how projected sea level rise shortens return periods of extreme storm surges, potentially reducing their return periods from centuries to decades. Sea-level rise will not only inundate low-lying land, like filling a bath tub, but can contribute to dynamic changes in sandy beaches, through shifting the beach position landward and higher. How to calculate the amount of retreat is controversial (e.g., see recent attempt by Ranasinghe *et al.*, 2012 and the Bayesian network forecast approach by Gutierrez *et al.*, 2011). Scientists disagree on whether tropical cyclones will become more intense and/or frequent in the future, although recent, as of yet non-conclusive, evidence suggests extra-tropical storm tracks may change in a warmer future extending Atlantic winter storms eastward impacting Europe to a greater degree than present (Schwierz *et al.*, 2010; Woolings *et al.*, 2012.). Such increase in storminess could accelerate erosion of beaches and dunes. Coastal squeeze will accelerate with rising sea level. Finding, and funding, sufficient sand to artificially rebuild beaches and dunes will likely become increasingly difficult.

5.4.1.3. Estuaries, Tidal Flats, and Lagoons

Sea-level rise will have consequences for the partitioning of habitats within estuaries and for the landward extension of estuaries. Global warming has consequences for the physics, chemistry and biology of estuaries. Most of the time stratification, dominated by salinity, is a natural process, but long-term global warming, climate-related precipitation changes and altered riverine discharge may increase the extent, duration and frequency of estuarine stratification with consequences for ecosystem metabolism, biogeochemical processes and organism distribution patterns. For instance increasing persistence of stratification in the estuarine plume of the Mississippi river will lead to more increasing hypoxia (Rabalais *et al.*, 2009).

1
2 Anthony *et al.* (2010) projected that climate change will generate sediment redistribution as well as increased
3 erosion and shoreward migration of barriers in coastal lagoons. The flushing rate, which is a key parameter
4 controlling biogeochemical processes such as primary production (Smith *et al.*, 2005b; Webster and Harris, 2004),
5 could either increase due to barrier breaching or lower freshwater supply or decrease if the input of freshwater
6 decreases (Anthony *et al.*, 2010).
7

8 The loss of benthic macrophytes is projected in some lagoons due to increased mortality and decreased net primary
9 production driven by increased temperature and lower light availability resulting from sea level rise and increased
10 inputs of nutrients and suspended solids (Lloret *et al.*, 2008). Since benthic macrophytes play a key role to intercept
11 and store nutrients (Grall and Chauvaud, 2002), their demise could increase the occurrence and magnitude of
12 eutrophication (Lloret *et al.*, 2008).
13
14

15 5.4.1.4. Deltas

16

17 Deltas are influenced by climatic drivers from drainage basins, e.g., changes of precipitation, and from oceans, e.g.,
18 sea-level rise and changes of waves and storm surges, and also non-climatic drivers, e.g., changes of water and
19 sediment discharges, subsidence, land-use change. Projected impacts on deltas are caused mainly by fluvial floods,
20 future sea-level rise, changes of waves and storm surges, resulting in increased coastal flooding, wetland decrease,
21 increased coastal erosion and increased salinization of cultivated land and ground water (McLeod *et al.*, 2010; Day
22 *et al.*, 2011). As non-climatic drivers, e.g., reduction in sediment delivery, subsidence, and land-use changes, have
23 impacted deltas for the last 50 years more than climatic drivers (Syvitski, 2008), combined impacts of both drivers
24 on deltas will be projected. Flooding of surface areas of 33 deltas chosen to represent global delta variability will
25 increase by 50 % for the future sea-level rise projected for 2100 using the IPCC AR4 scenario (Syvitski *et al.*, 2009).
26 Future sea-level rise and storm surges by potentially stronger tropical cyclones will present a threat to deltas,
27 particularly densely populated megadeltas in Asia (Murray *et al.*, 2012).
28
29

30 5.4.1.5. Mangroves and Salt Marshes

31

32 Sea-level rise may be problematic for mangrove systems in case mangrove-derived peat accumulation and/or
33 sediment supply and thus accumulation cannot keep pace with sea-level rise and drowning will occur. Geological
34 record shows that these systems migrate landwards during transgressions.
35

36 Global warming will have effect on the geographical distribution patterns of salt marshes, with likely increases at
37 high latitudes and decreases at lower latitudes, but this is rather uncertain at the moment (Bromberg Gedan *et al.*,
38 2009). Salt marsh plant may become more productive at temperature rises but respiration losses also increase by
39 about 20% (Kirwan and Blum, 2011). The balance between increase in production due to temperature and carbon
40 dioxide increases and increase in respiration due to elevated temperature appears to be in favour of mineralization
41 processes, suggesting that coastal marshes in a high carbon dioxide, high temperature world would be less resilient
42 to sea-level rise (Kirwan and Blum, 2011).
43

44 Submergence-accretion and productivity-submergence feedbacks couple rates of accretion to sea-level rise and may
45 limit drowning of marshes due to accelerated sea-level rise (Kirwan and Temmerman, 2009; Bromberg Gedan *et al.*,
46 2011).
47

48 The direct effect of atmospheric carbon dioxide increase on saltmarshes will be differential depending on whether
49 C3 (Phragmites) or C4 (e.g. Spartina) plants dominate, because the latter are usually rather insensitive to direct CO₂
50 effects (Rozema *et al.*, 1991).
51
52
53

5.4.1.6. Coral Reefs

Considering the maximum rate of vertical accretion recorded during the last deglaciation (about 20 mm y⁻¹; Dullo, 2005; Montaggioni *et al.*, 2005), modern reefs appear to be able to keep up with the present rate of sea level rise but this may be compromised with a decreased rate of accretion driven by climatic and non-climatic drivers. A 0.5 to 1.0 m rise in sea level would increase coastal erosion, sediment resuspension, and the duration of high turbidity events on exposed reef flats of Molokai, Hawaii, leading to decreased light availability for photosynthesis and increased sediment-induced stress (Storlazzi *et al.*, 2011). Model results suggest that the thermal tolerance of reef-building corals are likely to be exceeded every year within the next few decades (Hoegh-Guldberg, 1999). The combination of ocean acidification and temperature being synergistic in several reef-builders (Reynaud *et al.*, 2003; Anthony *et al.*, 2008), it is very likely that the impacts will be more dramatic in the next decades than observed today near CO₂ vents, which do not simulate future warming (Fabricius *et al.*, 2011). Under the A1B CO₂ emission scenario, frequency and intensity of bleaching events will increase considerably during the period 2000-2100, with 99% of the reef cells experiencing at least one severe bleaching event in 2090-2099 (Figure 5-6; Teneva *et al.*, 2011; see also Meissner *et al.*, 2012). There is a high level of confidence that a large decline of coral cover will occur in the Hawaiian Archipelago during the 21st century (A1B CO₂ emission scenario; Hoeke *et al.*, 2011). A global model suggests that all coral reefs will stop growing and start dissolving when atmospheric CO₂ reach 560 ppm (Silverman *et al.*, 2009).

[INSERT FIGURE 5-6 HERE

Figure 5-6: Percent of reef locations (1°x1° latitude/longitude cells which have coral reefs) that experience no bleaching (green), at least one mild bleaching event (reddish-brown), or at least one severe bleaching event (purple) for each decade. Observed bleaching events are summarized from the ReefBase dataset (Kleypas *et al.* 2008). In the observations, some of the “no bleaching” cells may have experienced bleaching but it was either not observed or not reported. Modeled bleaching events are averages of data from four ensemble runs of the Community Climate System Model version 3 using the standard degree heating month formula (Teneva *et al.*, 2011). The labels of values ≤ 1% are not shown.]

5.4.1.7. Seagrasses and Algae

Based on current observations indicating that seagrass meadows are already under stress due to realised climate change, it is projected that seagrass meadows will experience a decline with further warming (e.g., Marbà and Duarte, 2010, Rasheed and Unsworth, 2011).

Seagrass meadows may, however, expand their poleward ranges with warming, particularly towards the Arctic, along the coasts of Greenland, Norway, Siberia and North America. However, a lack of reports on the dynamics of seagrass meadows at high latitudes precludes the assessment of whether the expected poleward expansion is already occurring (Duarte, 2000).

Increased CO₂ is expected to increase seagrass photosynthetic rates (Hemminga and Duarte, 2000; Hendriks *et al.*, 2010).

Macroalgae in the north temperate zone are expected to extend their distribution into the High Arctic towards the end of the 21st century, but retreat along the northeastern Atlantic coastline (Müller *et al.*, 2009), whereas Antarctic seaweeds are not expected to alter their distribution substantially (Müller *et al.*, 2009). However, range shifts of macroalgae may be slow (Hinz *et al.*, 2011) and poleward shifts have been documented for warm-water species rather than for cold-water ones (Lima *et al.*, 2007). Hence, the expectation of poleward range shifts of macroalgae due to increasing temperature should be considered with caution as it does not seem to be a universal process (Lima *et al.*, 2007).

Macroalgae are, in general, not expected to be negatively affected by ocean acidification (Hendriks *et al.*, 2010). However, calcifying macroalgal species may be affected by ocean acidification, as macroalgae calcification rates

1 have been shown to be inhibited by elevated CO₂ concentrations (Gao *et al.*, 1993; Kuffner *et al.*, 2008).
2 Examination of community structure along volcanic areas, naturally enhanced in CO₂ suggests that turf algae may
3 be impacted by the acidification levels expected by 2100 (Porzio *et al.*, 2011), and research on coral reefs along
4 naturally CO₂ enriched reefs near volcanic areas suggests that macroalgal cover increases at high CO₂ (Fabricius *et*
5 *al.*, 2011).
6
7

8 **5.4.2. Impacts on Human Systems**

9

10 **5.4.2.1. Human Settlements**

11

12 The most important effects of climate change on the coastal cities include the effects of sea-level rise, effects of
13 extreme events on built infrastructure (such as wind storms, storm surges, floods, heat extremes and droughts),
14 effects on health, food and water-borne disease, effects on energy use, and effects on water availability and
15 resources (Hunt and Watkiss, 2010). Considering that projected changes up to 2100 indicate that it is very likely that
16 mean sea level rise will contribute to upward trends in extreme coastal high water levels, there is a high confidence
17 that human settlements currently experiencing coastal inundation and erosion, will continue to do so in the absence
18 of changes in other contributing factors.
19

20 An assessment of coastal flooding on 136 port cities around the world each with >1 m inhabitants in 2005 indicated
21 40 million inhabitants to be exposed to a 1 in 100 year coastal flood event. By 2070 this would trebled to 150
22 million. The top 10 exposed cities in terms of exposed population are Mumbai, Guangzhou, Shanghai, Miami, Ho
23 Chi Minh City, Kolkata, Greater New York, Osaka-Kobe, Alexandria and New Orleans, almost equally split
24 between the developed and developing countries. In terms of assets exposed, 60% are from the USA, Japan and the
25 Netherlands. The total assets exposed in 2005 across all cities are estimated to be US\$3,000 billion, which would
26 increase to US\$35,000 billion by 2070s (Nicholls *et al.*, 2008; Hanson *et al.*, 2011).
27

28 Land subsidence is greater than the effect of sea-level rise in a number of Asian coastal cities. For example, parts of
29 Jakarta are subjected to regular flooding on a near-monthly basis. Under current conditions, the estimated damage
30 by extreme coastal flood events with return periods of 100 and 1000 years is €4 billion and €5.2 billion
31 respectively. Under a scenario for 2100, damage is increased by a factor of 4-5 (Ward *et al.*, 2011). Semarang, 400
32 km east of Jakarta, is already subject to coastal hazards due to tidal inundation and land subsidence. With a scenario
33 of 1.2 m inundation, nearly 4600 ha would be affected at a cost of €1.8 billion (Marfai and King, 2008).
34

35 Based on the assumption that projected increase in extreme sea levels is largely due to an increase in MSL, Hunter
36 (2010) described a method of combining observations of present sea level extremes with the projections of sea level
37 rise to obtain a “sea level allowance.” This allowance is calculated so that the expected frequency of flooding events
38 is preserved. It is based on the projected rise in mean sea level and its uncertainty, and on the variability of tides and
39 storm surges (which are parameterised by the scale parameter of their Gumbel distribution). The method was applied
40 to 198 tide gauge stations over the globe, yielding estimates of the scale parameter (a measure of the variability of
41 high sea levels), which varied between 0.05 and 0.20 m for 90% of the stations considered (Hunter, 2010).
42
43

44 **5.4.2.2. Industry, Transport, and Infrastructures**

45

46 Climate change and especially sea-level rise may impact critical infrastructures located at the coastal area.
47

48 Transportation facilities serve as the lifeline to communities. Sea-level rise poses a risk to transportation in ensuring
49 reliable and sustained transportation services since due to the network configuration, inundation of even the smallest
50 component of an intermodal system can result in a much larger system disruption. For instance, even though a
51 transportation terminal may not be affected, the access roads to it could be, thus forcing the terminal to cease or
52 reduce operation (CCSP, 2008). Some low-lying railroads, tunnels, ports and roads are already vulnerable to
53 flooding and a rising sea level will only exacerbate the situation by causing more frequent and more serious
54 disruption of transportation services. Furthermore, sea-level rise will reduce the extreme flood return periods and

1 will lower the design critical elevations of infrastructure such as airports, tunnels, and ship terminals (Jacob *et al.*,
2 2007). For example, it is estimated that a hypothetical 1 m rise in relative sea level projected for the Gulf Coast
3 region between Alabama and Houston over the next 50-100 years would permanently flood a third of the region's
4 roads as well as putting more than 70% of the regions ports at risk (CCSP, 2008).

5
6 Although not completely coastal, the estimated costs of climate change to Alaska's public infrastructure could add
7 US\$3.6-6.1 billion (+10% to 20% above normal wear and tear) from now to 2030 and US\$5.6-7.6 billion (+10% to
8 12%) from now to 2080 (Larsen *et al.*, 2008). Higher costs of climate change for coastal infrastructure are expected
9 due to its proximity to the marine environment.

10
11 Other projected impacts are beneficial for the transportation system. For example, decline of Arctic sea-ice coverage
12 could extend seasonal accessibility to high-latitude shipping routes such as the northwest shipping route that
13 connects the Atlantic to the North Pacific (MCCIP, 2008).

14
15 Coastal infrastructural instability may result from natural hazards triggered by groundwater-level (GWL) variations
16 resulting from rising sea level. For earthquake-prone coasts, this could be exacerbated by earthquake liquefaction if
17 GWL increases with sea-level rise (Yasuhara *et al.*, 2007).

18
19 Transportation infrastructure is not the only sector affected. There are several lifeline, infrastructures and industry
20 facilities traditionally located at or close to the shoreline that play a very relevant role to the human system. A
21 number of these existing facilities are located at lower elevations and if extreme climate events (storm surges,
22 extreme winds and waves) become more frequent and intense, there will be increased stress on all of these
23 infrastructure systems (Zimmerman and Faris 2010). Table 5-3 summarizes potential impacts of sea level rise,
24 coastal floods and storms on critical coastal infrastructure in the communications, energy, transportation and water
25 waste sectors.

26
27 [INSERT TABLE 5-3 HERE

28 Table 5-3: Impacts of sea level rise, coastal floods, and storms on critical coastal infrastructure by sector.]

29 30 31 5.4.2.3. *Fisheries, Aquaculture, and Agriculture*

32
33 The potential for climate change to affect food security in the coastal zone may arise through one or more climate
34 drivers such as sea-level rise, increases in ocean temperature, acidity and changes in rainfall patterns. The impact of
35 such drivers will be enhanced where other non-climate drivers such as pollution and poor management practices
36 occur. For aquaculture, it is anticipated that negative impacts of rising ocean temperatures will be felt in the
37 temperate regions due to exceedance of the optimal temperature range of organisms currently cultured in this region
38 whereas positive impacts will be felt in the tropical and subtropical regions (De Silva and Soto, 2009).

39 40 41 5.4.2.4. *Coastal Tourism and Recreation*

42
43 Globally, coastal tourist destinations are affected by sea-level rise (coastal erosion), sea surface temperature and
44 acidification (coral reefs), and increased frequency and intensity of tropical storms (damage to infrastructure and
45 tourist attractions). Future sea-level rise aggravates coastal erosion leading to the loss of beaches. Even small rises in
46 sea level could result in significant erosion and submergence of land, increased floods, contamination of freshwater
47 aquifers, loss of protective coral reefs, mangroves and beaches which will increase exposure to extreme weather
48 events including tourism destinations which will reduce amenity (Scott *et al.*, 2008).

49
50 The Caribbean with many high-dependency tourism islands would be impacted by climate change and sea-level rise.
51 St. Kitts and Nevis, Antigua and Barbuda, Barbados, St. Vincent and the Grenadines and Grenada are particularly
52 affected with high annual costs due to degrading beach assets and inundation. The estimated capital costs to rebuild
53 tourist resorts are US\$10-23 billion in 2050 and US\$24.5-73.9 billion in 2080. A hypothetical 1-m sea-level rise
54 would result in the loss or damage of 21 CARICOM airports, inundation of land surrounding 35 ports, and at least

1 307 multi-million dollar tourism resorts damaged or lost from erosion to the coastal beach areas (Simpson *et al.*,
2 2011).

3
4 For tropical islands and coasts dependent on corals for tourism, there has been a concern about coral bleaching and,
5 in recent years, about the impacts of acidification. While large communities of coral reefs may persist under
6 scenarios of atmospheric carbon dioxide at 375 ppm (+1°C) and 450-500 ppm (+2°C), will degrade carbonate reef
7 structures under a scenario of >500 ppm (+3°C) with serious consequences for tourism destinations in Australia,
8 Caribbean and other small island nations (Hoegh-Gulberg *et al.*, 2007).

9
10 Another result of climate change on coastal tourism would be the coastal squeeze exacerbated by coastal
11 construction and tourist hotels built within the zone at risk to flooding and erosion (Schleupner, 2008). Dykes, as an
12 adaptation measure for sea-level rise have a negative impacts on tourist coasts as shown in the coastal districts of
13 Schleswig-Holstein, Germany, where an increase in the length of dikes resulted in the reduction in the average price
14 of accommodation.

15 16 17 5.4.2.5. *Water Resources*

18
19 Salinization is often considered the major impact on coastal aquifers as a result of sea-level rise. A typical situation
20 is where coastal geological conditions favoured saltwater intrusion into the recharge zone. For example, Guyana
21 with more than 90% of its ¾ million population in a narrow coastal strip less than 10 km from the coast is protected
22 by a system of dikes constructed mainly of concrete and clay. The flat coastal plains extend more than 40 km inland
23 and, under a hypothetical 1 m sea level rise, saltwater is estimated to intrude into the recharge zones for a distance of
24 1-12.5 km. Water extraction will exacerbate the intrusion (Narayan, 2006).

25 26 27 5.4.2.6. *Health*

28
29 The expansion of brackish and saline water bodies in the coastal areas associated with rising sea levels is now
30 recognized as a potential health hazard to coastal communities (Ramasamy and Surendran, 2011). In Bangladesh
31 increased salinity in drinking water will increase the risk of diarrhea and skin diseases. Inland intrusion of saltwater
32 may turn former freshwater habitats into saltmarsh areas acting as breeding ground of saltmarsh mosquitoes and
33 increase vector-borne diseases in the coastal areas of the country. The construction of embankments as a response to
34 sea-level rise, may favour visceral leishmaniasis vectors and result in increased cases of visceral leishmaniasis
35 (Sahid, 2009).

36
37 Kolstad and Johansson (2011) contend that sparsity of empirical climate–health data is a major contributing factor to
38 uncertainties in projections of future climate change on health.

39 40 41 5.5. **Assessing Vulnerabilities, Risks, and Costs**

42 43 5.5.1. *Approaches*

44
45 Vulnerability and risk are ambiguously defined concepts and a wide variety of methods are applied for assessing
46 these (e.g., Adger, 2006; Hinkel, 2011). When following the IPCC definition of vulnerability, assessments of
47 vulnerability ought to go beyond assessments of potential impacts in that they also include information on
48 adaptation. This can either be done explicitly by including adaptation within an impact model or implicitly by using
49 generic indicators of adaptive capacity. Roughly two interpretations of risk can be distinguished in the literature.
50 The first one refers to risks of gradual changes of mean climate variables (e.g., global mean sea-level rise). Under
51 this interpretation, approaches applied for assessing risks are identical to those applied for assessing vulnerability.
52 The second interpretation refers to extreme events and interprets risk as a measure of the probability of occurrence
53 of extreme events and their consequences (i.e., damages). Under this second interpretation, coastal assessments

1 focus on flood risk. In summary, the following seven types of approaches for assessing coastal vulnerability and risk
2 can be distinguished in the literature:

- 3 - *Submergence exposure approaches* that use GIS to assess the exposure (in terms of people, assets,
4 ecosystems) to permanent inundation under a given level of global mean sea-level rise (e.g., Dasgupta *et al.*,
5 2008; Boateng, 2012).
- 6 - *Flood exposure approaches* that use GIS to assess the exposure to temporary inundation during a coastal
7 flood event, either assuming present sea-levels or future ones by combining the extreme water level of the
8 flood event with a given level of global mean sea-level rise (e.g., Dasgupta *et al.*, 2009; Kebede and
9 Nicholls, 2012).
- 10 - *Indicator-based approaches* that aggregate data on the current state of the coastal systems into vulnerability
11 indices. The indicating variables used can either be biophysical (e.g., Yin *et al.*, 2012; Bosom and Jimenez,
12 2011), socioeconomic (Cinner *et al.*, 2012) or both (e.g., Bjarnadottir *et al.*, 2011; Yoo *et al.*, 2012; Li and
13 Li, 2012). The socioeconomic variables used refer to the social system's current and usually generic (i.e.,
14 not impact specific) capacity to adapt.
- 15 - *Impact model-based approaches* that include adaptation explicitly in computational models and simulate
16 impacts attained over time under various socio-economic and climate scenarios as well as adaptation
17 strategies. Examples of this approach are the applications of the DIVA and FUND models discussed in
18 Section 5.5.3 and 5.5.4.
- 19 - *Hydrodynamic flood damage approaches* that use hydrodynamic models to simulate the consequences of
20 particular extreme water level events (e.g., Xia *et al.*, 2011; Lewis *et al.*, 2011). The probabilities used for
21 calculating risks are either the ones of extreme-water levels or, in the case of existing defences, those of
22 defence failure (e.g., Reeve, 1998; Dawson *et al.*, 2009; Dutta, 2011).
- 23 - *Expected flood damage approaches* that assess current and future flood risks as mathematical expectation
24 using the full distribution of extreme water levels (e.g., the flood component of the DIVA model as done by
25 Hinkel *et al.*, 2010; Hinkel *et al.*, 2011; Hinkel *et al.*, 2012).
- 26 - *Qualitative approaches* that assess coastal vulnerability and risk using a range of methods including
27 literature review, expert interviews and participatory methods.

28
29 In terms of cost assessment, most studies use the direct cost method, in which exogenous prices are multiplied with
30 quantities and aggregated. For instance, the amount of land lost to sea level rise is multiplied by the value of land.
31 Estimates of land values are based on data from the respective markets. For wetland loss, direct costs are used too
32 but the “prices” are taken from monetary valuation studies for non-market goods and services (Woodward and Wui,
33 2001; Brander *et al.*, 2006; Barbier, 2012). Some studies use computable general equilibrium models and growth
34 models to study the indirect and dynamic impacts of climate change, including sea level rise. These studies are
35 reviewed in Chapter 10.

36 37 38 **5.5.2. Coastal Systems**

39
40 [Section 5.3 and 5.4 overlaps to be resolved in this section]

41
42 Since AR4, no new global study of the vulnerability of coastal wetlands to sea-level rise has been conducted.
43 Dasgupta *et al.* (2009) estimated that 88,224 – 347,400 km² of wetland area are exposed to sea-level rises of 1-5 m
44 across 84 developing nations, but it is unknown how much of this area will be lost, as wetlands are able to keep pace
45 with rising sea-levels if sufficient sediment supply and migration space is available (McFadden *et al.*, 2007). The
46 DIVA model considered this process (Hinkel and Klein, 2009) but has not been applied globally for this impact.
47 McLeod *et al.* (2010a) applied DIVA to the Coral Triangle Nations and find that under 0.54m of SLR by 2100, the
48 Solomon Islands may lose 68%, the Philippines 51% and East Timor 50% of their total coastal wetland area. This
49 study also found that the composition of wetland area changed toward a greater percentage of mangroves and
50 smaller percentage of unvegetated wetlands. The majority of work on the vulnerability of coastal habitats and
51 ecosystems is carried out at local scales. The remainder of this chapter summarizes results of these studies for
52 various types of coastal ecosystems.

5.5.2.1. Rocky Shores

The FAR identified differences in the vulnerability of rocky shores to erosion due to various climate factors, primarily in terms of cliff retreat patterns and rates, and concluded that cliffs formed in softer lithologies are likely to retreat more rapidly in the future. Since then research has confirmed the large spatial and temporal variability of these processes in different parts of the world and has examined the control of various factors on the rates of retreat. Findings show that the main risks for this type of coastal systems appear to arise from sea-level rise.

Jackson and MacIvvenny (2011) estimated potential changes for the coasts of northern Scotland due to sea-level rise. They found rising sea levels to lead to a significant decrease and steepening of the intertidal area in rocky shores, which could have further impacts on the ecology of intertidal areas. In a modelling study, Trenhaile (2011) found sea-level rise to trigger faster rates of cliff recession, especially in coasts that are already retreating fast. However, results further suggested that coasts currently retreating slowly would experience the largest proportional increases in retreat rates. Increases in storminess appear to have smaller effects on rocky shores (Trenhaile, 2011; Dawson *et al.*, 2009). Importantly, the study of Dawson *et al.* (2009) in the East Anglian coast of England demonstrated the potential of cliff erosion in reducing flood risk, thus highlighting the tradeoffs that exist between cliff erosion and flood impacts.

Other factors may also affect the response of rocky shores to changes in climate. However, the study of Dornbusch *et al.* (2008) who examined cliff retreat rates in East Sussex, England for a period of 130 years concluded that climate factors, such as precipitation and frost, appear to have a limited role in cliff retreat. Currently, our understanding of the response of rocky coastlines to changes in climate is limited, partly due to its large spatial and temporal variability. Research also suggests that the response of these systems to larger changes, particularly in the rates of sea-level rise, may lead to increased sensitivity (Dawson *et al.*, 2009; Jackson and McIlveny, 2011), which needs to be further explored.

5.5.2.2. Beaches and Sand Dunes

Beaches and sand dunes are particularly vulnerable to increased erosion under sea-level rise. Vulnerability of beaches to sea-level rise is assessed qualitatively (Abuodha and Woodroffe, 2010; Sheik Mujabar and Chandrasekar, 2011) or quantitatively (Snoussi *et al.*, 2009). However, the widely employed Bruun rule has been under criticism (Cooper and Pilkey, 2004; Ranasinghe and Stive, 2009), particularly when used for national to local assessments, and alternative models are proposed (Ranasinghe *et al.*, 2012). Also, advanced numerical models, such as the Long-Term Configuration (LTC) model (Coelho *et al.*, 2007) have been applied for evaluating the effects of wave climate and sea-level rise on beach erosion.

Further to erosion, beach and dune recession and inundation will lead to loss of coastal biodiversity. Defeo *et al.* (2009) discussed the impacts of climate change on sandy beach ecosystems, with particular reference to erosion and rising temperatures and how those factors can affect habitats and food supply. Some studies that assess impacts of sea-level rise on beach ecosystems exist. For example, Fuentes *et al.* (2010) discussed the impacts of sea-level rise on green turtle population in the northern Great Barrier Reef, Australia. Nevertheless, the links between climate change and coastal biodiversity in beach ecosystems have not so far been adequately studied.

5.5.2.3. Estuaries

Estuaries are vulnerable to a range of climate drivers including sea-level rise, changes in runoff and storminess. The vulnerability of estuaries varies geographically and is determined by both direct changes in environmental variables as well as changes in the geomorphology of estuaries (e.g. build-up of sand bars across estuary entrances) resulting from factors sensitive to climate (e.g. runoff, erosion, salinization) (Gillanders *et al.*, 2011).

Some studies found that estuaries are particularly vulnerable to fluctuations in salinity which may result from changes in precipitation extremes and freshwater supply. Pollack *et al.* (2011) assessed long-term trends in the

1 response of benthic macrofauna to climate variability in Lavada-Colorado estuary, Texas and found the abundance
2 of benthic macrofauna to be significantly correlated with salinity but not with temperature. Levinton *et al.* (2011)
3 simulated salinities for a period of 100 years for the Hudson estuary and found it vulnerable to changes in
4 precipitation and discharge with respect to oyster mortality. Fujii and Raffaelli (2008) examined the spatial patterns
5 of benthic macrofaunal biomass in the Humber estuary, UK, and related them to variables such as salinity, sediment
6 characteristics and morphological factors. Using model simulations they estimated that sea-level rise and associated
7 changes in these variables could have significant implications for the intertidal habitats of the estuary.

8
9 Changes in estuaries will also affect fisheries. Changes in salinity have led to reduced species abundance (Zampatti
10 *et al.*, 2010) but the response of the ecosystem is complex as species have been found to react to these changes by
11 moving in other areas of the estuary (Sakabe and Lyle, 2010).

14 5.5.2.4. *Temperate Lagoons*

15
16 Climate-induced changes in coastal lagoons can include, among others, the redistribution of sediments and altered
17 sedimentation patterns as well as changes in turbidity, salinity and temperature (Anthony *et al.*, 2009). The response
18 of lagoon systems to these changes will vary greatly, both spatially and temporally, between lagoons and will
19 depend on the physical characteristics of the lagoons and on the processes that control these characteristics.

20
21 Vulnerability of coastal lagoons to different types of stresses, such as sea-level rise, storminess or increased water
22 temperature, has been studied for several lagoons around the world (Bruneau *et al.*, 2011; Brito *et al.*, 2010; Nixon
23 *et al.*, 2009) and some characteristics of individual responses of specific lagoon systems can be extrapolated for
24 other lagoon systems (Canu *et al.*, 2010). Further work is required to understand the response of lagoons to different
25 drivers. At the same time, uncertainties exist as to how human and climate induced pressures will affect vulnerable
26 lagoon systems and how the implications of potential management schemes for maintaining lagoon ecosystem
27 services (Chapman, 2010) will impact coastal lagoons.

30 5.5.2.5. *Salt Marshes*

31
32 The future development of coastal salt marshes is subject to a variety of natural and anthropogenic risks. As the
33 vertical growth of salt marshes tends towards an equilibrium with sea-level rise (Redfield, 1972) salt-marsh
34 accretion rates lag behind accelerated sea-level rise rate by about 20-30 years (Kirwan and Temmerman, 2009).
35 Accelerating sea-level rise rates may therefore result in marsh submergence, retrogressive vegetation succession, or
36 complete drowning of the marsh (Craft *et al.*, 2009). Salt marshes appear to be more resilient to sea-level rise in
37 macro-tidal environments than in micro-tidal environments (Kirwan *et al.*, 2010).

38
39 Subsidence of salt marshes is also triggered by human activities, such as the extraction of underground natural
40 resources (Day Jr. *et al.*, 1999; Dijkema, 1997), or the use of salt marshes as pasturelands for livestock, inducing
41 sediment compaction and shallow subsidence of the salt marsh surface. Natural shallow subsidence within the salt
42 marsh layer is observed as a consequence of autocompaction of the marsh sediments and is especially large on
43 highly organic marshes (Allen, 2000; Cahoon *et al.*, 1995).

44
45 Besides relative sea-level rise, potential changes in storminess can play a role in the survival of marshes. The natural
46 development of salt marshes, continuously migrating inland as sea level rises and eroding at the seaward edge, is
47 inhibited at many coastlines since salt marshes have been embanked in historic times or coastlines were fixed in
48 place for coastal protection reasons (Doody, 2004). This effect, often referred to as coastal squeeze, is amplified
49 through lateral erosion of the salt marshes, if storminess and associated wave heights are increasing (van der Wal
50 and Pye, 2004). However, increasing storminess can also increase the resilience of salt marshes towards drowning as
51 vertical accretion is enhanced through increasing inundation heights and inundation frequencies (Schuerch *et al.*,
52 2012).

5.5.2.6. *Mangroves*

Sea-level rise poses the greatest challenge to mangroves (Gilman *et al.*, 2008). Various studies indicate an increased vulnerability of mangrove forests to rises in sea-level. Lovelock *et al.* (2011) found subsidence due to organic sediment compaction in western Moreton Bay to negate surface accretion processes, rendering the mangrove forest vulnerable to sea-level rise. At the same time, mangroves in the eastern part of the bay appeared to be stable with current rates of sea level rise, mostly due to sufficient sediment supply. In Micronesia, Krauss *et al.* (2010) also reported differential susceptibilities of mangrove forests to sea-level rise and found fringe mangrove forests to be most vulnerable. However, due to the importance of subsurface root accumulation for mangrove elevation (McKee *et al.*, 2007; McKee, 2011) and therefore in mangroves keeping up with rising sea level, further work is needed for understanding the response of mangrove forests to changes in different climate factors. Langley *et al.* (2009) suggested that increased CO₂ can stimulate elevation gain for wetlands and can counterbalance sea-level rise. Furthermore, changes in precipitation patterns may also affect the spatial distribution and growth of mangroves by influencing salinity and fluvial supply of sediment and nutrients. However the response of mangroves to variability in precipitation remains largely unexplored (Gilman *et al.*, 2008).

5.5.2.7. *Seagrass Meadows*

Currently, a worldwide decline of seagrass beds is being observed (Lotze *et al.*, 2006; Waycott *et al.*, 2009; Short *et al.*, 2011). In most cases the decline is caused by direct human impact such as eutrophication, increased sedimentation as well as habitat and ecological degradation, but other drivers are expected to be also relevant (Orth *et al.*, 2006).

Increased sea surface temperatures are suggested as an important seagrass driver and shifts in seagrass distribution, alteration of growth rates, physiological functions and sexual reproduction are expected, even though the extent of impacts depends on the thermal tolerance of individual species (Orth *et al.*, 2006; Short *et al.*, 2011). Elevated temperatures may also increase growth of algae and epiphytes which are in competition to seagrass, especially for sunlight (Short and Neckles, 1999; Holmer *et al.*, 2011). This situation will have further implications as seagrasses (*Zostera marina*) show increased light requirements with higher temperatures as their photosynthetic capacities decrease (Zimmerman *et al.*, 1989; Moore *et al.*, 1997).

Rising sea levels are also expected to have multiple effects on seagrasses. Increases in water depths above present meadows will lead to reduced light conditions, while increased hydrodynamics (wave and currents), often associated with a higher sea level, can aggravate this effect by increasing the amount of suspended matter in the water. As studies in the U.S. and the Wadden Sea have shown, seagrass is particularly sensitive to increased hydrodynamics (Fonseca and Bell, 1998; Schanz and Asmus, 2003) and to their indirect effects (Cabaço and Santos, 2007; Dolch and Reise, 2010). Such conditions can be exacerbated by changes in storminess, leading to a significant threat for seagrass communities (Dolch and Reise, 2010). Further risks for seagrass from storms can be caused by heavy rainfall. Assessing the vulnerability of seagrasses to hurricanes, Carlson Jr. *et al.* (2010) showed that increased runoff following the 1997-98 El Nino event resulted in large declines in seagrass in Florida. These impacts were possibly due to light stress caused by increased turbidity, dissolved organic matter and phytoplankton blooms and were more damaging than the physical impacts of moderate tropical cyclones (Carlson Jr. *et al.*, 2010).

5.5.3. *Human Activities*

Since the publication of AR4 progress has been made in quantifying risks and vulnerabilities of coastal human systems to climate change and sea-level rise. A large number of regional, national and sub-national scale studies has been conducted using the range of methodologies presented in 5.5.1. Table 5-5 highlights some of these studies. A comprehensive account of vulnerabilities of specific regions is given in the regional chapters.

At the global scale, a number of studies have been conducted since AR4. Generally, these studies confirm, as also highlighted in AR4, that coastal vulnerability and risks are strongly influenced by non climatic drivers, in particular

1 socio-economic development, which influences vulnerability in two ways: It determines the level of future exposure
2 and it determines the level of future capacity to adapt (Nicholls *et al.*, 2007).

3
4 For densely populated coastal areas, socio-economic development is expected to be the major driver of exposure and
5 hence impacts during the 21st century (Nicholls and Cazenave, 2010). Under the UN medium population projections,
6 the population of 126 major port cities exposed to a 1-in-100 year flood event is expected to increase by 50% in
7 2070 through sea-level rise alone and by 150% if socio-economic development is considered on top of this (Hanson
8 *et al.*, 2011). Comparatively, Asia contains the greatest exposure in terms of population, assets and productivity and
9 the largest increases in exposure through socio-economic development are expected in Asia and Africa (Dasgupta,
10 2009; Hanson, 2011; Kebede and Nicholls, 2012).

11
12 Two recent global studies using the DIVA model indicate that the potential impacts on human activities in the 21st
13 century are substantial, but also that impacts can be reduced substantially through adaptation. Nicholls *et al.* (2011)
14 estimated that without adaptation 72-187 million people would be displaced due to submergence and erosion by
15 2100 assuming sea level increases of 0.5-2.0 m by 2100. About 70% of these people are from East, Southeast and
16 South Asia. Adaptation reduces these impacts roughly by three orders of magnitude, with annual incremental
17 adaptation costs between US\$25 and \$270 billion. Hinkel *et al.* (2012) estimated the expected number of people
18 flooded annually in 2100 to reach 170-260 million per year in 2100 without adaptation and two orders of magnitude
19 smaller with adaptation assuming global mean sea-level rises of 0.6-1.3m by 2100.

20
21 Few studies have assessed the effects of mitigation on coastal risks. Nicholls and Lowe (2004) estimated that
22 stabilizing emission at 550 ppm reduces the number of people flooded annually by factor 50% in 2110 compared to
23 a scenario of unmitigated emissions. Tol (2007) found that stabilizing emissions at 550 ppm reduces global impacts
24 on wetlands and drylands by about 10% in 2100 compared to scenario of unmitigated emissions. Hinkel *et al.* (2012)
25 reported that stabilizing emissions at 450 ppm-CO₂-eq reduces flood risks by 23-29%. These numbers need to be
26 taken with caution, as all three studies assume that the effects of lower levels of global warming due to mitigation on
27 global mean sea-level rise are assumed to be small (about 10-20cm). This may not be the case as unmitigated
28 global warming increases the risk of an accelerated melting of the ice sheets of Greenland and West Antarctica
29 leading to higher rates of sea-level rise, which were not considered in the studies listed above.

30
31 Recent studies also underpin the AR4 conclusions that there are significant regional differences in coastal
32 vulnerability (Nicholls *et al.*, 2007). Most countries in South, South East and East Asia are particularly vulnerable to
33 sea-level rise due to rapid economic growth and coast-ward migration of people into urban coastal areas together
34 with high rates of anthropogenic subsidence in deltas where many of the densely populated areas are located
35 (Nicholls and Cazenave, 2010). At the same time, economic growth also increases the capacity to adapt and the
36 benefits of adaptation are also highest in these regions (Nicholls *et al.*, 2010). On the contrary, while many African
37 countries experience a similar trend in rapid urban coastal growth, the level of economic development is generally
38 lower and hence the capacity to adapt is smaller (Hinkel *et al.*, 2011; Kebede and Nicholls, 2012).

39
40 [INSERT TABLE 5-4 HERE

41 Table 5-4: Summary of Coastal Vulnerability Assessments conducted at national or regional scale. Studies that
42 either costed (Adapt\$) or considered adaptation options (Adapt) are indicated. In studies that have considered
43 adaptation: WOA=without adaptation; WA=with adaptation. CVI=Coastal Vulnerability Index; GIS=Geographic
44 Information System, DIVA=Dynamic Interactive Vulnerability Assessment model. RSLR=relative sea level rise.]

45 46 47 5.5.4. Costs

48
49 A comprehensive picture on coastal vulnerability needs to take into account costs, in particular the costs of inaction
50 in relation to the costs of adaptation and residual damages. Since the AR4, a large number of studies have assessed
51 costs of sea-level rise to regions and countries. These will be discussed comprehensively in the Regional Chapters.
52 There are only a few studies that systematically compare costs across the countries of the world and provide a
53 comprehensive and internally consistent estimate of the costs at global scales (Table 5-5). These studies are difficult

1 to compare due to differences in scenarios used, impacts considered, methodologies applied and impact indicators
2 used. Nevertheless, some robust findings emerge.

3
4 [INSERT TABLE 5-5 HERE

5 Table 5-5: Global assessments of costs of sea-level rise.]
6

7 Recent global modeling studies indicate that while the cost of sea-level rise in the 21st are substantial, the cost of
8 inaction are larger than the sum of adaptation and residual damages costs for large parts of the world. Nicholls and
9 Tol (2006) applied FUND to estimate the direct costs of dry land loss due to submergence and wetland loss under a
10 global sea-level rise of 0.35m by 2080 for all countries in the world for 4 of the SRES scenarios. Direct costs are the
11 costs of coastal protection, plus residual land loss, plus wetland loss due to both sea level rise and coastal protection.
12 They found that coastal protection is cheap relative to the value of the land protected and, using the cost-benefit
13 model of Fankhauser (1995), concluded that it would be economically justified to protect the bulk of low-lying,
14 populated coastline. They assumed that the value of land increases with economic growth but the costs of coastal
15 protection do not. Thus, more than 80% of the exposed coast is protected in all but 15 countries in all scenarios, and
16 coastal protection is stronger in the scenarios that assume more rapid economic growth. The annual cost of coastal
17 protection is below 0.1% of GDP in over 180 countries. Land loss nowhere exceeds more than 3% of the country's
18 total area. As the most valuable areas are protected first, the economic impact would be smaller than that (but is
19 unreported by Nicholls and Tol, 2006). Using the same methodology, Anthoff *et al.* (2010) confirmed that applying
20 protection is economically rational for most countries even for a global mean sea-level rise of 2m by 2100. Studies
21 using DIVA come to similar conclusions. Hinkel *et al.* (2012) estimated that the direct cost of coastal flooding
22 (comprising dike upgrade, dike maintenance and residual damage cost) reaches US\$ 300 billion per year in 2100
23 without adaptation and US\$90 billion per year with adaptation under a 1.26 m sea-level rise scenario.
24

25 Even though the risks of sea-level rise seem manageable for human activities in large parts of the developed world
26 through enhancing coastal protection, the risk of low probability high impact events will increase with protection
27 and rising sea-levels. This holds particularly true for many coastal cities such as London, Tokyo, Shanghai,
28 Hamburg and Rotterdam that already rely heavily on coastal defences (Nicholls *et al.*, 2007) and defence failure
29 could have severe consequences.
30

31 From the perspective of less-wealthier and small island countries, annual costs of sea-level rise can amount to
32 several percentage points of national GDP. Nicholls and Tol (2006) reported the highest coastal protection costs for
33 Palau, Tuvalu and the Federated States of Micronesia. For the later country, protection costs reach 14% of GDP in
34 the 2080s in the B2 scenario. Hinkel *et al.* (2012) reported that the direct annual cost of coastal flooding (including
35 adaptation and residual damages) in 2100 lies between 5% and 9% of GDP for Kiribati, the Solomon Islands,
36 Vanuatu and Tuvalu under 0.64m of sea-level rise.
37

38 These regional differences in vulnerability are also highlighted in continental scale studies. For the European Union,
39 Hinkel *et al.* (2010) estimated that without adaptation the total monetary damage caused by flooding, salinity
40 intrusion, erosion and migration is estimated to US\$17 billion per year under global mean sea-level rise scenarios of
41 0.35m-0.45m. Adaptation reduces these damages by one order of magnitude and costs 2.6-3.5 billion US\$ per year
42 in 2100. For Africa, Hinkel *et al.* (2011) estimated the expected annual damage costs due to flooding, erosion and
43 salinity intrusion to US\$5-9 billion under scenarios of global mean sea-level rises from 0.64-1.26 m by 2100, if no
44 adaptation takes place. Adaptation cuts damage costs in half by 2100, but requires substantial investments of US\$2-
45 US\$6 billion per year. The adaption costs reported are incremental costs that do not take into account the current
46 adaptation deficit, which is expected to be substantial for developing countries but has not been assessed in detail
47 (Parry *et al.* 2009). Hinkel *et al.* (2011) estimated that overcoming Africa's current adaptation deficit with respect to
48 coastal flooding would require an initial investment of US\$300 billion for building dikes and an additional US\$3
49 billion per year for future maintenance.
50

5.5.5. *Uncertainties and the Long-Term Commitment to Sea-Level Rise*

The three major sources of uncertainty in the assessment of coastal vulnerability, risk and costs are (i) sea-level rise including local subsidence, (ii) socioeconomic development including coastward migration and urbanization patterns, and (iii) the level of adaptation that will take place. The available studies have only explored a small fraction of this uncertainty. Only few assessments consider global mean sea-level rise scenarios beyond the range given in the AR4, thus excluding the impacts of a possible large contribution of the melting of the ice sheets of Greenland and Antarctica to global mean sea-level rise. Studies considering regional patterns of climate-induced sea-level rise are missing. Many studies rely on few or only a single socio-economic scenario.

Only few studies consider adaptation in the estimation of vulnerabilities and risks, which gives an incomplete picture as the question to what extent society can handle the potential impacts that is omitted (Hinkel, 2012). In the case where adaptation is considered, most studies focus on protection via hard structures, because protection via 'soft' options such as dune or mangrove rehabilitation is difficult to simulate at broad scales and cost estimates are less developed (e.g., Linham and Nicholls, 2010; Hinkel *et al.*, 2011). Many more adaptation options are, however, available including retreat and accommodation (See Section 5.6) and future work needs to consider these. With future socio-economic development being often the major risk factor, steering development away from low-lying areas seems to be a potentially effective adaptation strategy (Kebede and Nicholls, 2012) which, however, has not been explored in studies.

For deltas, another major source of uncertainty is human-induced subsidence due to anthropogenic sediment compaction as a result of the withdrawal of ground fluids such as oil, water and gas. Human-induced subsidence may lead to rates of local sea-level rise that are an order of magnitude higher than current rates of climate-induced global-mean sea level rise (Syvitski *et al.*, 2009). Densely populated deltas are particularly vulnerable and many of the world's mega-cities are situated in deltas have subsided by several meters during the 20th century (Nicholls, 1995). It is difficult to predict how this trend will continue as information on annual rates of human-induced subsidence is extremely limited (Hanson *et al.*, 2011). Furthermore, while some cities such as Shanghai and Tokyo have managed to control subsidence rates through policy measures, other cities such as Bangkok, Manila and Jakarta continue to subside at high rates (Nicholls 1995; Hanson *et al.*, 2011).

The available studies also only cover a limited range of impacts. There is no recent global study on the long-term impacts on coastal wetlands and no global consistent study taking into account all major impacts including land loss due erosion, land loss due to submergence, wetland loss, flood damage and salinity intrusion. In particular, studies either focus on land loss due to submergence or on flood damage. The majority of research on coastal flood risks only assesses exposure and not damages (see Table 5-4). There is also a lack of intermediate scale methodologies for assessing coastal flood risk. Assessments are either carried out at local scales using hydrodynamic models or at global scales using damage functions. As there are so few studies of the costs of sea level rise at a global level, confidence is low. Uncertainties are largely unknown. The need for further research is large. Finally, there is insufficient knowledge on indirect impacts and costs of coastal flooding including health impacts and those related to the disruption of economic activities through a flood.

Most studies assess risks of single hazards. Integrated studies assessing multiple hazards and trade-offs between adaptation options are rare. Flood defences, for example, prevent the inland migration of coastal habitat when sea level is rising leading to a loss of intertidal habitat through coastal squeeze (Nicholls *et al.*, 2007; McFadden *et al.*, 2007). Furthermore, coastal armoring in one location may have negative consequences on other locations as reduced long-shore sediment transport through protection measures may increase both flooding and erosion risk at other locations (e.g., Dawson *et al.*, 2009). These trade-offs and processes need to be further explored.

Finally, few studies take into account that vulnerability and risk will increase beyond the 21st century as sea level will continue to rise for centuries due to global warming until reaching equilibrium conditions even if climate forcing is stabilized, which was termed commitment to sea-level rise in the AR4 (Meehl *et al.*, 2007; Nicholls *et al.*, 2007). Sea-level rise due to thermal expansion is estimated to reach equilibrium in the year 3000 at levels of about 0.5m per °C of warming if deep water-formation is sustained and higher values if not (Meehl *et al.*, 2007). The

1 contribution of the melting of the ice-sheets to long-term sea-level rise is potentially much larger and while the
2 timescales involved are unknown and subject to intense research, there is concern about a potential rapid melting
3 due to self-amplification mechanisms involved (Levermann *et al.*, 2011).

4 5 6 **5.6. Adaptation and Managing Risks**

7 8 **5.6.1. Approaches**

9
10 Adaptation to coastal risks from climate variability and change occurs in the context of existing governance and
11 socio-ecological systems, regardless of whether adaptation is proactive and planned or reactive and ad hoc. To
12 discuss adaptation outside of these contextual factors is theoretical at best and misleading at worst. Governance
13 involves the legal and institutional context of coastal management; ownership rules related to coastal land and
14 resources; stakeholders typically involved in coastal management decisions, and the social norms, rules, and
15 dynamics that guide their interactions.

16
17 The socio-ecological system within which governments and individuals act is intricately connected to the
18 governance system, but it helps to single it out as its own co-determinant of adaptation. The socio-economic context
19 of adaptation includes the general state of the (local) economy; prevalent urban and economic sectors dependent on
20 or located in the coastal zone; past, present and planned development decisions; the degree of demographic
21 concentration and the resulting build-up; the technologies employed; the state of human welfare; as well as any past
22 or existing social conflict and social capital; access to power and relationships among power holders and affected
23 stakeholders. Cultural factors play important roles, e.g., on world views, gender, class or caste relationships,
24 concurrent pressures and trends of cultural transformation.

25
26 The physical and ecological contexts of relevance for adaptation is the geologic/geomorphologic type of coastline,
27 the prevalent climate, the local ecosystems and existing biodiversity; the local rate of relative sea-level rise and
28 interacting climate change impacts (temperature, precipitation, storm regime, sediment supply, and salinity
29 changes); and concurrent non-climatic environmental or human pressures and trends on coastal geo-ecological
30 systems. Together, the particular determinants of physical-ecological processes present a range of what adaptation
31 options are physically feasible or environmentally appropriate.

32
33 _____ START BOX 5-4 HERE _____

34 35 **Box 5.4. Case Study – Paradigm Shift in Adaptation to Rising Sea Levels in The Netherlands**

36
37 Inhabitants in the low-lying coastal area of the Netherlands have been facing coastal and river flooding for centuries.
38 Extreme sea levels together with long-term delta subsidence have been responsible for major catastrophic events
39 causing thousands of fatalities over the last ten centuries. The expansion of habitable land and the increase in the
40 population exposed to catastrophic events have co-existed with technological improvements for coastal defence and
41 large investments resulting in a dramatic decrease in mortality rate (Van Baars and Van Kempen, 2009). Most of the
42 coastal protection has been implemented during the second half of the twentieth century, resulting in the
43 construction of large and numerous infrastructures altering the appearance of an important part of the Dutch coastal
44 area (Kabat *et al.*, 2009).

45
46 [INSERT FIGURE 5-7 HERE

47 Figure 5-7: Case study: paradigm shift in adaptation to rising sea levels in the Netherlands. Source: Stive *et al.*,
48 2011.]

49
50 At present, nine million residents live in coastal areas at an elevation below sea level, where roughly 65% of gross
51 national product is generated (Stive *et al.*, 2011). The anticipation of a changing climate, including sea-level rise,
52 during the twenty-first century, together with an audit carried out in 2007 giving evidence of an aging flood
53 protection system and an increasing vulnerability has renewed the demand for new plans for water and coastal
54 management.

1
2 In this context the Dutch Government has set out far-reaching recommendations on how to keep the country flood-
3 proof over the 21st century by considering a paradigm shift, namely addressing coastal protection ‘working with
4 nature’ instead of only ‘fighting’ the forces of nature with engineered structures and providing ‘room for river’.
5 Some of the recommendations include soft and environmentally friendly solutions such as preserving land from
6 development to accommodate increased river inundation; raising the level of lakes, to ensure a continuous supply of
7 fresh water; removing existing flooding protecting structures to restore natural estuary and tidal regimes; improving
8 the standards of flood protection by a factor of 10 until 2050; maintaining flood protection by beach nourishment,
9 expanding the coast seaward in the next century and putting in place the necessary political-administrative, legal and
10 financial resources. The estimated total cost of implementing this ambitious plan is €2.5-3.1 billion a year to 2050,
11 representing a 0.5% of the current Dutch annual gross national product (Stive *et al.*, 2011).
12

13 Aerts *et al.* (2008) estimated today’s economic damage from flooding as approximately €190 billion covering both
14 direct and indirect damage. The estimated future potential damage would increase to €400 to €800 billion in 2040
15 and €3700 billion in 2100 in the absence of any measures, given a sea-level rise of 24 to 60 cm in 2040 and 150 cm
16 in 2100. The factors that govern calculations of estimated future potential damage are economic growth combined
17 with indirect damage. The Delta Committee suggested that, under the umbrella of this paradigm shift in the
18 approach to water and coastal management, climate change offers key challenges that can result into societal and
19 economic growth and evolution, moving the Netherlands into a sustainable country (Kabat *et al.*, 2009).
20

21 _____END BOX 5-4 HERE _____
22

23 Coastal management typically needs to balance multiple goals that can and often do conflict, and frequently are
24 adjudicated among in an unbalanced fashion. Among the most relevant coastal management goals for adaptation are
25 the minimization of risks and impacts from coastal hazards to ensure public safety and welfare; economic
26 development and use of coastal resources; and protection of coastal environmental resources, natural assets, and
27 ecosystems.
28

29 Optimizing solutions taking into account the three goals is a common problem in coastal zones. Many approaches
30 have been developed over time to achieve greater integration, better social, ecological, and economic outcomes
31 when trade-offs are inevitable, and smoother governance, including Integrated Coastal Management (e.g., Sales,
32 2009; Christie *et al.*, 2005), Community-Based Adaptation (e.g., Dumaru, 2010; Huq and Reid, 2007; Reid *et al.*,
33 2009), Ecosystem-Based Adaptation (e.g., Zeitlin *et al.*, 2012; Vignola *et al.*, 2009; IUCN, 2008), and Disaster Risk
34 Reduction and Management (Shaw *et al.*, 2010; IPCC, 2012).
35

36 _____ START BOX 5-5 HERE _____
37

38 **Box 5-5. Case Study – Climate Change Adaptation and Integrated Coastal Zone Management in Developing** 39 **Countries** 40

41 Integrated Coastal Zone Management (ICZM) promotes sectoral and spatial integration of various activities in the
42 coastal zone by establishing coordination across varying sectors and government institutions with a view to
43 sustainably develop coastal resources and protect the environment. This makes it feasible for combining climate
44 change adaptation with ICZM as a part of integrative effort in coastal management. However, the difficulties of
45 integration and coordination are present both in developed and developing economies and could arise if and when
46 climate change adaptation is mainstreamed into ICZM.
47

48 ICZM in developing countries fostered by international organizations under the UN or (non)governmental units, in
49 particular, struggle to meet the goals of ICZM (Isager, 2008). The drawbacks that are present in the implementation
50 of ICZM in developing countries consequently act as constraints to enforcing climate change adaptation within
51 ICZM. For example, inadequate financial commitment follows when initial funding by the external organizations
52 disappears, and national governments have to step up to finance the cost of ICZM (Ibrahim and Shaw, 2012).
53 Ineffective coastal governance is more visible in cases where the capacity of actors is low, the operation of single
54 agency is dysfunctional, and subsequently, the integration of multiple coastal agencies is beyond the reach of many

1 developing countries (Ibrahim and Shaw, 2012; Martinez *et al.*, 2011). Politics are also a strong force because of the
2 involvement of various stakeholders, the hierarchy of government agencies and ministries, and the power of the
3 majority political party or political leaders (Tabet and Fanning, 2012; Isager, 2008). Furthermore, the nature of
4 public participation in developing countries differs from that in developed countries, and different norms and
5 cultures need to be taken into account to assess public participation central to ICZM (Barale and Özhan, 2010).

6
7 As such, legal and institutional capabilities critical to the implementation of ICZM are often not available in
8 developing countries. Instead, governments generate climate change adaptation strategies that are part of shoreline
9 management plan, regional development, disaster management, and coastal resource management. In addition, cases
10 of adaptation strategies specific to climate change in practice are few. Most are at the planning stage equipped with
11 scenarios. Existing strategies derive from responses to coastal disasters and economic and social change affecting
12 coastal livelihoods. For example, the Bangladesh case illustrates the benefits of trained volunteers readily accessible
13 to affected population to disseminate cyclone warnings, evacuate people, and conduct rescue missions (WRI, 2007).
14 This could be translated into a strategy for climate change adaptation. Other strategies in progress or at the trial stage
15 include forested buffer zones (Mustelin *et al.*, 2011) and incremental migration. The long-term results and their
16 replicability need to be further assessed.

17
18 To anticipate climate change and adapt, scholars have considered a variety of strategies that could be of use in the
19 future. No or low-regret options provide co-benefits to the goals of sustainable development, livelihood
20 improvement, and human well-being (IPCC, 2012). Combining different strategies are also increasingly under
21 consideration (Cheong, 2011; Cheong *et al.*, 2012; Cartwright, 2009). They include the blend of ecology and
22 engineering such as mangrove planting, buffer zones and land use, and insurance and structural coastal defence
23 (Yohe *et al.*, 2011). Adaptation planning by geographic scale is at work. Regional adaptation strategies (Martinez *et al.*,
24 2011) and community-based adaptation (Cutter *et al.*, 2012) are both on the rise as scale economies provided by
25 adaptation at the regional level as well as the critical role of local physical and cultural attributes and local priorities
26 are valuable in adaptation planning. Although ICZM can be a valuable policy framework to integrate climate change
27 adaptation and coastal management, no studies of the effectiveness of ICZM combined with climate change
28 adaptation in developing countries exist yet to assess its utility.

29
30 _____ END BOX 5-5 HERE _____

31
32 Adaptation – as it becomes integral to what coastal managers do –will face the same multi-purpose challenges, as
33 different interests, needs, and stakeholder viewpoints have to be addressed and as climate-driven and non-climatic
34 pressures on coastal environments grow (Tobey *et al.*, 2010). Indeed, experience to date shows that the challenges
35 with (integrative) adaptive coastal management is not radically different from those encountered with historical
36 coastal management (Tobey *et al.*, 2010). However, climate change-conscious coastal management would adjust
37 these approaches to the dynamic nature of coastal areas, long-term trends (as opposed to assuming static baselines)
38 and thus greater uncertainty and longer time frames in planning (beyond 30 years), the long-term commitments
39 inherent in climate change, the potential for physical and ecological thresholds or tipping points, and the long lead
40 times often required for making changes in coastal management (see references in Table 5-6). Garmendia *et al.*
41 (2010) suggested that improving the integration of various expertise and values can guide to the definition of
42 appropriate policy options and adequate decisions when complexity, value conflict and uncertainty exist as is,
43 generally, the case in coastal zones.

44
45 [INSERT TABLE 5-6 HERE

46 Table 5-6: Approaches to integrative, adaptive coastal management.]

47
48 To date, despite experimentation with these novel or adapted coastal management approaches, meeting the multiple
49 goals, improving governance, accounting for the most vulnerable populations and sectors and fully integrating
50 consideration of natural ecosystems is still largely aspirational. Meanwhile development in high risk areas grows,
51 coastal ecosystems continue to degrade in many regions, coastal freshwater resources are being overdrawn in many
52 highly populated areas, and vulnerability to coastal disasters grow (e.g., Jentoft, 2009; McFadden, 2008; Mercer,
53 2010; Shipman and Stojanovic, 2007).

5.6.2. Practices

Coping with the dynamics of physical processes and rapid population growth and investment in coastal areas has built a body of knowledge and tools applicable to many of the potential impacts associated with climate changes. These tools include the structural, planning and regulatory, hazard response planning, biological, and market-based tools as well as physical and integrated assessment modeling to assist in identifying possible impacts (Bedsworth and Hanak, 2010; Horstman *et al.*, 2009; Rosenzweig *et al.*, 2011). Climate change and related impacts raise new considerations, including greater degrees of uncertainty, and continue to confront long-standing analytical challenges.

Since the AR4, there has been further progress in impacts modeling and integrated assessment efforts. General adaptation tenants for climate change conservation strategies are advancing with more specific ecological and social conditions and context recommendations. The differences among coastal impact models as applied to environmental conservation goals result in important trade-offs of human and financial resources required for implementation, feedbacks and impacts represented and the degree of spatial resolution provided. The difficulty in obtaining critical information regarding appropriate uses, required data inputs and outputs, range of costs and expertise required have been identified as potential obstacles to their wider appropriate use (McLeod *et al.*, 2010a).

The scope of scale of integration is advancing. The development of successful coastal adaptation strategies needs combining scenarios of climate change and socio-economic conditions, and risk assessment (Kirshen *et al.*, 2011). For example, Dawson *et al.* (2009) employed climate, coastal management, and socioeconomic scenarios in conjunction with physical models extended over larger spatial and temporal scales to evaluate probabilistic predictions of coastal behavior with an assessment of expected annual damages and illustrated trade-offs associated with different management approaches .

Inundation models benefit from the increased availability of more accurate lidar data of coastal elevations (Gesch, 2009), although these data are not widely available. Numerical modeling linking spectral wave transformation model with calculation of gradients in potential long shore sediment transport rate can be used to project magnitudes of potential coastal erosion and accretion, under proscribed deep water wave conditions (e.g. Adams *et al.*, 2011).

Integrated assessment models continue to differ in their approaches to representing interactions among regions and sectors with the result that the ability to represent impacts and adaptation continues to involve significant limitations (UNFCCC, 2010a). For instance, these models do not consistently incorporate the interaction between impacts in one sector and human adaptation to impacts in another sector and other significant interactions (Warren, 2011). The majority of integrated assessment models address adaptation as an implicit rather than explicit process at an aggregated level with assumptions that may result in overly optimistic representations of the amount of adaptation and underrepresentation of costs (Patt *et al.*, 2010).

Efforts to develop improved vulnerability indices and to identify hotspots which serve to focus or prioritize management efforts continue to evolve although significant differences exist among them (e.g., McLaughlin and Cooper, 2010; Mustafa *et al.*, 2011; Ozyurt and Ergin, 2009). Diversity among coastal environments, local governments, institutions, economies, technologies, and cultures contribute to difficulty in generalization. Selection and availability of indicators as well as scale also contribute to differences in the sensitivity and applicability of these models across places and hazards. Consequently, tradeoffs occur between detailed locally actionable analyses and representation of broader patterns. Our ability to quantify vulnerability continues to be restricted by limits to our understanding of human adaptive capacity, broad social dynamics, and relationships between ecosystem and human well-being (Farhan and Lim, 2011; Raudsepp-Hearne *et al.*, 2010; Tol *et al.*, 2008).

Since the AR4, new information is available on the likelihood of increased rates of sea-level rise and ocean acidification. Policy recommendations for addressing ocean acidification at the local and regional levels, rather than through international mitigation efforts, are beginning to emerge. Application of existing water quality laws, land-use management to protect biological integrity, local mitigation efforts, and increased focus on data collection to inform future regulation have been proposed (Kelly *et al.*, 2011).

1
2 As adaptation planning has begun in some places, there is an emerging body of literature to inform decision-making,
3 public participation and communication efforts. Efforts to support decision-making recognize that information alone
4 may not fully serve managers needs and could be supplemented by financial and technical assistance resources as
5 well as organizations which serve as an interface between science and practice (Tribbia and Moser, 2008). Newly
6 developed mapping and visualization approaches may contribute to these processes in several ways, however there
7 is an important need for testing and evaluation of these technologies in public participation processes (Jude, 2008;
8 Sheppard *et al.*, 2011). These participation processes carry with them the challenges or power relationships met in
9 other public arenas and differences of opinion may be magnified by the uncertainty and longtime horizons
10 associated with climate change making (Few *et al.*, 2007).

11 12 13 **5.6.3. Adaptation Costs**

14
15 Efforts to assess the costs and benefits of adaptation options are continuing to evolve, although significant further
16 work is needed (Nicholls *et al.*, 2010, Yohe *et al.*, 2011). The cost of adaptation is only one part of the overall cost
17 induced by climate change, which includes also the costs of mitigation and the costs of the residual impacts.
18 Adaptation costs are defined as "the costs of planning, preparing for, facilitating, and implementing adaptation
19 measures including transition costs" and adaptation benefits are "the avoided damage costs or the accrued benefits
20 following the adoption and implementation of adaptation measures" (AR4 WG2). There are several potential
21 assessment techniques; prominent among them is the cost-benefit analysis, although it has limitations with respect to
22 the treatment of nonmarket values. All assessment types need to consider the distribution of burdens and benefits
23 across groups, sectors, or other entities (UNFCCC, 2010b). A major review of methods and findings oriented
24 towards national planning needs has also been produced under the Nairobi Work Plan (UNFCCC, 2010a). Coastal
25 assessment also differ as some take an aggregate approach working at larger levels with generalizing assumptions
26 while others take it disaggregated approach.

27
28 The coastal zone, along with water resources in agriculture, tends to have a deeper body of research on the costs and
29 benefits of adaptation options than other sectors (Argawala and Fankhauser, 2008). Within the body of research,
30 several methodological issues have been identified. These include the determination of baseline conditions;
31 treatment of uncertainty and equity including distributional impacts, ancillary benefits and public-private efforts and
32 economic valuation (UNFCCC, 2010a).

33
34 Argawal and Fankhauser (2008) summarized key features of the large number of studies that focus on the costs of
35 sea level rise impacts and adaptation. They identified three main themes: that there is extensive information
36 available on regional and global costs of adaptation, although generally only for 1-m sea-level rise; the optimal
37 percentage of coastline that should be protected in order to minimize costs (protection plus residual damage) is often
38 quite high, however that is dependent on population density and land value; and, the annualized cost estimates for
39 optimal protection are often less than 0.1% of national GDP, with the caveat that there is significant regional
40 variation and higher costs particularly for small island states.

41
42 Newer studies have emerged using a wider range of scenarios, expanded on the impacts considered, and integrated
43 other adaptation options (Anthoff *et al.*, 2010; Ciscar *et al.*, 2011; Nicholls *et al.*, 2011; Nicholls *et al.*, 2010). For
44 example, cost-benefit analyses of 0.5, 1.0, and 2.0 m sea-level rise using the FUND model show significant benefits
45 from protection, however authors caution that these findings might overestimate the extent of protection likely to be
46 implemented (Anthoff *et al.*, 2010). The UNFCCC study estimated additional adaption costs of \$4-11 billion/year in
47 2030 (Nicholls, 2007). However, those costs may be higher in the case of high-end sea level rise scenarios, and may
48 also be underestimated because the analysis focuses mainly on the incremental adaptation costs with little attention
49 to residual damages and no consideration of the adaptation deficit (Parry *et al.*, 2009). These authors go on to
50 remark that it is quite possible that the cost of addressing the adaptation deficit for coastal protection will exceed the
51 \$11 billion/year (Nicholls, 2007); however, that deficit is not well understood and requires further definition and
52 quantitative analysis (Parry *et al.*, 2009).

1 Economic models and valuation studies are emphasizing the need to address ecosystem impacts and the value of
2 ecosystem goods and services. Projected investments in coastal protection and beach nourishment would both entail
3 environmental costs (Parry *et al.*, 2009). While there has been a rapid growth in research on ecosystem services,
4 there is a substantial research agenda, including some longstanding challenges in valuation, to be addressed in both
5 the ecological and economic dimensions (Anton *et al.*, 2010; Balmford *et al.*, 2011; Mendelsohn and Olmstead,
6 2009; Polasky and Segerson, 2009). The lack of understanding of the connections between ecosystem services and
7 human well-being (Raudsepp-Hearne *et al.*, 2010) is also a barrier to valuation.

10 5.6.4. Constraints

11
12 The principal finding in the coastal chapter of the AR4 was that “there are limits to the extent to which natural and
13 human coastal systems can adapt even to the more immediate changes in climate variability and extreme events,
14 including in more developed countries” (Nicholls *et al.*, 2007, p. 342). A variety of studies have been published in
15 the interim, reinforcing this finding, and producing a better understanding of the nature of the barriers and limits to
16 adaptation both generally (Biesbroek *et al.*, 2011; Dupuis and Knoepfel, 2011; Gifford, 2011; Sietz *et al.*, 2011;
17 Amudsen *et al.*, 2010; Burch *et al.*, 2010; Larson, 2010; Lonsdale *et al.*, 2010; Moser and Ekstrom, 2010; Adger *et al.*,
18 2009a,b; Mitchell *et al.*, 2006; Huang *et al.*, 2011); and more specifically in the coastal sector (e.g., Lata and
19 Nunn, 2011; Mozumber *et al.*, 2011; Storbjörk and Hedrén, 2011; Bedsworth and Hannak, 2010; Frazier *et al.*,
20 2010; Saroar *et al.*, 2010; Moser *et al.*, 2008; Tribbia and Moser, 2008; Ledoux *et al.*, 2005).

21
22 Since the AR4, a clearer definition of limits and barriers has emerged. Adaptation *limits* are defined as “obstacles
23 that tend to be absolute in a real sense: they constitute thresholds beyond which existing activities, land uses,
24 ecosystems, species, sustenance, or system states cannot be maintained, not even in a modified fashion” (Moser and
25 Ekstrom, 2010, p. 22026). Coastal research since the AR4 has examined particularly physical limits to natural
26 (unassisted) adaptation, e.g., of coastal marshes (Kirwan *et al.*, 2010a, b; Craft *et al.*, 2009; Langley *et al.*, 2009;
27 Mudd *et al.*, 2009). In their experimental study, Kirwan *et al.* (2010a) found that coastal marshes – due to nonlinear
28 feedbacks among inundation, tidal range, plant growth, organic matter accretion, and sediment deposition – can
29 adapt to conservative rates of sea-level rise (A1B), so long as there is sufficient sediment supply. By contrast, even
30 coastal marshes with high sediment supplies are hard-pressed to adapt to more aggressive rates of SLR (Rahmstorf,
31 2007). Marshes accustomed to large tidal ranges show greater capability to adapt than micro-tidal marshes (Kirwan
32 *et al.*, 2010b). Other studies show how different climate change impacts interact to reduce the viability of coastal
33 ecosystems sooner than when only a single driver is considered (e.g., Desantis *et al.*, 2007; Spalding and Hester,
34 2007).

35
36 By contrast, social, economic, institutional, informational and other *barriers* constitute mutable “obstacles that can
37 be overcome with concerted effort, creative management, change of thinking, prioritization, and related shifts in
38 resources, land uses, institutions, etc.” (Moser and Ekstrom, 2010, p. 22027). As Adger *et al.* (2009b) argued, most
39 social obstacles (even if they appear as limits to the involved), are barriers in that they “can be overcome with
40 sufficient political will, social support, resources, and effort” (Moser and Ekstrom, 2010, p. 22027). The common
41 thread among all barriers is that they make adaptation less efficient or less effective or may require significant
42 changes that can lead to missed opportunities, difficult trade-offs, or higher costs.

43
44 Researchers have categorized barriers in different ways, and they have placed variable emphasis on certain barriers.
45 For example, common barriers identified include negative environmental consequences, technological feasibility,
46 costs, institutional settings, entitlements and entrenched habits, political calculus, deeply held cultural values,
47 worldviews and beliefs, lack of awareness, knowledge or location-specific information, social justice concerns, or
48 negative interactions between different policy goals. Table 5-7 provides some examples of barriers found in the
49 literature specific to coastal adaptation.

50
51 [INSERT TABLE 5-7 HERE

52 Table 5-7: Common barriers to coastal adaptation.]

1 The wide range of barriers identified in Table 5-7 reflects different coastal management contexts, different foci on
2 levels of governance and actors/decision-makers, as well as different methods used in identifying them. This
3 diversity does not allow for a quantitative meta-analytical integration, and yet critical insights have emerged since
4 the AR4. First, the commonly heard claim that lack of information is the main constraint to (coastal) adaptation is
5 refuted by the wide range of barriers identified in the sampled literature listed in Table 5-7, and many of them are
6 empirically shown to be more important than lack of locally relevant, credible information. While information is
7 clearly important, it matters differently for certain actors, at certain times in the adaptation process. Second, different
8 constraints typically do not act as barriers in isolation, but come in interacting bundles. For example, Moser and
9 Tribbia (2006/2007) and Mozumber *et al.* (2011) showed that lack of staff time is related to and often correlated
10 with overall lack of resources for planning and implementation; lack of awareness is often related to both lack of
11 experience and lack of communication and education (Saroar and Routray, 2010); social resistance to certain
12 adaptation options is related to attitudes, worldviews, (spiritual) beliefs, cultural norms, place attachment, and
13 economic investment and options (Barnett and Campbell, 2010; Lata and Nunn, 2011). Third, it is therefore difficult
14 to predict which barriers matter most in any specific context but instead multiple barriers need to be addressed if
15 adaptation is to move successfully through the different stages of the management process (Moser and Ekstrom,
16 2010; Storbjörk, 2010; Lonsdale *et al.*, 2010). Nonetheless, there are some non-surprising yet important
17 commonalities: studies focused on government staff show the predominance of intra- and cross-institutional as well
18 as budgetary constraints, with informational, communication, political, and public support barriers playing important
19 additional roles (e.g., Storbjörk and Hedrén, 2011; Moser and Tribbia, 2006/2007; Ledoux *et al.*, 2005). By contrast,
20 studies focused on individuals and their views on potentially unplanned, reactive adaptation show a predominance of
21 psychosocial (place attachment, social support, social norms, identity), cultural-cognitive (beliefs, worldviews,
22 values, awareness, education) and economic barriers (e.g., Adger *et al.*, 2011; Saroar and Routray, 2010). Fifth,
23 some factors can act as enablers and added capacity to adapt, while acting in barriers at others (Burch, 2010;
24 Storbjörk, 2010). For example, strong leadership in a government agency can help motivate and advance adaptation
25 internally, while hindering cross-agency ownership of the challenges and responsibilities to plan and implement
26 adaptation (Storbjörk, 2010). A complementary insight is that some capacities or factors can compensate for other
27 present barriers, thus rendering them less severe (e.g., leadership can compensate to some extent for lack of
28 information and economic resources).

29
30 Finally, as the Ledoux *et al.* (2005) study showed explicitly, and as emerges as a common concern from wide-
31 ranging literature reviews (Biesbroek *et al.*, 2011; Ekstrom *et al.*, 2011), some critical barriers arise from the
32 interactions across policy domains, existing laws and regulations, and historical legacies (long-term impacts of past
33 decisions and policies). Dawson *et al.* (2009), for example, showed that – due to the interconnectivity of
34 geomorphologic processes within a littoral cell – attempts to reduce one coastal climate risk (e.g., erosion) may well
35 increase the exposure to another coastal climate risk (e.g., flooding). Such trade-offs can reduce the ultimate
36 effectiveness of one or all of the interacting adaptation options.

37 38 39 **5.6.5. Links between Adaptation and Mitigation**

40
41 For the foreseeable future, coastal areas will be preoccupied with managing interacting stresses from sea-level rise,
42 temperature increases, precipitation changes, changing storm regimes, runoff from coastal watersheds into near-
43 coastal waters as well as non-climatic stressors such as population and development increases in vulnerable areas,
44 pollution from land use and industrial activities, and threats from infectious diseases (e.g., Melbourne-Thomas *et al.*,
45 2011; Bunce *et al.* 2010a; Halpern *et al.*, 2008a). At the same time, successful adaptive coastal management of
46 climate risks will involve assessing and minimizing potential trade-offs with other non-climate policy goals (e.g.,
47 economic development, enhancement of coastal tourism) and interactions between adaptation and mitigation (e.g.
48 Bunce *et al.*, 2010b; Barbier *et al.*, 2008; Tol 2007; Brown *et al.*, 2002).

49
50 A range of studies suggest that adaptation will be the predominant approach to reducing climate risks to coastal
51 communities, populations, resources and activities over the 21st century due to the enormous momentum involved in
52 sea-level rise and the time lag between emission reductions, temperature changes and impacts on global sea levels
53 (Nicholls *et al.*, 2011; Nicholls *et al.*, 2007). Systematic assessment of potential synergies and tradeoffs between
54 mitigation, adaptation, and other, non-climatic policy goals and efforts to maintain or increase flexibility to enable

1 policy adjustments in the future have been proposed as strategies to recognize, avoid and minimize the risk of
2 negative policy interactions (e.g., Vermaat *et al.* 2005; Nicholls *et al.*, 2011). Positive synergies and
3 complementarities between mitigation and adaptation in the coastal sector exist because many coastal zone-based
4 activities and various coastal management activities involve emissions of greenhouse gases and will be impacted to
5 varying degrees by climate change (Section 5.3). The first few items in Table 5-8 show examples of such positive
6 interactions. In addition to positive interactions, the possibility for negative interactions (or tradeoffs) exists as well.

7
8 [INSERT TABLE 5-8 HERE

9 Table 5-8: Positive synergies and tradeoffs between selected coastal adaptation and mitigation options.]

10
11 Klein *et al.* (2007, p. 749) defined trade-offs between mitigation and adaptation as the “balancing of adaptation and
12 mitigation when it is not possible to carry out both activities fully at the same time (e.g., due to financial or other
13 constraints)”. This definition has been criticized as being too broad and potentially obscuring important differences
14 between tradeoffs (Moser, 2011). A finer differentiation would distinguish various types of constraints that may
15 prevent the full implementation of selected adaptation and mitigation measures either because of insufficient
16 supporting means and conditions or due to concerns over unwanted outcomes. Such undesirable outcomes may
17 include, but not be constrained to, negative environmental consequences, undesirable social implications, political
18 repercussions, equity concerns such as distributional or intergenerational impacts, and so on (see references in Table
19 5-8). The second and third sections of Table 5-8 list a range of adaptation and mitigation options and show their
20 respective potential negative implications for the complementary goal.

21 22 23 **5.7. Uncertainties and Data Gaps**

24
25 This chapter has updated knowledge about the impacts of climate change on the coastal ecosystems including a
26 better understanding of the coastal ecosystems not on their own right but also from the impacts of overexploitation
27 and habitat destruction that have been responsible for most of the historical changes. There is a better understanding
28 of the varying impacts of weather and climate extremes and long-term sea-level rise on human systems. For
29 example, the projected increase in both tropical storm intensity and population in the next 20 years can expect to
30 greatly increase the number of people exposed and exacerbate disaster risk (Knutson *et al.*, 2010). While knowledge
31 has increased it has yet to provide a complete understanding of new issues, e.g., ocean acidification impacts, or old
32 issues, e.g., adaptation costs and options. The complexity of adaptation issues, especially involving interacting with
33 human systems has been discussed. Adaptation has widened in scope to cover areas where policymakers would like
34 to have more information to act in future, e.g., vulnerability assessment, costs of adaptation and more adaptation
35 options.

36
37 Although a better understanding of shoreline response to future sea-level rise has been made by recent
38 improvements in technology (e.g., satellite imagery) to investigate and characterize large-scale changes in shoreline,
39 quantitative predictions of future coastal change remains difficult. This is due to the complexity of coastal systems,
40 influence of infrequent storm events and insufficient understanding of coastal systems over decadal timescales.
41 Shoreline response is more complicated than simple drowning alone because of factors such as sediment supply,
42 offshore geology, engineering structures, and wave forcing (Ashton *et al.*, 2008). For example, for many
43 sedimentary coasts, one fundamental question is the sediments and rate of sedimentation in response to sea-level
44 rise. In the long-term, we “need to eventually develop the capability to predict at least a regionally averaged
45 shoreline response to a given change in the rate of sea-level rise” (Ashton *et al.*, 2008: 737).

46
47 Although sea level is predicted to rise in future, there are uncertainties in evaluating the historical changes, modeling
48 future climatic change and estimating site-specific impacts. Many SLR assessments are not at spatial or temporal
49 scales most relevant for decision makers who required information on baseline conditions and projections of change
50 (Kettle, 2012). The local data required for SLR assessment are also not easily available. For example, LIDAR data
51 are only easily available for the USA coasts (NOAA Digital Coast Data Access Viewer website) but not for the rest
52 of the world yet. Only when such data become available in future, many developing countries, especially low-lying
53 countries, the deltaic areas of Asia and small island states could better assess the impact of sea-level rise.

1 There are significant gaps in vulnerability assessment of other specific coastal aspects. For example, climate
2 modeling of diseases that could affect the coastal areas is based mainly on the mean values of climate. There is a
3 need to incorporate effects of daily temperature variation into predictive models and show how that variation is
4 altered by climate change (Paaijmans *et al.*, 2010). Also, despite tourism as one of most important industries in the
5 coastal areas, not enough is known about tourists' likely behavioural reactions to projected climatic changes
6 (Moreno and Amelung, 2009).

7
8 The available vulnerability studies only explore a small fraction of the uncertainty. Generally, studies do not
9 consider the full range of possible relative sea-level changes and often exclude a potential large contribution of ice
10 sheet melting to sea-level rise, regional variations in climate-induced sea-level change and local factors such as
11 human-induced subsidence. Many studies rely on few or only a single socio-economic scenario and exclude
12 adaption or consider only a few stylized options. The available studies also only cover a limited range of impacts.
13 Integrated studies considering various impacts and their interdependence are rare. In particular, studies either focus
14 on land loss due to submergence or on flood damage. Generally, few studies explore indirect impacts and indirect
15 costs.

16
17 A wide range of coastal management framework and measures is available and used in coastal adaptation to climate
18 change, and their scope of integration has increased by combining scenarios of climate change and socio-economic
19 conditions and risk assessment (Kirshen *et al.*, 2011). While various adaptation measures are available, at the local
20 level, apart from adaptation options such as dykes and beach nourishment, there is not enough information on
21 assessment of adaptation options. Knowledge gaps exist, data are missing or their reliability is insufficient. In some
22 cases, alternatives are clear, e.g., giant floodgates or floating houses and amphibious housing (e.g., UK,
23 Netherlands). For many developing countries with narrow coastal areas and small island nations, the issue of coastal
24 squeeze becomes an increasing pertinent issue as the coastal ecosystems are drowned and cannot migrate inland
25 because of coastal protection measures or coastal communities cannot move inland.

26
27 Of various adaptation approaches to climate change, the integrated coastal management (ICM) has developed as an
28 effective framework and been able achieve a number of goals : the minimization of risks and impacts from coastal
29 hazards, economic development and use of coastal resources, and protection of coastal environmental resources,
30 natural assets, and ecosystems. However, the ICM still faces the limitation and uncertainty of the longer time frames
31 for sea-level rise and ocean acidification, the potential for physical and ecological thresholds or tipping points, and
32 the long lead times often required for making changes in coastal management, due to system lags in socioeconomic
33 systems.

34
35 There is an increasing trend to merge the practice of DRR (disaster risk reduction) and CCA (climate change
36 adaptation) particularly in developing countries (Berse *et al.*, 2011; CCD 2009). However, DRR differs from CCA
37 in spatial and temporal scales, knowledge and norms and DRR goals, strategies and measures need to be revised or
38 modified to meet the goals of CCA more effectively (Birkman and von Teichman, 2010).

39
40 The coastal zone has developed a body of research on the costs and benefits of adaptation options (Argawala and
41 Fankhauser, 2008). There is a continued evolution on the assessment of costs and benefits of adaptation options with
42 a wider range of scenarios, expanded impacts considered and integrated adaptation options. However, several key
43 issues in methodological development still exist: these include the determination of baseline conditions; treatment of
44 uncertainty and equity including distributional impacts, ancillary benefits and public-private efforts and economic
45 valuation (UNFCCC, 2010a).

46
47 Developing a knowledge platform for adaptation with communication between scientists, policy makers,
48 stakeholders and the general public could be considered as a priority area for coastal areas of a large or regional area
49 affected by climate change and sea-level rise. This is well developed in European Union (European Commission
50 Climate Action website), the Mediterranean (PAP website) and Australia (OzCoasts website), and but less so in the
51 developing countries, except in certain regions, e.g. Caribbean islands (CCCCC website), Pacific Islands (SPREP
52 website). An Adaptation Knowledge Platform has been developed for Asia-Pacific (Adaptation Knowledge Platform
53 website) but no coastal portal is available for Southeast Asia and East Asia.

1 Lastly, coastal research relating to climate change needs to be positioned in a proper context and in line of what has
2 been noted in the 21st century. Based on Science Citation Index, Li *et al.* (2011) concluded that temperature,
3 environment, precipitation, greenhouse gas, risk and biodiversity will be the foci of climate research in the 21st
4 century. The implications for coasts would be on biodiversity and flooding which is more coast-bound. Future
5 technological advances can be significant, e.g., new forms of energy and food production, information and
6 communication technology (ICT) for risk monitoring (Delta Commission, 2008). This would be useful for flood
7 risks and food production in deltas and coastal systems (aquaculture).
8
9

10 **5.8. Conclusion**

11
12 Since the AR4, there has been much research on the impacts of climate change on the coasts. While the observed
13 and future increase in weather and climate extremes can be variable, the rate of sea-level rise would seem to be
14 critical to many issues related to the response of both coastal ecosystems and human systems.
15

16 In some way, the human systems in the coastal areas are critical than the coastal ecosystems considering the fact that
17 they exacerbate the impacts of climate change on the coastal ecosystems. At the same time, it is evident that
18 increased vulnerability and exposure to climate change and sea-level rise would be exacerbated by rapid population
19 growth and increased urbanization in the LECZ. Such hotspots would be in the developing world, particularly, the
20 megadeltas in Asia and small island states.
21

22 Some assessments of vulnerabilities and costs for coasts at the global, regional and the national level are available.
23 But these assessments have uncertainties and limitations and not suitable for local areas to take appropriate action.
24 While some local areas have relied on traditional practices or use current adaptation measures, many have
25 difficulties in deciding what appropriate or effective options can be made for the future.
26

27 Despite various problems, issues and limits to which natural and human coastal systems can adapt, adaptation will
28 remain the predominant approach to reduce climate risks to coastal communities. While costs assessments have yet
29 to be fully developed for easy application, adaptation has to be taken now. As coastal areas are also affected by non-
30 climate related disasters, e.g., earthquakes, tsunamis, there is increasing options for climate change adaptation
31 (CCA) to incorporate disaster risk reduction (DRR) to maintain sustainable coastal communities.
32
33

34 **Frequently Asked Questions**

35 ***FAQ 5.1: How does climate change affect coastal ecosystems?***

36 The major global stressors, mostly caused by increased concentration of CO₂ in the atmosphere, affecting coastal
37 ecosystems are extreme weather and climate events, sea-level rise, ocean warming and ocean acidification. The
38 impact of sea level is mostly related to the capacity by animals (e.g. corals) and plants (e.g. mangroves) to keep up
39 with the vertical rise of the sea. Warming affects all organisms, increasing their metabolism and causing mass
40 mortality events of those living close their upper thermal limit. Ocean acidification negatively impacts many
41 organisms that build shells and skeletons but its effects are poorly known at the ecosystem level.
42
43

44 ***FAQ 5.2: How is climate change contributing to coastal erosion?***

45 Erosion is the process of wearing away material from the coastal profile due to imbalance in the supply and export
46 of material from a certain section: dunes, beaches, cliffs, etc. It is mainly caused by winds, waves and sea level. Any
47 rise in mean sea level will result in a landward and upward displacement of the cross-shore seabed profile and a
48 retreat of the shoreline. Increasing waves heights can cause the sand bars to move seawards and high storm surges
49 (sea levels) also produce an offshore movement of sand due to non-equilibrium in the profile. Higher waves and
50 surges may increase the probability of sand barriers and dunes overwash or breaching. Changes in wave direction or
51 propagation due to changing wave heights or sea levels may result in increasing gradient in the sediment transport
52 rate and consequently erosion.
53

54 ***FAQ 5.3: Does adequate planning of the coastal uses contributes to reduce climate change impacts?***

1 Yes, adequate planning of the coastal uses contributes to reduce climate change impacts. Such planning is normally
2 supported by national legislation and considers both the problems of both climate change and coastal hazards,
3 especially coastal flooding. Regional coastal strategies and plans are established with guidelines for local
4 governments to implement. For measures to be taken, the focus is on precautionary measures irrespective of future
5 climate change. An important paradigm change of planning land uses to reduce climate change is to use the buffer
6 zone as a response to coastal inundation. The strategy is to work with nature rather than against nature, e.g. in the
7 Netherlands.

8
9 **FAQ 5.4: Is the sea level rising equally in all regions?**

10 No, there are spatial variations in sea-level changes that can add to or subtract from the global average rise. The
11 spatial variations can result from a variety of processes such as ocean circulation, where sloping sea level balances
12 the Coriolis force; changes in seawater temperature, where warming of seawater causes it to expand and raise sea
13 level in a thermal expansion; and changes in gravity, where the loss of mass from ice sheet melting changes the local
14 gravity field and seawater moves away, raising sea level at distant locales.

15
16 **FAQ 5.5: What climate change impacts are getting more severe for the coasts?**

17 The primary coastal impact becoming more severe is elevated water levels due to sea-level rise (SLR). SLR not only
18 inundates low-lying land, like filling a bath tub, but can contribute to dynamic changes in sandy beaches, through
19 shifting the beach location landward and higher, and in wetlands, by drowning marsh vegetation. Scientists disagree
20 on whether tropical cyclones will become more intense and/or frequent in the future. However, for some coasts,
21 where beaches cannot shift landward and upward, storm surges will increasingly impact coastal development due to
22 their superposition on top of a rising sea.

23
24 **FAQ 5.6: How can coastal communities adapt to climate change impacts?**

25 Various adaptation options are available which range from accommodation to retreat. Adaptation through structural
26 measures include hard (e.g. sea walls) and soft (e.g. coastal revegetation) management options. Non-structural
27 measures include land-use planning (e.g. rolling easements that require relocation of vulnerable infrastructure as
28 critical risk thresholds are crossed). Education is also important for building community resilience. Risk transfer
29 mechanisms (e.g. insurance) address residual risk although where risks are too high, retreat from coastal areas may
30 be the only viable response. A combination of strategies, tailored to suit the particular coastal community, may be
31 required and will need to be reviewed and adjusted as circumstances change in the future.

32
33
34 **References**

- 35
36 Abal, E. G., W. C. Dennison, Seagrass Depth Range and Water Quality in Southern Moreton Bay,
37 **Abuodha, P. A. O. and C. D. Woodroffe (2010):** Assessing vulnerability to sea-level rise using a coastal sensitivity
38 index: a case study from southeast Australia. *Journal of Coastal Conservation* **14(3)**: 189-205.
39 **Acker, J. et al.,** *Ieee Geoscience and Remote Sensing Letters* **6**, 209 (2009).
40 **Adams P.N., D.L. Inman and J.L. Lovering, 2011:** Effects of climate change and wave direction on longshore
41 sediment transport patterns in Southern California. *Climatic Change*, **109, Suppl. 1**: 211-228.
42 **Adey, W. H., R. S. Steneck, 2001:** Thermogeography over time creates biogeographic regions: a
43 temperature/space/time-integrated model and an abundance-weighted test for benthic marine algae. *Journal of*
44 *Phycology* **37**, 677-698.
45 **Adger, N., 2006:** Vulnerability. *Global Environmental Change*, **16(3)**, 268-281.
46 **Adger, N., J. Barnett, F. Chapin, and H. Ellemor, 2011:** This must be the place: Under-representation of identity and
47 meaning in climate change decision-making. *Global Environmental Politics* **11 (2)**:1-25.
48 **Adger, W. N., I. Lorenzoni, and K. L. O'Brien eds. 2009a:** *Adapting to Climate Change: Thresholds, Values,*
49 *Governance.* Cambridge University Press.
50 **Adger, W. N., S. Dessai, M. Goulden, M. Hulme, I. Lorenzoni, D. Nelson, L.-O. Naess, J. Wolf, and A. Wreford,**
51 **2009b:** Are there social limits to adaptation to climate change? *Climatic Change* **93 (3-4)**:335 - 354.
52 **Adjeroud, M., F. Michonneau, P. J. Edmunds, Y. Chancerelle, T. Lison de Loma, L. Penin, L. Thibaut, J. Vidal-**
53 **Dupiol, B. Salvat, R. Galzin, 2009:** Recurrent disturbances, recovery trajectories, and resilience of coral
54 assemblages on a South Central Pacific reef. *Coral Reefs* **28**, 775-780.

- 1 **Aerts, J., T. Sprong, and B. Bannink** 2008: eds. *Attention to Safety. BSIK Klimaat foor Ruimte*. DG Water. The
2 Hague, The Netherlands, Report 009/2008 (in Dutch).
- 3 **Allen, J. R. L.** 1995: Salt-marsh growth and fluctuating sea level: implications of a simulation model for Flandrian
4 coastal stratigraphy and peat-based sea-level curves. *Sedimentary Geology* **100**, 21–45.
- 5 **Allen, J.R.L.,** 2000: Morphodynamics of holocene salt marshes: A review sketch from the atlantic and southern
6 north sea coasts of europe. *Quaternary Science Reviews*, **19(12)**, 1155-1231.
- 7 **Alongi, D. M.,** 2002: Present state and future of the world's mangrove forests. *Environmental Conservation* **29**
8 **(3)**:331-349.
- 9 **Alongi, D. M.,** 2008: Mangrove forests: Resilience, protection from tsunamis, and response to global climate
10 change. *Estuarine, Coastal and Shelf Science* **76**: 1-13.
- 11 **Alongi, D. M., A. D. McKinnon,** 2005: The cycling and fate of terrestrially-derived sediments and nutrients in the
12 coastal zone of the Great Barrier Reef shelf. *Marine Pollution Bulletin* **51**, 239-252.
- 13 **Alongi, D. M., A. I. Robertson,** 1995: Factors regulating benthic food chains in tropical river deltas and continental
14 shelf areas. *Geo-Marine Letters* **15**, 145-152 .
- 15 **Alvarez-Filip, L., M. Millet-Encalada, H. Reyes-Bonilla,** 2009: Impact of Hurricanes Emily and Wilma on the
16 Coral Community of Cozumel Island, Mexico: *Bulletin of Marine Science* **84**, 295-306.
- 17 **Alvarez-Filip, L., I. Gil,** 2006: Effects of Hurricanes Emily and Wilma on coral reefs in Cozumel, Mexico. *Coral*
18 *Reefs* **25**, 583.
- 19 **Amundsen, H., F. Berglund, and H. Westskog,** 2010: Overcoming barriers to climate change adaptation - a question
20 of multilevel governance? *Environment and Planning C: Government and Policy* **28 (2)**:276-289.
- 21 **Andersen, T.J., S. Svinth, and M Pejrup,** 2011: Temporal variation of accumulation rates on a natural salt marsh in
22 the 20th century - The impact of sea level rise and increased inundation frequency. *Marine Geology* **279**, 178–
23 187.
- 24 **Andersson A. J., Mackenzie F. T. and Gattuso J.-P.,** 2011: *Effects of ocean acidification on benthic processes,*
25 *organisms, and ecosystems.* In: Gattuso J.-P. and Hansson L. (Eds.), *Ocean acidification*, pp. 122-153. Oxford:
26 Oxford University Press.
- 27 **Anthoff, D., R.J. Nicholls, and R.S.J. Tol,** 2010: The economic impact of substantial sea-level rise. **15(4)**, 321-335.
- 28 **Anthony K. R. N., Kline D. L., Diaz-Pulido G., Dove S. and Hoegh-Guldberg O.,** 2008: Ocean acidification causes
29 bleaching and productivity loss in coral reef builders. *Proceedings of the National Academy of Science U.S.A.*
30 **105**:17442–17446.
- 31 **Anthony, A., J. Atwood, P. August, C. Byron, S. Cobb, C. Foster, C. Fry, A. Gold, K. Hagos, L. Heffner, D.Q.**
32 **Kellogg, K. Lellis-Dibble, J.J. Opaluch, C. Oviatt, A. Pfeiffer-Herbert, N. Rohr, L. Smith, T. Smythe, J. Swift,**
33 **and N. Vinhateiro,** 2009: Coastal lagoons and climate change: Ecological and social ramifications in US atlantic
34 and gulf coast ecosystems. *Ecology and Society*, **14(1(8))**.
- 35 **Anton, C., J. Young, P.A. Harrison, M. Musche, G. Bela, C.K. Feld, R. Harrington, J.R. Haslett, G. Pataki, M.D.A.**
36 **Rounsevell, M. Skourtos, J.P. Sousa, M.T. Sykes, R. Tinch, M. Vandewalle, A. Watt, and J. Settele,** 2010:
37 Research needs for incorporating the ecosystem service approach into EU biodiversity conservation
38 policy. **19(10)**, 2979-2994.
- 39 **Aoyama, M., H. Goto, H. Kamiya, I. Kaneko, S. Kawae, H. Kodama, Y. Konishi, K.-I. Kusumoto, H. Miura, E.**
40 **Moriyama, K. Murakami, T. Nakano, F. Nozaki, D. Sasano, T. Shimizu, H. Suzuki, Y. Takatsuki and**
41 **A.Toriyama,** 2008: Marine biogeochemical response to a rapid warming in the main stream of the Kuroshio in
42 the western North Pacific *Fisheries Oceanography* **17**, 206.
- 43 **Argawala, S. and S. Fankhauser,** 2008: Economic aspects of adaptation to climate change: Costs, benefits, and
44 policy instruments. *OECD*, Paris.
- 45 **Ashton, A.D., J.P. Donnelly, and R.L. Evans** 2008: A discussion of the potential impacts of climate change on the
46 shorelines of the Northeastern USA. *Mitigation and Adaptation Strategies for Global Change* **13**: 719-743.
- 47 **Aunan, K., and B. Romstad,** 2008: Strong coasts and vulnerable communities: Potential implications of accelerated
48 sea-level rise for Norway. *Journal of Coastal Research* **24(2)**, 403-409
- 49 **Azzellino, A., S. A. Gaspari, S. Airoidi, C. Lanfredi,** 2008: Biological consequences of global warming: does sea
50 surface temperature affect cetacean distribution in the western Ligurian Sea? *J. Mar. Biol. Assoc. U.K.* **88**,
51 1145-1152.
- 52 **Badjeck, M-C., E.H. Allison, A.S. Halls, and N.K. Dulvy** 2010: Impacts of climate variability and change on
53 fishery-based livelihoods. *Marine Policy* **34**, 375-383.

- 1 **Baker**, A. C., P. W. Glynn, B. Riegl, 2008: Climate change and coral reef bleaching: An ecological assessment of
2 long-term impacts, recovery trends and future outlook. *Estuarine Coastal and Shelf Science* **80**, 435.
- 3 **Bakun** A., 1990. Global climate change and intensification of coastal ocean upwelling. *Science* **247**: 198-201.
- 4 **Ball** MC, Pidsley SM., 1995: Growth responses to salinity in relation to distribution of two mangrove species,
5 *Sonneratia alba* and *S. lanceolata*. *Functional Ecology* **9**:77–85.
- 6 **Ball**, M. C., 1998: Mangrove Species Richness in Relation to Salinity and Waterlogging: A Case Study Along the
7 Adelaide River Floodplain, Northern Australia. *Glob. Ecol. Biogeogr. Lett.* **7**, 73-82.
- 8 **Balmford**, A., B. Fisher, R. E. Green, R. Naidoo, B. Strassburg, R. K. Turner and A. S. L. Rodrigues 2011:
9 Bringing Ecosystem Services into the Real World: An Operational Framework for Assessing the Economic
10 Consequences of Losing Wild Nature. *Environmental and Resource Economics* **48(2)**, 161-175.
- 11 **Bancroft**, W.J., Garkaklis, M.J. & Roberts, J. D., 2004: Continued expansion of the Wedge-tailed Shearwater,
12 *Puffinus pacificus*, nesting colonies on Rottnest Island, Western Australia. *Emu Austral Ecology* **104**: 79-82.
- 13 **Barale**, V. and E. Özhan 2010: Advances in integrated coastal management for the Mediterranean and Black Sea.
14 *Journal of Coast Conservation* **14**: 249–255.
- 15 **Barange**, M. and R.I. Perry 2009: Physical and ecological impacts of climate change relevant to marine and inland
16 capture fisheries and aquaculture. In *Climate Change Implications for Fisheries and Aquaculture: Overview of*
17 *Current Scientific Knowledge* [Cochrane, K., C. De Young, D. Soto, and T. Bahri (eds)]. FAO Fisheries and
18 Aquaculture Technical Paper n° **530**. Rome, FAO. pp. 7–106.
- 19 **Barbier**, E. B., E. W. Koch, B. R. Silliman, S. D. Hacker, E. Wolanski, J. Primavera, E. F. Granek, S. Polasky, S.
20 Aswani, L. A. Cramer, D. M. Stoms, C. J. Kennedy, D. Bael, C. V. Kappel, G. M. E. Perillo, and D. J. Reed.
21 2008. Coastal Ecosystem-Based Management with Nonlinear Ecological Functions and Values. *Science* **319**
22 (5861):321-323.
- 23 **Barbier**, E.B., 2012: Progress and challenges in valuing coastal and marine ecosystem services. *Review of*
24 *Environmental Economics and Policy* **6(1)**, 1-19.
- 25 **Baric**, A., Grbec, B., Bogner, D. 2008: Potential Implications of Sea-Level Rise for Croatia. *J. Coastal Res.* **24(2)**
26 299-305.
- 27 **Barlow** P.M., and Reichard, E.G. 2010: Saltwater intrusion in coastal regions of North America. *Hydrogeology*
28 *Journal*, **18**: 247–260.
- 29 Barnett and Campbell, 2010
- 30 **Barry** J. P., Widdicombe S. and Hall-Spencer J. M., 2011: Effects of ocean acidification on marine biodiversity and
31 ecosystem function. In: Gattuso J.-P. and Hansson L. (Eds.), *Ocean acidification*, pp. 192-209. Oxford: Oxford
32 University Press.
- 33 **Barry**, J. P., C. H. Baxter, R. D. Sagarin, S. E. Gilman, 1995: Climate-Related, Long-Term Faunal Changes in a
34 California Rocky Intertidal Community. *Science* **267**, 672-675.
- 35 **Barton**, A., Hales, B., Waldbusser, G. G., Langdon, C., & Feely, R. A., 2012: The Pacific oyster, *Crassostrea gigas*,
36 shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean
37 acidification effects. *Limnology and Oceanography*, **57(3)**, 698–710.
- 38 **Baulcomb**, Corinne. 2011: *Review of the Evidence Linking Climate Change to Human Health for Eight Diseases of*
39 *Tropical Importance*. Land Economy Working Paper Series No **63**. Scottish Agricultural College. Available at:
40 <http://www.sac.ac.uk/mainrep/pdfs/leergworkingpaper63.pdf>
- 41 **Beare**, D. J. F. Burns, A. Greig, E. G. Jones, K. Peach, M. Kienzle, E. McKenzie, D. G. 2004: Reid Long-term
42 increases in prevalence of North Sea fishes having southern biogeographic affinities. *Mar. Ecol.-Prog. Ser.* **284**,
43 269-278.
- 44 Beaugrand, G., P. C. Reid, F. Ibanez, J. A. Lindley, M. Edwards, *Science* **296**, 1692 (May, 2002).
- 45 Beaugrand, G., P. C. Reid, *Global Change Biology* **9**, 801 (Jun, 2003).
- 46 Beaugrand, G., C. Luczak, M. Edwards, *Global Change Biology* **15**, 1790 (Jul, 2009).
- 47 **Bedsworth**, L. W., and E. Hanak, 2010: Adaptation to Climate Change: A Review of Challenges and Tradeoffs in
48 Six Areas. *J. Am. Plann. Assoc.* **76(4)**, 477-495.
- 49 **Berkelmans** R. and van Oppen M. J. H., 2006: *The role of zooxanthellae in the thermal tolerance of corals: a*
50 *'nugget of hope' for coral reefs in an era of climate change*. Proceedings of the Royal Society of London.
51 Series B: Biological Sciences **273**: 2305-2312.
- 52 **Berse**, K., P. Tran and J. Velasquez 2011: *At the Crossroads – Climate Change Adaptation and Disaster Risk*
53 *Reduction in Asia and the Pacific*. UNISDR Asia and Pacific Secretariat, 162 pp.

- 1 **Bertness, M. D., S. C. Pennings**, in *Concepts and Controversies in Tidal Marsh Ecology*, M. P. Weinstein, D. A.
2 Kreeger, Eds. (Kluwer Academic Publishers, Dordrecht, 2000), pp. 39–57.
- 3 **Bessat F. and Buigues D.**, 2001: Two centuries of variation in coral growth in a massive Porites colony from
4 Moorea (French Polynesia): a response of ocean-atmosphere variability from south central Pacific.
5 *Palaeogeography*, **175(1-4)**: 381-392.
- 6 **Biesbroek R., Klostermann J., Termeer C., Kabat P.**, 2011: Barriers to climate change adaptation in the Netherlands.
7 *Climate Law* **2**: 181-199.
- 8 **Biesbroek, G. R., R. J. Swart, and W. G. M. van der Knaap**, 2009: The mitigation-adaptation dichotomy and the
9 role of spatial planning. *Habitat International* **33 (3)**:230-237.
- 10 **Biggs D.**, 2011: *Case study: the resilience of the nature-based tourism system on Australia's Great Barrier Reef*.
11 Report prepared for the Australian Department of Sustainability, Environment, Water, Population and
12 Communities on behalf of the State of the Environment 2011 Committee Government. 32 p.
- 13 **Bird E.C.F.** 2000: *Coastal Geomorphology: An Introduction*. John Wiley, Chichester, 322 pp.
- 14 **Birkemeier, W.A.** 1985: Field Data on Seaward Limit of Profile Change. *Journal of Waterway, Port, Coastal and*
15 *Ocean Engr.* Vol **111** No **3**.
- 16 **Birkman, J. and K. von Teichman** 2010: Integrating disaster risk reduction and climate change adaptation: key
17 challenges – scales, knowledge and norms. *Sustainability Science* **5**: 171-184.
- 18 **Bjarnadottir, S., Y. Li, and M.G. Stewart**, 2011: Social vulnerability index for coastal communities at risk to
19 hurricane hazard and a changing climate. *Natural Hazards* **59(2)**, 1055-1075.
- 20 Blackburn, S. I., G. Cresswell, *Australian Journal of Marine and Freshwater Research* **44**, 253 (1993).
- 21 Blasco, F., P. Saenger, E. Janodet, *Catena* **27**, 167 (1996).
- 22 Blomqvist, S., M. Peterz, *Mar. Ecol.-Prog. Ser.* **20**, 85 (1984).
- 23 **Blum, M.D. and H.H. Roberts**, 2009: Drowning of the Mississippi Delta due to insufficient sediment supply and
24 global sea-level rise. *Nature Geoscience* **2**, 488–491.
- 25 **Boateng, I.** 2012a: GIS assessment of coastal vulnerability to climate change and coastal adaption planning in
26 Vietnam. *Journal of Coastal Conservation* **16**: 25–36
- 27 **Boateng, I.** 2012b: An assessment of the physical impacts of sea-level rise and coastal adaptation: a case study of
28 the eastern coast of Ghana. *Climatic Change*.
- 29 **Bocanegra, E., Da Silva Jr., G.C. Custodio, E., Manzano, M. and Montenegro, S.**, 2010: State of knowledge of
30 coastal aquifer management in South America, *Hydrogeology Journal* **18**: 261–267
- 31 **Boden, T. G. Marland, and B. Andres**, 2011: *Global CO2 Emissions from Fossil-Fuel Burning, Cement*
32 *Manufacture, and Gas Flaring: 1751-2008*. Carbon Dioxide Information Analysis Center, Oak Ridge National
33 Laboratory, Oak Ridge, TN. Emissions data available at:
34 http://cdiac.ornl.gov/ftp/ndp030/global.1751_2008.ems.
- 35 **Boehlert, G. W., and A. B. Gill.**, 2010: *Environmental and Ecological Effects of Ocean Renewable Energy*
36 *Development: A Current Synthesis*. Rockville, MD: The Oceanography Society. Available at:
37 <http://hdl.handle.net/1957/16152>.
- 38 **Bollmann, M., T. Bosch, F. Colijn, R. Ebinghaus, R. Froese, K. Güssow et al** 2010: *World Ocean Review 2010*.
39 Maribus, Hamburg, 232 pp.
- 40 **Bongaerts, P., T. Ridgway, E. Sampayo and O. Hoegh-Guldberg** (2010): Assessing the “deep reef refugia”
41 hypothesis: focus on Caribbean reefs. *Coral Reefs*, **29(2)**, 309-327.
- 42 Booth, D. T., C. Freeman, *Coral Reefs* **25**, 629 (2006).
- 43 **Borges A. V.** 2005: Do we have enough pieces of the jigsaw to integrate CO2 fluxes in the coastal ocean? *Estuaries*
44 **28**:3–27
- 45 **Borges A. V. and Abril G.**, 2011: Carbon dioxide and methane dynamics in estuaries. In: Wolanski E. M., D (Ed.),
46 *Treatise on Estuarine and Coastal Science*, pp. 119-161. Waltham: Academic Press.
- 47 **Borges A. V. and Gypens N.** 2010: Carbonate chemistry in the coastal zone responds more strongly to
48 eutrophication than to ocean acidification. *Limnology and Oceanography* **55**:346-353.
- 49 Borum, J. *et al.*, *J. Ecol.* **93**, 148 (2005).
- 50 **Bosom, E. and J.A. Jimenez**, 2011: Probabilistic coastal vulnerability assessment to storms at regional scale -
51 application to Catalan beaches (NW mediterranean). *Natural Hazards and Earth System Sciences*, **11(2)**, 475-
52 484.
- 53 Boteng, 2012
- 54 Boudouresque, C. F., G. Bernard, G. Pergent, A. Shili, M. Verlaque, *Botanica Marina* **52**, 395 (2009).

- 1 **Bouillon**, S., V.H. Rivera-Monroy, R.R. Twilley and J.G. Kairo J. G., 2009: Mangroves. In: D. Laffoley & G.
2 Grimsditch (eds.). *The Management of Natural Coastal Carbon Sinks*, pp. 13-20. Gland, Switzerland: IUCN.
- 3 **Bouwman** L. Klein Goldewijk K, Van Der Hoek K.W. Beusena A.H.W., Van Vuurena, D.P., Willems J., Rufino
4 M.C. and E Stehfest (2011): *Exploring global changes in nitrogen and phosphorus cycles in agriculture*
5 *induced by livestock production over the 1900–2050 period*. PNAS pnas.
- 6 **Brakenridge**, G.R., Syvitski, J.P.M., Overeem, I., Higgins, S.A., Kettner, A.J., Stewart-Moore, J.A., Westerhoff, R.,
7 2012: Global Mapping of Storm Surges and the Assessment of Delta Vulnerability. *Natural Hazards*, special
8 issue on storm surges.
- 9 **Brander**, L.M., R.J.G.M. Florax, and J.E. Vermaat, 2006: The empirics of wetland valuation: A comprehensive
10 summary and a meta-analysis of the literature. *Environmental and Resource Economics*, **33(2)**, 223-250.
- 11 **Breitburg** D. L., Hondorp D. W., Davias L. A. and Diaz R. J., 2009. Hypoxia, nitrogen, and fisheries: integrating
12 effects across local and global landscapes. *Annual Reviews of Marine Science* **1**:329-349.
- 13 **Brito**, A., A. Newton, P. Tett, and T.F. Fernandes, 2010: Sediment and water nutrients and microalgae in a coastal
14 shallow lagoon, ria formosa (Portugal): Implications for the water framework directive. *Journal of*
15 *Environmental Monitoring*, **12(1)**, 318-328.
- 16 Bromberg Gedan et al., 2009
- 17 Bromberg Gedan et al., 2011
- 18 **Brown** K., E. L. Tompkins, and W. N. Adger., 2002: *Making Waves: Integrating Coastal Conservation and*
19 *Development*. London: Earthscan.
- 20 Bruce, A. J., v. A. P. Biol. 2, New York, pp. 37–94., in *Biology and Geology of Coral Reefs*, O. A. Jones, R.
21 Endean, Eds. (Academic Press, New York, 1976), vol. 3, pp. 37-94.
- 22 **Bruneau**, N., A.B. Fortunato, G. Dodet, P. Freire, A. Oliveira, and X. Bertin, 2011: Future evolution of a tidal inlet
23 due to changes in wave climate, sea level and lagoon morphology (obidos lagoon, Portugal). *Continental Shelf*
24 *Research*, **31(18)**, 1915-1930.
- 25 **Bruno** J. F. and Selig E. R., 2007: *Regional decline of coral cover in the Indo-Pacific: timing, extent, and*
26 *subregional comparisons*. PLoS One 2:e711.
- 27 **Bruno** John F., Hugh Sweatman, William F. Precht, Elizabeth R. Selig, and Virginia G. W. Schutte, 2009:
28 Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* **90**:1478–1484.
- 29 **Buddemeier** R.W. and Smith S.V., 1988: Coral reef growth in an era of rapidly rising sea level: predictions and
30 suggestions for long-term research. *Coral Reefs* **7**: 51-56.
- 31 **Bunce** M., K. Brown, and S. Rosendo, 2010a: Policy misfits, climate change and cross-scale vulnerability in coastal
32 Africa: how development projects undermine resilience. *Environmental Science and Policy* **13 (6)**:485-497.
- 33 **Bunce** M., S. Rosendo, and K. Brown, 2010b: Perceptions of climate change, multiple stressors and livelihoods on
34 marginal African coasts. *Environment, Development and Sustainability* **12 (3)**:407-440.
- 35 **Buranapratheprat**, A., T. Yanagi, K. O. Niemann, S. Matsumura, P. Sojisuporn, *Journal of Oceanography* **64**, 639
36 (2008).
- 37 **Burch**, S. 2010: Transforming barriers into enablers of action on climate change: Insights from three municipal case
38 studies in British Columbia, Canada. *Global Environmental Change* **20 (2)**:287-297.
- 39 **Burke** L., K. Reynter, M. Spalding and A. Perry 2011: *Reefs at Risk Revisited*. Washington, DC: World Resources
40 Institute, 114 p.
- 41 **Burt** J., Al-Harhi S. and Al-Cibahy A., 2011: Long-term impacts of coral bleaching events on the world's warmest
42 reefs. *Marine Environmental Research* **72**:225-229.
- 43 **Bustnes**, J. O., T. Anker-Nilssen, S. H. Lorentsen, *J. Ornithol.* **151**, 19 (Jan, 2010).
- 44 **Byrkjedal**, I., O. R. Godo, M. Heino, Northward range extensions of some mesopelagic fishes in the Northeastern
45 Atlantic, *Sarsia* **89**, 484-489 (2004).
- 46 **Bythell**, J. C., Z. M. Hillis-Starr, C. S. Rogers, *Mar. Ecol.-Prog. Ser.* **204**, 93 (2000).
- 47 **Cabaco**, S. and R. Santos, 2007: Effects of burial and erosion on the seagrass *Zostera noltii*. *Journal of Experimental*
48 *Marine Biology and Ecology*, **340(2)**, 204-212.
- 49 **Cahoon**, D.R., D.J. Reed, and J.W. Day, 1995: Estimating shallow subsidence in microtidal salt marshes of the
50 southeastern united states: Kaye and barghoorn revisited. *Marine Geology*, **128(1-2)**, 1-9.
- 51 **Cai** W.-J., Hu X., Huang W.-J., Murrell M. C., Lehrter J. C., Lohrenz S. E., Chou W.-C., Zhai W., Hollibaugh J. T.,
52 Wang Y., Zhao P., Guo X., Gundersen K., Dai M. and Gong G.-C., 2011: Acidification of subsurface coastal
53 waters enhanced by eutrophication. *Nature Geoscience* **4**:766–770.
- 54 **Cai**, F., Su, X., Liu, J., Li, B., and Lei, G. 2009: Coastal erosion in China under the condition of global climate

- 1 change and measures for its prevention, *Progress in Natural Science*, **19**, 415-426.
- 2 **Campbell**, A., V. Kapos, A. Chenery, N. Doswald, S.I. Kahn, M. Rashid, J. Scharlemann and B. Dickson 2009. The
3 linkages between biodiversity and climate change adaptation – a review of the recent scientific literature. In :
4 *Review of the Literature on the Links between Biodiversity and Climate Change: Impacts, Adaptation and*
5 *Mitigation* [Campbell, A., V. Kapos, J.P.W. Scharlemann, P. Bubba, A. Chenery, L. Coad, B. Dickson, N.
6 Doswald, M.S.I. Khan, F. Kershaw and M. Rashid, eds.], Secretariat of the Convention on Biological Diversity,
7 Montreal. Technical Series No. 42, p. 49-87.
- 8 Campbell, S. J., L. J. McKenzie, S. P. Kerville, *Journal of Experimental Marine Biology and Ecology* **330**, 455
9 (Mar, 2006).
- 10 **Canu**, D.M., C. Solidoro, G. Cossarini, and F. Giorgi, 2010: Effect of global change on bivalve rearing activity and
11 the need for adaptive management. *Climate Research*, **42(1)**, 13-26.
- 12 **Cardoso** P. G., Raffaelli D. and Pardal M. A., 2008: The impact of extreme weather events on the seagrass *Zostera*
13 *noltii* and related *Hydrobia ulvae* population. *Marine Pollution Bulletin* **56**:483-492.
- 14 **Caribbean Community Climate Change Centre** (CCCCC) website:
15 <http://www.caribbeanclimate.bz/cpacc/cpacc.html>
- 16 **Carlson Jr.**, P.R., Yarbrow, L.A., Kaufman, K.A., Mattson, R.A., 2010: Vulnerability and resilience of seagrasses to
17 hurricane and runoff impacts along Florida's west coast. *Hydrobiologia* **649**, 39-53.
- 18 Carlsson, P., E. Graneli, P. Tester, L. Boni, *Mar. Ecol.-Prog. Ser.* **127**, 213 (1995).
- 19 **Carpenter** K., E., Abrar M., Aeby G., Aronson R., B., Banks S., Bruckner A., Chiriboga A., Cortes J., Delbeek J.,
20 Charles, DeVantier L., Edgar G., J., Edwards A., J., Fenner D., Guzman H., M., Hoeksema B., W., Hodgson G.,
21 Johan O., Licuanan W., Y., Livingstone S., R., Lovell E., R., Moore J., A., Obura D., O., Ochavillo D., Polidoro
22 B., A., Precht W., F., Quibilan M., C., Reboton C., Richards Z., T., Rogers A., D., Sanciangco J., Sheppard A.,
23 Sheppard C., Smith J., Stuart S., Turak E., Veron J., E. N., Wallace C., Weil E. and Wood E., 2008. One-third
24 of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* 1159196-
25 1159196.
- 26 **Carrington** E., 2002. Seasonal variation in the attachment strength of blue mussels: causes and consequences.
27 *Limnology and Oceanography* **47**:1723-1733.
- 28 **Cartwright**, A. 2009: *Coastal vulnerability in the context of climate change: A South African perspective*.
29 [http://www.90x2030.org.za/oid%5Cdownloads%5CCoastal%20vulnerability%20South%20African%20perspec](http://www.90x2030.org.za/oid%5Cdownloads%5CCoastal%20vulnerability%20South%20African%20perspective.pdf)
30 [tive.pdf](http://www.90x2030.org.za/oid%5Cdownloads%5CCoastal%20vulnerability%20South%20African%20perspective.pdf)
- 31 Castro, P., *Mar. Biol.* **46**, 237 (1978).
- 32 Castro, P., *Symbiosis* **5**, 161 (1988).
- 33 **CCD**, Commission on Climate Change and Development 2009. *Closing the Gaps*. Stockholm, 80 pp.
- 34 **CCSP**, 2008: *Impacts of Climate Change and Variability on Transportation Systems and Infrastructure: Gulf Coast*
35 *Study, Phase I*. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global
36 Change Research. Department of Transportation, Washington, DC, USA, 445 pp.
- 37 **Chaibi**, M. and Sedrati, M., 2009, Coastal Erosion Induced by Human Activities: The Case of Two Embayed
38 Beaches on the Moroccan Coast, *Journal of Coastal Research*.
- 39 Chaloupka, M., N. Kamezaki, C. Limpus, *Journal of Experimental Marine Biology and Ecology* **356**, 136 (Mar,
40 2008).
- 41 Chambers, L. E. "The Impact of Climate on Little Penguin Breeding Success" (Bureau of Meteorology, 2004).
- 42 Chapman, P. M. (In Press). "Management of coastal lagoons under climate change." *Estuarine, Coastal and Shelf*
43 *Science*.
- 44 **Chen** C-TA and Borges AV. 2009 : Reconciling opposing views on carbon cycling in the coastal ocean: continental
45 shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂. *Deep-Sea Res. II* **56**:578–90
- 46 **Chen**, F., S. Miao, M. Tewari, J.-W. Bao, and H. Kusaka, 2011: A numerical study of interactions between surface
47 forcing and sea-breeze circulations and their effects on stagnation in the greater Houston area. *Journal of*
48 *Geophysical Research-Atmospheres*, 2011; DOI: 10.1029/2010JD015533
- 49 **Cheong**, S-M. 2011. Editorial. Guest editor, Special Issue: Coastal Adaptation. *Climatic Change* **106(1)**: 1-4.
- 50 **Cheong**, S-M., B. Silliman and P.P. Wong (2012): Managing uncertainties in coastal adaptation. *Conservation*
51 *Letters* (under revision).
- 52 **Chhatre** A., and A. Agrawal. 2009: *Trade-offs and synergies between carbon storage and livelihood benefits from*
53 *forest commons*. *Proceedings of the National Academy of Sciences* 106 (42):17667-17670.

- 1 **Chmura**, G. L., 2009: Tidal salt marshes. In: D. Laffoley & G. Grimsditch (eds.). *The Management of Natural*
2 *Coastal Carbon Sinks*, pp. 5-11. Gland, Switzerland: IUCN.
- 3 **Chou**, W.C., J.L. Wu, Y.C. Wang, H. Huang, F.C. Sung, and C.Y. Chuang 2010: Modeling the impact of climate
4 variability on diarrhea-associated diseases in Taiwan (1996-2007). *Science of the Total Environment* **409**, 43-
5 51.
- 6 **Christie**, P., K. Lowry, A. T. White, E. G. Oracion, L. Sievanen, R. S. Pomeroy, R. B. Pollnac, J. M. Patlis, and R.-
7 L. V. Eisma. 2005. Key findings from a multidisciplinary examination of integrated coastal management
8 process sustainability. *Ocean and Coastal Management* **48 (3-6)**:468-483.
- 9 **Chu**, Z.X., X.G. Sun, S.K. Zhai, and K.H. Xu, 1996: Changing pattern of accretion/erosion of the modern Yellow
10 River (Huanghe) subaerial delta, China: Based on remote sensing images. *Marine Geology*, **227**, 13–30.
- 11 **Chust**, G., A. Caballero, M. Marcos, P. Liria, C. Hernandez, and A. Borja, 2010: Regional scenarios of sea level rise
12 and impacts on basque (Bay of Biscay) coastal habitats, throughout the 21st century. *Estuarine, Coastal and*
13 *Shelf Science*, **87(1)**, 113-124.
- 14 **Cinner**, J.E., T.R. McClanahan, N.A.J. Graham, T.M. Daw, J. Maina, S.M. Stead, A. Wamukota, K. Brown, and O.
15 Bodin, 2012: Vulnerability of coastal communities to key impacts of climate change on coral reef
16 fisheries. *Global Environmental Change-Human and Policy Dimensions*, **22(1)**, 12-20.
- 17 **Ciscar**, J.C., A. Iglesias, L. Feyen, L. Szabo, D. Van Regemorter, B. Amelung, R. Nicholls, P. Watkiss, O.B.
18 Christensen, R. Dankers, L. Garrote, C.M. Goodess, A. Hunt, A. Moreno, J. Richards, and A. Soria, 2011:
19 Physical and economic consequences of climate change in europe **108(7)**, 2678-2683.
- 20 **Coelho**, C., R. Silva, F. Veloso-Gomes, and F. Taveira-Pinto, 2009: Potential effects of climate change on northwest
21 portuguese coastal zones. *ICES Journal of Marine Science*, **66(7)**, 1497-1507.
- 22 **Cole**, R. G., R. C. Babcock, V. Travers, *New Zealand Journal of Marine and Freshwater Research* **35**, 17 (2001).
- 23 **Coleman**, J.M., O.K. Huh, and D.W. Braud Jr., 2008: Wetland loss in world deltas. *Journal of Coastal Research* **24**,
24 1–14.
- 25 **Connell**, J. H., *Coral Reefs* **16**, S101 (1997).
- 26 **Conservation International**, 2008: *Economic Values of Corals Reefs, Mangroves, and Seagrasses. A Global*
27 *Compilation*. Center for Applied Biodiversity Science, Conservation International, Arlington, VA, USA, 35 pp.
- 28 **Cooper**, J.A.G. and O.H. Pilkey, 2004: Sea-level rise and shoreline retreat: Time to abandon the bruun rule. *Global*
29 *and Planetary Change*, **43(3-4)**, 157-171.
- 30 **Costello**, A. M. Abbas, A. Allen, S. Ball, S. Bell, R. Bellamy, et al. 2009: *Managing the health effects of climate*
31 *change*. *Lancet*, 373, 1693-1733.
- 32 **Craft**, C., J. Clough, J. Ehman, S. Joye, R. Park, S. Pennings, H. Guo, and M. Machmuller. 2009. Forecasting the
33 effects of accelerated sea-level rise on tidal marsh ecosystem services. *Front. Ecol. Environ* **7**:73-78.
- 34 **Crain** C. M., Halpern B. S., Beck M. W. and Kappel C. V., 2009: Understanding and managing human threats to the
35 coastal marine environment. *Year in Ecology and Conservation Biology 2009* **1162**:39-62.
- 36 **Criales**, M. M. et al., *Caribbean Journal of Science* **38**, 52 (2002).
- 37 **Croquer**, A., E. Weil, *Dis. Aquat. Org.* **87**, 33 (2009).
- 38 **Crossland** C. J., Baird D., Ducrotoy J.-P. and Lindeboom H. J., 2005: The coastal zone- a domain of global
39 interactions. In: *Coastal fluxes in the anthropocene*. [Crossland, C.J., H.H. Kremer, H.J. Lindeboom, J.I.
40 Marshall Crossland, and M.D.A. Le Tissier(eds.)]. Springer-Verlag, Berlin, pp. 1-37.
- 41 **Custodio**, E. 2010: Coastal aquifers of Europe: an overview. *Hydrogeology Journal* **18**: 269–280
- 42 **Cutter** S., B. Osman-Elasha, J. Campbell, S-M. Cheong, S. McCormick, R. Pulwarty, S. Supratid, and G. Ziervogel,
43 2012: Managing risks from climate extremes at the local level. In : *Managing the Risks of Extreme Events and*
44 *Disasters to Advance Climate Change Adaptation*. [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken,
45 K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. A
46 Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change University of
47 Cambridge Press, Cambridge, U.K., p. 291-338.
- 48 **Dai** A., Qian T., Trenberth K. E. and Milliman J. D., 2009: Changes in continental freshwater discharge from 1948
49 to 2004. *Journal of Climate* **22**:2773-2792.
- 50 **Dasgupta**, S., B. Laplante, C. Meisner, D. Wheeler, and J. Yan, 2009: The impact of sea level rise on developing
51 countries: a comparative analysis. *Climatic Change* **93(3-4)**, 379-388
- 52 **Dasgupta**, S., B. Laplante, et al., 2010: Exposure of developing countries to sea-level rise and storm surges.
53 *Climatic Change*.
- 54 **Daw**, T. Adger, W.N. Brown, K. 2009: Climate change and capture fisheries: potential impacts, adaptation and

- 1 mitigation. In K. Cochrane, C. De Young, D. Soto and T. Bahri (eds). *Climate change implications for fisheries*
2 *and aquaculture: overview of current scientific knowledge*. FAO Fisheries and Aquaculture Technical Paper No
3 530. Rome, FAO, pp.107-150.
- 4 **Dawson**, R.J., M.E. Dickson, R.J. Nicholls, J.W. Hall, M.J.A. Walkden, P.K. Stansby, M. Mokrech, J. Richards, J.
5 Zhou, J. Milligan, A. Jordan, S. Pearson, J. Rees, P.D. Bates, S. Koukoulas, and A.R. Watkinson, 2009:
6 Integrated analysis of risks of coastal flooding and cliff erosion under scenarios of long term change. *Climatic*
7 *Change* **95(1-2)**, 249-288.
- 8 **Day Jr**, J.W., J. Rybczyk, F. Scarton, A. Rismondo, D. Are, and G. Cecconi, 1999: Soil accretionary dynamics, sea-
9 level rise and the survival of wetlands in venice lagoon: A field and modelling approach. *Estuarine, Coastal*
10 *and Shelf Science*, **49(5)**, 607-628.
- 11 **Day**, J.W. and L. Giosan, 2008: Survive or subside? *Nature Geoscience* **1**, 156–157.
- 12 **Day**, J.W., C. Ibáñez, F. Scarton, D. Pont, P. Hensel, J. Day, and R. Lane, 2011: Sustainability of Mediterranean
13 deltaic and lagoon wetlands with sea-level rise: The importance of river input. *Estuaries and Coasts* **34**, 483–
14 493.
- 15 Dayton, P. K., M. J. Tegner, *Science* **224**, 283 (1984).
- 16 **De Bruin**, K., R.B. Dellink, A. Ruijs, L. Bolwidt, A. van Buuren, J. Graveland, R.S. de Groot, P.J. Kuikman, S.
17 Reinhard, R.P. Roetter, V.C. Tassone, A. Verhagen and E.C. van Ierland 2009: Adapting to climate change in
18 The Netherlands: an inventory of climate adaptation options and ranking of alternatives. *Climatic Change* **95**:
19 23-45.
- 20 De Carlo, E. H., D. J. Hoover, C. W. Young, R. S. Hoover, F. T. Mackenzie, *Applied Geochemistry* **22**, 1777 (2007).
- 21 **De Groot**, R.S., Wilson, M.A., Boumans, R.M.J., 2002 : A typology for the classification, description and valuation
22 of ecosystem functions, goods and services. *Ecological Economics* **41** : 393–408.
- 23 **De la Vega-Leinert**, A.C. and R.J. Nicholls 2008 : Potential implications of sea-level rise for Great Britain. *Jour. of*
24 *Coastal Research* **24**, 342-357.
- 25 **De Silva**, S.S. and D. Soto 2009: Climate change and aquaculture: potential impacts, adaptation and mitigation. In
26 *Climate Change Implications for Fisheries and Aquaculture: Overview of Current Scientific Knowledge*
27 [Cochrane, K., C. De Young, D. Soto, and T. Bahri (eds)] FAO Fisheries and Aquaculture Technical Paper No
28 530. Rome, FAO. pp. 151-212.
- 29 **De'ath** G., Lough J. M. and Fabricius K. E., 2009: Declining coral calcification on the Great Barrier Reef. *Science*
30 **323**:116-119.
- 31 **Defeo**, O., A. McLachlan, D.S. Schoeman, T.A. Schlacher, J. Dugan, A. Jones, M. Lastra, and F. Scapini, 2009:
32 Threats to sandy beach ecosystems: A review. *Estuarine, Coastal and Shelf Science*, **81(1)**, 1-12.
- 33 Delange, W. P., P. J. Delange, *J. Coast. Res.* **10**, 539 (1994).
- 34 **DeLaune**, R. D., and J. R. White, 2011: Will coastal wetlands continue to sequester carbon in response to an
35 increase in global sea level? a case study of the rapidly subsiding Mississippi river deltaic plain. *Climatic*
36 *Change* online first; DOI 10.1007/s10584-011-0089-6.
- 37 **Delta Commission**, 2008: *Working together with water : Summary and conclusions*. 23 pp.
- 38 **Department of Climate Change**, 2009: *Climate Change Risks to Australia's Coast. A First Pass National*
39 *Assessment*. Report published by the Department of Climate Change, Australian Government, Canberra,
40 Australia, 172 pp.
- 41 **Desantis**, L.R.G., Bhotika, S., Williams, K., and Putz, F.E. 2007: Sea-level rise and drought interactions accelerate
42 forest decline on the Gulf Coast of Florida, USA *Glob Change Biol* **13**:2349-2360
- 43 DeVantier, L. M., G. De'ath, E. Turak, T. J. Done, K. E. Fabricius, *Coral Reefs* **25**, 329 (2006).
- 44 Devney, C. A., M. Short, B. C. Congdon, *Wildlife Research* **36**, 368 (2009).
- 45 Deysher, L. E., T. A. Dean, *Journal of Experimental Marine Biology and Ecology* **103**, 41 (Dec, 1986).
- 46 **Diaz** R. J. and Rosenberg R., 2008: Spreading dead zones and consequences for marine ecosystems. *Science*
47 **321**:926-929.
- 48 **Díaz-Almela** E, Marbà N and Duarte CM. 2007: Consequences of Mediterranean warming events in seagrass
49 (*Posidonia oceanica*) flowering records. *Global Change Biology* **13**: 224–235.
- 50 **Díaz-Almela**, E., N. Marbà, R. Martínez, R. Santiago and C. M. Duarte, 2009: Seasonal dynamics of *Posidonia*
51 *oceanica* in Magalluf Bay (Mallorca, Spain): temperature effects on seagrass mortality. *Limnology and*
52 *Oceanography* **54**: 2170–2182
- 53 **Diaz-Pulido** G., Gouezo M., Tilbrook B., Dove S. and Anthony K. R., 2011: *High CO2 enhances the competitive*
54 *strength of seaweeds over corals*. *Ecology Letters* **14(2)**:156-162.

- 1 **Dijkema, K.S.**, 1997: Impact prognosis for salt marshes from subsidence by gas extraction in the wadden
2 sea. *Journal of Coastal Research*, **13(4)**, 1294-1304.
- 3 *Dis. Aquat. Org.* **87**, 5-18 (2009).
- 4 **Dixon T.H.**, F. Amelung, A. Ferretti, F. Novali, F. Rocca, R. Dokka, G. Sella, S.-W. Kim, S. Wdowinski, and D.
5 Whitman, 2006: subsidence and flooding in New Orleans. *Nature* 441, 587–588.
- 6 **Dodd, C. K.**, M. J. Dreslik, *Journal of Zoology* **275**, 18 (2008).
- 7 **Dodet, G.**, X. Bertin, and R. Taborda, 2010: Wave climate variability in the North-East Atlantic Ocean over the last
8 six decades. *Ocean Modelling* **31**, 120-1313.
- 9 **Dolch, T.** and K. Reise, 2010: Long-term displacement of intertidal seagrass and mussel beds by expanding large
10 sandy bedforms in the northern wadden sea. *Journal of Sea Research*, **63(2)**, 93-101.
- 11 **Donato D. C.**, Kauffman J. B., Murdiyarto D., Kurnianto S., Stidham M. and Kanninen M., 2011: Mangroves
12 among the most carbon-rich forests in the tropics. *Nature Geoscience* **4**:293-297.
- 13 **Doney S. C.**, Mahowald N., Lima I., Feely R. A., Mackenzie F. T., Lamarque J.-F. and Rasch P. J., 2007: *Impact of*
14 *anthropogenic atmospheric nitrogen and sulfur deposition on ocean acidification and the inorganic carbon*
15 *system*. Proceedings of the National Academy of Science U.S.A. 104:14580-14585.
- 16 **Doody, J.**, 2004: 'Coastal squeeze' - an historical perspective. *Journal of Coastal Conservation*, **10(1)**, 129-138.
- 17 **Dornbusch, U.**, D.A. Robinson, C.A. Moses, and R.B.G. Williams, 2008: Temporal and spatial variations of chalk
18 cliff retreat in east sussex, 1873 to 2001. *Marine Geology*, **249(3-4)**, 271-282.
- 19 **Dragani, W.C.**, P.B. Martin, C. G. Simionato, and M. I. Campos, 2010: Are wind wave heights increasing in south-
20 eastern south American continental shelf between 32°S and 40°S? *Continental Shelf Research*, **30**, 481-490.
- 21 **Duarte C. M.**, Dennison W. C., Orth R. J. W. and Carruthers T. J. B., 2008: The charisma of coastal ecosystems:
22 addressing the imbalance. *Estuaries and Coasts* **31**:233-238.
- 23 **Duarte C. M.**, Middelburg J. J. and Caraco N., 2005: Major role of marine vegetation on the oceanic carbon cycle.
24 *Biogeosciences* **2**: 1-8.
- 25 **Duarte, C. M.**, N. Marbà, E. Gacia, J. W. Fourqurean, J. Beggins, C. Barrón, and E. T. Apostolaki 2010: Seagrass
26 community metabolism: assessing the carbon sink capacity of seagrass meadows, *Global Biogeochem. Cycles*,
27 **24**, GB4032.
- 28 **Duarte, C.M.** 2002: The future of seagrass meadows. *Environmental Conservation* **29**: 192-206
- 29 **Dullo W. C.**, 2005: Coral growth and reef growth: a brief review. *Facies* **51**:33-48.
- 30 **Dumaru, P.** 2010: Community-based adaptation: enhancing community adaptive capacity in Druadrua Island, Fiji.
31 Wiley Interdisciplinary Reviews: *Climate Change* **1** (Sept/Oct):751-763.
- 32 **Dunlop, J. N.**, C. A. Surman, R. D. Wooller, *Emu* **101**, 19 (2001).
- 33 **Dunlop, J. N.**, R. D. Wooller, *Records of the Western Australian Museum* **12**, 389 (1986).
- 34 **Dupuis J.**, and P. Knoepfel 2011: Les barrières à la mise en œuvre des politiques d'adaptation au changement
35 climatique: le cas de la Suisse. *Swiss Political Science Review* **17 (2)**:188–219.
- 36 **Dutta, D.**, W. Wright, and P. Rayment, 2011: Synthetic impact response functions for flood vulnerability analysis
37 and adaptation measures in coastal zones under changing climatic conditions: A case study in gippsland coastal
38 region, Australia. *Natural Hazards* **59(2)**, 967-986.
- 39 **Dwarakish, G.S.**, Vinay, S.A., Natesan, U., Toshiyuki, A., Kakinuma, T., Venkataramana, K., Pai, B.J. and Babita,
40 M.K. 2009: *Ocean and Coastal Management* **52**, 467–478.
- 41 **Ebi, K.L.**, J. Balbus, P.L. Kinney, E. Lipp, D. Mills, M.S. O'Neill, and M. Wilson, 2008: *Effects of Global Change*
42 *on Human Health. In: Analyses of the effects of global change on human health and welfare and human*
43 *systems*. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change
44 Research [Gamble, J.L. (ed.), K.L. Ebi, F.G. Sussman, T.J. Wilbanks, (Authors)]. U.S. Environmental
45 Protection Agency, Washington, DC, USA, p. 39–87
- 46 **Edmiston, H. L. et al.**, *J. Coast. Res.*, 38 (2008).
- 47 **Edwards, M.**, A. J. Richardson, *Nature* **430**, 881 (2004).
- 48 **Edwards, M.**, *Mar. Ecol.-Prog. Ser.*, 1 (2004).
- 49 **Ehlers, A.**, B. Worm, T. B. H. Reusch, *Mar. Ecol.-Prog. Ser.* **355**, 1 (2008).
- 50 **Ekstrom, J.A.**, Moser, S.C., and Torn, M., 2010: *Barriers to Adaptation: A Diagnostic Framework*. Final Project
51 Report. California Energy Commission, Sacramento, CA.
- 52 **Ellison, A. M.**, E. J. Farnsworth, *J. Ecol.* **84**, 717 (1996).
- 53 **Ellison, J. C.**, D. R. Stoddart, *J. Coast. Res.* **7**, 151 (1991).
- 54 **Ellison, J. C.**, *Estuarine Coastal and Shelf Science* **37**, 75 (1993).

- 1 **Elvidge, C.**, J. Dietz, R. Berkelmans, S. Andréfouët, W. Skirving, A. Strong and B. Tuttle 2004: Satellite
2 observation of Keppel Islands (Great Barrier Reef) 2002 coral bleaching using IKONOS data. *Coral Reefs*
3 **23**:123-132.
- 4 **Erdner, D.L.**, J. Dyble, M.J. Parsons, R.C. Stevens, K.A. Hubbard, M.L. Wrabel et al. 2008: Centers for oceans and
5 human health: a unified approach to the challenge of harmful algal blooms. *Environmental Health* **7**(Suppl 2),
6 S2.
- 7 **Ericson, J.P.**, C.J. Vorosmarty, S.L. Dingman, L.G. Ward and M. Meybeck, 2006: Effective sea-level rise and deltas:
8 causes of change and human dimension implications. *Global Planet Change*, **50**, 63-82.
- 9 **Espinosa-Romero, M. J.**, Chan, K. M. A., McDaniels, T. and Dalmer, D. M. 2011: Structuring decision-making for
10 ecosystem-based management. *Marine Policy* **35**(5): 575-583.
- 11 **Essink, G.H.P.**, E.S. van Baaren, and P.G.B. de Louw 2010: Effects of climate change on coastal groundwater
12 systems: A modeling study in the Netherlands. *Water Resources Research* **46**, W00F04.
- 13 **European Commission Climate Action:** http://ec.europa.eu/clima/sites/change/what_is_eu_doing/marine_en.htm
- 14 **Fabricius K. E.**, C. Langdon, S. Uthicke, C. Humphrey, S. Noonan, G. De'ath, R. Okazaki, N. Muehllehner, M.S.
15 Glas and J.M. Lough 2011: Losers and winners in coral reefs acclimatized to elevated carbon dioxide
16 concentrations. *Nature Climate Change* **1**: 165-169.
- 17 **Falaleeva M.**, C. O'Mahony, S. Gray, M. Desmond, J. Gault, and V. Cummins, 2011: Towards
18 climate adaptation and coastal governance in Ireland: Integrated architecture for effective management? *Marine*
19 *Policy* **35**: 784-993.
- 20 **Fankhauser, S.** 1995: Protection versus retreat: the economic costs of sea-level rise. *Environment and Planning*,
21 **27**(2), 299-319
- 22 **Farhan, A.R.** and S. Lim, 2011: *Resilience assessment on coastline changes and urban settlements: A case study in*
23 *seribu islands, Indonesia*. **54**(5), 391-400.
- 24 Feary, D. A., G. R. Almany, M. I. McCormick, G. P. Jones, *Oecologia* **153**, 727 (Sep, 2007).
- 25 **Feely R. A.**, Alin S. R., Newton J., Sabine C. L., Warner M., Devol A., Krembs C. and Maloy C., 2010: The
26 combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized
27 estuary. *Estuarine, Coastal and Shelf Science* **88**,442-449.
- 28 **Feely R. A.**, Sabine C. L., Hernandez-Ayon J. M., Ianson D. and Hales B., 2008: Evidence for upwelling of
29 corrosive "acidified" water onto the continental shelf. *Science* **320**:1490-1492.
- 30 **Feller I. C.**, Lovelock C. E., Berger U., McKee K. L., Joye S. B. and Ball M. C., 2010: Biocomplexity in mangrove
31 ecosystems. *Annual Reviews of Marine Science* **2**:395-417.
- 32 **Ferreira, O.**, Dias, J.A. and Taborda, R., 2008: Implications of Sea-Level Rise for Continental Portugal. *J. Coastal*
33 *Res.* **24**(2) 317-324.
- 34 **Few, R.**, K. Brown, and E.L. Tompkins, 2007: *Public participation and climate change adaptation: Avoiding the*
35 *illusion of inclusion*. **7**(1), 46-59.
- 36 **Findlay H. S.**, Burrows M. T., Kendall M. A., Spicer J. I. and Widdicombe S., 2010: Can ocean acidification affect
37 population dynamics of the barnacle *Semibalanus balanoides* at its southern range edge? *Ecology* **91**:2931-2940.
- 38 Findlay, H. S., M. A. Kendall, J. I. Spicer, S. Widdicombe, *Estuarine Coastal and Shelf Science* **86**, 675 (Mar,
39 2010).
- 40 Fish, M. R. *et al.*, *Conservation Biology* **19**, 482 (2005).
- 41 **FitzGerald, D.M.**, M.S. Fenster, B.A. Argow, I.V. Buynevich, 2008: Coastal impacts due to sea-level rise. *Annu.*
42 *Rev. Earth Planet. Sci.* **36**, 601-647.
- 43 **Fonseca, M.S.** and S.S. Bell, 1998: Influence of physical setting on seagrass landscapes near beaufort, north
44 carolina, USA. *Marine Ecology - Progress Series*, **171**, 109-121.
- 45 **Foufoula-Georgiou, E.**, J. Syvitski, C. Paola, C.T. Hoanh, P. Tuong, C. Vörösmarty, H. Kremer, Brondizio, Y.
46 Saito, R. Twilley, 2011: *International Year of Deltas 2013: A proposal*. *Eos, Transactions of American*
47 *Geophysical Union*, 92, 340-341.
- 48 Francour, P., C. F. Boudouresque, J. G. Harmelin, M. L. Harmelinviven, J. P. Quignard, *Marine Pollution Bulletin*
49 **28**, 523 (1994).
- 50 **Frazier, T. G.**, N. Wood, and B. Yarnal, 2010: Stakeholder perspectives on land-use strategies for adapting to
51 climate-change-enhanced coastal hazards: Sarasota, Florida. *Applied Geography* **30**(4):506-517.
- 52 Fu, F. X., *et al.*, *Harmful Algae* **7**, 76 (Jan, 2008).
- 53 Fu, F. X., M. E. Warner, Y. H. Zhang, Y. Y. Feng, D. A. Hutchins, *Journal of Phycology* **43**, 485 (Jun, 2007).

- 1 Fuentes, M. M. P. B., M. Hamann, C. J. Limpus, *Journal of Experimental Marine Biology and Ecology* **383**, 56
2 (2010).
- 3 **Fuentes**, M.M.P.B., C.J. Limpus, M. Hamann, and J. Dawson, 2010: Potential impacts of projected sea-level rise on
4 sea turtle rookeries. *Aquatic Conservation - Marine and Freshwater Ecosystems*, **20(2)**, 132-139.
- 5 **Fujii**, T. and D. Raffaelli, 2008: Sea-level rise, expected environmental changes and responses of intertidal benthic
6 macrofauna in the humber estuary, UK. *Marine Ecology - Progress Series*, **371**, 23-35.
- 7 Galbraith, H. *et al.*, *U S Forest Service General Technical Report PSW* **191**, 1119 (2005).
- 8 Gambaiani, D. D., P. Mayol, S. J. Isaac, M. P. Simmonds, *J. Mar. Biol. Assoc. U.K.* **89**, 179 (Feb, 2009).
- 9 **Gao**, K., Aruga, Y., Asada, K., Ishihara, T., Akano, T., Kiyohara, M. 1993: Calcification in the articulated coralline
10 alga *Corallina pilulifera* with special reference to the effect of elevated CO₂ concentration. *Marine Biology* **117**:
11 129-132.
- 12 **Garcia** S. M. and de Leiva Moreno I., 2003: Global overview of marine fisheries. In: Sinclair M. and Valdimarsson
13 G. (Eds.), *Responsible fisheries in the marine ecosystem*, pp. 103–123. Wallingford: CAB International.
- 14 Gardner, T. A., I. M. Cote, J. A. Gill, A. Grant, A. R. Watkinson, *Ecology* **86**, 174 (2005).
- 15 **Garmendia** E., G. Gamboa, J. Franco, J.M. Garmendia, P. Liria, and M. Olazabal 2010: Social multi-criteria
16 evaluation as a decision support tool for integrated coastal zone management. *Ocean and Coastal Management*,
17 **53**: 385–403.
- 18 Garza, C., C. Robles, *Mar. Biol.* **157**, 673 (2010).
- 19 **Gazeau** F., Quiblier C., Jansen J. M., Gattuso J.-P., Middelburg J. J. & Heip C. H. R., 2007: *Impact of elevated CO₂*
20 *on shellfish calcification*. *Geophysical Research Letters* **34**, L07603.
- 21 **Gedan** K. B., Kirwan M. L., Wolanski E., Barbier E. B. and Silliman B. R., 2011: The present and future role of
22 coastal wetland vegetation in protecting shorelines: answering recent challenges to the paradigm. *Climatic*
23 *Change* **106**:7-29.
- 24 **Gero**, A., Meheux, K. and Dominey-Howes, D. 2011: Integrating community based disaster risk reduction and
25 climate change adaptation: examples from the Pacific. *Nat. Hazards Earth Syst. Sci.* **11**: 101-113.
- 26 **Gesch**, Dean B. 2009: Analysis of Lidar Elevation for Improved Identification and Delineation of Lands Vulnerable
27 to Sea-Level Rise. *Journal of Coastal Research* **58**
- 28 **Gifford**, R. 2011: The dragons of inaction: Psychological barriers that limit climate change mitigation and
29 adaptation. *American Psychologist* **66 (4)**:290-302.
- 30 **Gill**, A. B. 2005: Offshore renewable energy: ecological implications of generating electricity in the coastal zone.
31 *Journal of Applied Ecology* **42(4)**:605-615.
- 32 **Gillanders**, B.M., T.S. Elsdon, I.A. Halliday, G.P. Jenkins, J.B. Robins, and F.J. Valesini, 2011: Potential effects of
33 climate change on Australian estuaries and fish utilising estuaries: A review. *Marine and Freshwater*
34 *Research*, **62(9)**, 1115-1131.
- 35 Gilman, E. L., *et al.*, *Climate Research* **32**, 161 (2006).
- 36 **Gilman**, E.L., J. Ellison, N.C. Duke, and C. Field, 2008: Threats to mangroves from climate change and adaptation
37 options: A review. *Aquatic Botany*, **89(2)**, 237-250.
- 38 **Giridharan**, R., S. S. Y. Lau, S. Ganesan, and B. Givoni, 2007: Urban design factors influencing heat island
39 intensity in high-rise high-density environments of Hong Kong. *Building and Environment* **42(10)**:3669-3684.
- 40 Godley, B. J., *et al.*, *Journal of Experimental Marine Biology and Ecology* **263**, 45 (2001).
- 41 Goffart, A., J. H. Hecq, L. Legendre, *Mar. Ecol.-Prog. Ser.* **236**, 45 (2002).
- 42 **Gómez-Gesteira** M., de Castro M., Alvarez I. and Gómez-Gesteira J. L., 2008: Coastal sea surface temperature
43 warming trend along the continental part of the Atlantic Arc (1985-2005). *Journal of Geophysical Research*
44 **113**, C04010.
- 45 **González** M., Medina, R. 2001. On the application of static equilibrium bay formulations to natural and man-made
46 beaches. *Coastal Engineering*, Vol. **43**, 209-225
- 47 Goreau, T. F., *Science* **145**, 383 (1964).
- 48 Gorgula, S. K., S. D. Connell, *Mar. Biol.* **145**, 613 (2004).
- 49 **Graham** N. A., Chabanet P., Evans R. D., Jennings S., Letourneur Y., Macneil M. A., McClanahan T. R., Ohman
50 M. C., Polunin N. V. and Wilson S. K., 2011: *Extinction vulnerability of coral reef fishes*. *Ecology Letters*
51 **14**:341-348.
- 52 Graham, M. H. *Journal of Experimental Marine Biology and Ecology* **218**, 127 (Nov, 1997).
- 53 Graham, M. H., *et al.*, *Mar. Ecol.-Prog. Ser.* **148**, 269 (Mar, 1997).

- 1 **Grall J.** and Chauvaud L., 2002: Marine eutrophication and benthos: the need for new approaches and concepts.
2 *Global Change Biology* **8**:813-830.
- 3 **Grantham B. A.,** Chan F., Nielsen K. J., Fox D. S., Barth J. A., Huyer A., Lubchenco J. and Menge B. A., 2004.
4 Upwelling-driven nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific.
5 *Nature* **429**:749-754.
- 6 Graus, R. R., I. G. Macintyre, *Carbonates Evaporites* **13**, 43 (1998).
- 7 Greve, T. M., J. Borum, O. Pedersen, *Limnology and Oceanography* **48**, 210 (2003).
- 8 **Grevelle, G.** and Mimura, N. (2008): Vulnerability assessment of sea-level rise in Viti Levu, Fiji Islands.
9 *Sustainability Science*, **3**: 171-180.
- 10 Griffiths, S. P., *Estuarine Coastal and Shelf Science* **58**, 173 (2003).
- 11 Guadayol, O., *et al.*, *Mar. Ecol.-Prog. Ser.* **381**, 139 (2009).
- 12 Guillemot, N., P. Chabanet, O. Le Pape, (2010) Cyclone effects on coral reef habitats in New Caledonia (South
13 Pacific) *Coral Reefs*, **29**, 445–453
- 14 **Gulev, S.K.** and V. Grigoviera, 2006: Variability of the winter wind waves and swell in the North Atlantic and
15 North Pacific as revealed by the voluntary observing ship data. *Journal of Climate*, **19**, 5667-5685.
- 16 **Gutiérrez D.,** Bouloubassi I., Sifeddine A., Purca S., Goubanova K., Graco M., Field D., Méjanelle L., Velazco F.,
17 Lorre A., Salvatelli R., Quispe D., Vargas G., Dewitte B. and Ortlieb L., 2011: *Coastal cooling and increased*
18 *productivity* in Geophysical Research Letters 38:L07603.
- 19 **Gutierrez, B.,** Plant, N., and Thieler, E.R., 2011: A Bayesian network to predict coastal vulnerability to sea level
20 rise. *J. Geophys. Research* Vol. **116**, F02009, 15 pp.
- 21 **Gutt, J.** 2001: On the direct impact of ice on marine benthic communities, a review. *Polar Biology* **24**: 553-564.
- 22 **Gypens N.,** Borges A. V. & Lancelot C., 2009: Effect of eutrophication on air-sea CO₂ fluxes in the coastal
23 Southern North Sea: a model study of the past 50 years. *Global Change Biology* **15**:1040-1056.
- 24 **Gypens N.,** Lacroix G., Lancelot C. and Borges A. V., 2011: *Seasonal and inter-annual variability of air-sea CO₂*
25 *fluxes and seawater carbonate chemistry in the Southern North Sea.* *Progress in Oceanography* 88:59-77.
- 26 **Hadley, D.** 2009: Land use and the coastal zone. *Land Use Policy*, **265**: S198-S203.
- 27 Hale, J. A., T. K. Frazer, D. A. Tomasko, M. O. Hall, *Estuaries* **27**, 36 (2004).
- 28 **Hallegatte, S.** 2009: Strategies to adapt to an uncertain climate change. *Global Environmental Change* **19(2)**: 240-
29 247.
- 30 **Hallegatte, S.,** Ranger, N., Mestre, O., Dumas, P., Corfee-Morlot, J., Herweijer, C., Muir-Wood, R., 2011:
31 Assessing climate change impacts, sea level rise and storm surge risk in port cities: a case study on Copenhagen
32 . *Climatic Change* **104**:113–137
- 33 **Hall-Spencer J. M.,** Rodolfo-Metalpa R., Martin S., Ransome E., Fine M., Turner S. M., Rowley S. J., Tedesco D.
34 and Buia M.-C., 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*
35 **454**:96-99.
- 36 **Halpern B. S.,** Walbridge S., Selkoe K. A., Kappel C. V., Micheli F., D'Agrosa C., Bruno J. F., Casey K. S., Ebert
37 C., Fox H. E., Fujita R., Heinemann D., Lenihan H. S., Madin E. M. P., Perry M. T., Selig E. R., Spalding M.,
38 Steneck R. and Watson R., 2008a: A global map of human impact on marine ecosystems. *Science* **319**:948-952.
- 39 **Halpern, B. S.,** K. L. McLeod, A. A. Rosenberg, and L. B. Crowder, 2008b: Managing for cumulative impacts in
40 ecosystem-based management through ocean zoning. *Ocean and Coastal Management* **51(3)**:203-211.
- 41 **Handmer, J.,** Y. Honda, Z.W. Kundzewicz, N. Arnell, G. Benito, J. Hatfield, I.F. Mohamed, P. Peduzzi, S. Wu, B.
42 Sherstyukov, K. Takahashi, and Z. Yan, 2012: Changes in impacts of climate extremes: human systems and
43 ecosystems. In: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*
44 [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K.
45 Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. A Special Report of Working Groups I and II of the
46 Intergovernmental Panel on Climate Change (IPCC). Cambridge University Press, Cambridge, UK, and New
47 York, NY, USA, pp. 231-290.
- 48 **Hansen, H. S.** 2011: *Urban Land-Use Projections Supporting Adaptation Strategies to Climate Changes in the*
49 *Coastal Zone.* *Geocomputation, Sustainability and Environmental Planning: Studies in Computational*
50 *Intelligence* 348: 17-34.
- 51 **Hansen, L.,** J. Hoffman, C. Drews and E. Mielbrecht 2010: Designing Climate-Smart Conservation: Guidance and
52 Case Studies. *Conservation Biology* **24(1)**, 63-69.
- 53 **Hanson, S.,** Nicholls, R. Ranger, N., Hallegatte, S., Dorfee-Morlot, J., Herweijer, C. and Chateau, J., 2011: A global
54 ranking of port cities with high exposure to climate extremes. *Climatic Change* **104(1)**, 89-111.

- 1 **Hansson D.**, Eriksson C., Omstedt A. and Chen D. L., 2011: Reconstruction of river runoff to the Baltic Sea, AD
2 1500-1995. *International Journal of Climatology* **31**:696-703.
- 3 **Hapke, C.**, Himmelstoss, E., Kratzmann, M., List, J., and Thieler, E.R., 2011: *National Assessment of Shoreline*
4 *Change; historical shoreline change along the New England and Mid-Atlantic coasts*, Open-File Report, U.S.
5 Geological Survey, No. 2010-1118, 57 pp.
- 6 Harley, C. D. G. *et al.*, *Ecology Letters* **9**, 228 (2006).
- 7 Harley, M.D., Turner, I.L., Short, A.D., Ranasinghe, R. 2010. Interannual variability
8 Harris, G. P., F. B. Griffiths, L. A. Clementson, H. Lyne, H. van der Voe, *Journal of Plankton Research* **13**, 109
9 (1991).
- 10 Harty, C., *Coastal Management* **32**, 405 (2004).
- 11 **Hashizume, M.**, Armstrong, B., Hajat, S., Wagatsuma, Y., Faruque, A.S.G., Hayashi, T., and Sack, D.A. 2007:
12 Association between climate variability and hospital visits for non-cholera diarrhoea in Bangladesh: effects and
13 vulnerable groups. *International Journal of Epidemiology* **36**:1030–1037
- 14 **Hashizume, M.**, B. Armstrong, Y. Wagatsuma, A.S.G. Faruque, T. Hayashi, and D.A. Sack 2008: Rotavirus
15 infections and climate variability in Dhaka, Bangladesh: a time-series analysis. *Epidemiology and Infections*,
16 **136**, 1281-1289.
- 17 **Hawkins S. J.**, Moore P. J., Burrows M. T., Poloczanska E., Mieszkowska N., Herbert R. J. H., Jenkins S. R.,
18 Thompson R. C., Genner M. J. and Southward A. J., 2008. Complex interactions in a rapidly changing world:
19 responses of rocky shore communities to recent climate change. *Climate Research* **37**:123-133.
- 20 Hawkins, S. J., *et al.*, *Marine Ecology, Progress Series* **396**, 245 (2009).
- 21 **Heip, C.H.R.**, Goosen N.K., Herman, P.M.J. Kromkamp J., Middelburg, J.J. and Soetaert, K. (1995) Production and
22 consumption of biological particles in temperate tidal estuaries. *Oceanogr.Mar. Biol. Ann. Reviews* **33**, 1-150.
- 23 **Helmle K. P.**, Dodge R. E., Swart P. K., Gledhill D. K. and Eakin C. M., 2011. Growth rates of Florida corals from
24 1937 to 1996 and their response to climate change. *Nature Communications* **2**:215.
- 25 **Helmuth B.**, Harley C. D. G., Halpin P. M., Oandapos,Donnell M., Hofmann G. E. and Blanchette C. A., 2002:
26 Climate change and latitudinal patterns of intertidal thermal stress. *Science* **298**:1015-1017.
- 27 **Helmuth B.**, Mieszkowska N., Moore P. and Hawkins S. J., 2006: Living on the edge of two changing worlds:
28 forecasting the responses of rocky intertidal ecosystems to climate change. *Annual Review of Ecology Evolution*
29 *and Systematics* **37**, 373.
- 30 **Hemer, M.A.**, 2010: Historical trends in Southern Ocean storminess: Long-term variability of extreme wave heights
31 at Cape Sorell, Tasmania. *Geophysical Research Letters* **37**
- 32 **Hemer, M.A.**, J.A. Church, and J.R. Hunter, 2010: Variability and trends in the directional wave climate of the
33 Southern Hemisphere. *International Journal of Climatology* **30**, 475-491.
- 34 **Hemminga, M.A.**, and C.M. Duarte 2000: *Seagrass Ecology*. Cambridge Univ. Press, Cambridge.
- 35 **Hendriks, I. E.**, C.M. Duarte and M. Álvarez, 2010: Vulnerability of marine biodiversity to ocean acidification: a
36 meta-analysis. *Estuarine, Coastal and Shelf Estuarine Science* **86**: 157–164.
- 37 **Hiddink J. G.** and Hofstede R., 2008: Climate induced increases in species richness of marine fishes. *Global*
38 *Change Biology* **14**:453-460.
- 39 **Hinkel, J.** and R.J.T. Klein, 2009: The DINAS-COAST project: Developing a tool for the dynamic and interactive
40 assessment of coastal vulnerability. *Global Environmental Change*, **19(3)**, 384-395.
- 41 **Hinkel, J.** Nicholls, R.J., Vafeidis, A.T., Tol, S.J. and Avagianou, T. 2010: Assessing risk of and adaptation to sea-
42 level rise in the European Union: an application of DIVA. *Mitigation and Adaptation Strategies for Global*
43 *Change*, **5(7)**, 1-17.
- 44 **Hinkel, J.**, 2011: Climate impacts: From numbers to stories. In: *Reframing the problem of climate change: From*
45 *zero sum game to win-win solutions*. [Carlo C. Jaeger, Klaus Hasselmann, Gerd Leipold, Diana Mangalagu,
46 and J. David Tabara (eds.)]. Earthscan, Oxon, UK, New York, USA & Canada, pp. 35-53.
- 47 **Hinkel, J.**, Brown, S., Exner, L., Nicholls, R.J., Vafeidis, A.T. and Kebede, A.S. 2011: Sea-level rise impacts on
48 Africa and the effects of mitigation and adaptation: an application of DIVA. *Regional Environmental*
49 *Change*, **12**, 207-224.
- 50 **Hinkel, J.**, D.P.v. Vuuren, R.J. Nicholls, and R.J.T. Klein, 2012: *The effects of mitigation and adaptation on coastal*
51 *impacts in the 21st century*.
- 52 **Hinz H.**, Capasso E., Lilley M., Frost M. and Jenkins S. R., 2011: Temporal differences across a bio-geographical
53 boundary reveal slow response of sub-littoral benthos to climate change. *Marine Ecology Progress Series*
54 **423**:69-82.

- 1 **Hochachka** P. W. and Somero G. N., 2002: *Biochemical adaptation: mechanism and process in physiological*
2 *evolution*, xi, 466 p. New York: Oxford University Press.
- 3 **Hoegh-Guldberg** O. and Bruno J. F., 2010: The impact of climate change on the world's marine ecosystems.
4 *Science* **328**:1523-1528.
- 5 **Hoegh-Guldberg** O., 1999: Climate change, coral bleaching, and the future of the world's coral reefs. *Marine and*
6 *Freshwater Research* **50**:839-866.
- 7 **Hoegh-Guldberg** O., 2011: Coral reef ecosystems and anthropogenic climate change. *Regional Environmental*
8 *Change* **11**:215-227.
- 9 **Hoegh-Guldberg** O., Mumby P. J., Hooten A. J., Steneck R. S., Greenfield P., Gomez E., Harvell C. D., Sale P. F.,
10 Edwards A. J., Caldeira K., Knowlton N., Eakin C. M., Iglesias-Prieto R., Muthiga N., Bradbury R. H., Dubi A.
11 and Hatzilios M. E., 2007: Coral reefs under rapid climate change and ocean acidification. *Science* **318**:1737-
12 1742.
- 13 **Hoeke** R. K., Jokiel P. L., Buddemeier R. W. and Brainard R. E., 2011: *Projected changes to growth and mortality*
14 *of Hawaiian corals over the next 100 years*. PLoS One 6:e18038.
- 15 **Hofmann** A, Soetaert K and Middelburg JJ, 2008: Present nitrogen and carbon dynamics in the Scheldt estuary
16 using a novel 1-D model. *Biogeosciences* **5**, 981- 1006.
- 17 **Hofmann** A. F., Peltzer E. T., Walz P. M. and Brewer P. G., 2011: *Hypoxia by degrees: establishing definitions for*
18 *a changing ocean*. Deep-Sea Research (Part I, Oceanographic Research Papers) 58: 1212-1226.
- 19 **Hofmann** G. E., Smith J. E., Johnson K. S., Send U., Levin L. A., Micheli F., Paytan A., Price N. N., Peterson B.,
20 Takeshita Y., Matson P. G., Crook E. D., Kroeker K. J., Gambi M. C., Rivest E. B., Frieder C. A., Yu P. C. and
21 Martz T. R., 2011: *High-frequency dynamics of ocean pH: a multi-ecosystem comparison*. PLoS ONE
22 6:e28983.
- 23 **Holmer**, M., P. Wirachwong, and M.S. Thomsen, 2011: Negative effects of stress-resistant drift algae and high
24 temperature on a small ephemeral seagrass species. *Marine Biology*, **158(2)**, 297-309.
- 25 Hoover, R. S., *et al.*, *Mar. Ecol.-Prog. Ser.* **318**, 187 (2006).
- 26 **Horstman**, E. M., K. M. Wijnberg, A. J. Smale and Sjmh Hulscher 2009. Long-term Coastal Management
27 Strategies: Useful or Useless? *Journal of Coastal Research*: 233-237.
- 28 Horton and Rosenzweig. 2010
- 29 **Howarth** R., Chan F., Conley D. J., Garnier J., Doney S. C., Marino R. and Billen G., 2011. Coupled
30 biogeochemical cycles: eutrophication and hypoxia in temperate estuaries and coastal marine ecosystems.
31 *Frontiers in Ecology and the Environment* **9**:18-26.
- 32 Hsieh, C. H., H. J. Kim, W. Watson, E. Di Lorenzo, G. Sugihara, *Global Change Biology* **15**, 2137 (Sep, 2009).
- 33 **Hsu**, K-C., C-H. Wang, K-C. Chen, C-T. Chen, and K-W. Ma 2007: Climate-induced hydrological impacts on the
34 groundwater system of the Pingtung Plain, Taiwan. *Hydrogeology Jour.*, **15**, 903-913.
- 35 **Huang**, C., P. Vaneckova, X. Wang, G. FitzGerald, Y. Guo and S. Tong S. 2011. Constraints and barriers to public
36 health adaptation to climate change: A review of the literature. *American Journal of Preventive Medicine*, **40**:
37 183-190.
- 38 Huang, T., L. G. Sun, Y. H. Wang, X. D. Liu, R. B. Zhu, *Antarctic Science* **21**, 571 (2009).
- 39 Hughes, T. P., J. H. Connell, *Limnology and Oceanography* **44**, 932 (1999).
- 40 **Hunt**, A. and P. Watkiss 2011: Climate change impacts and adaptation in cities: a review of the literature. *Climatic*
41 *Change*, **104**, 13-49.
- 42 **Hunter**, J., 2010: Estimating sea level extremes under conditions of uncertain sea level rise. *Climatic Change* **99**,
43 331-350.
- 44 **Huntington** T. G., 2006: Evidence for intensification of the global water cycle: review and synthesis. *Journal of*
45 *Hydrology* **319**:83-95.
- 46 **Huq** S. and H. Reid. 2007: *Community-Based Adaptation: A vital approach to the threat climate change poses to*
47 *the poor*. IIED Briefing Paper. London: International Institute for Environment and Development.
- 48 **Ibrahim** H.S. and D. Shaw 2012: Assessing progress toward integrated coastal zone management: Some lessons
49 from Egypt. *Ocean and Coastal Management*, **58**: 26-35.
- 50 **IPCC, 2012**: Summary for policymakers. In: *Managing the Risks of Extreme Events and Disasters to Advance*
51 *Climate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D.
52 Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. A Special Report of
53 Working Groups I and II of the Intergovernmental Panel on Climate Change. Cambridge University Press,
54 Cambridge, UK, and New York, NY, USA, pp. 1-19.

- 1 **Irving** A. D., Connell S. D. and Russell B. D., 2011: Restoring coastal plants to improve global carbon storage:
2 reaping what we sow. *PLoS ONE* 6: e18311.
- 3 **Isager**, L. 2008: Coastal Zone Management in Developing Countries with Kenya as a Particular Example.
4 Geocenter, Danmark. http://geocenter.dk/projekter/2008_coastal_zone/index.html
- 5 **ISDR**, 2009: Global Assessment Report on Disaster Risk Reduction, International Strategy for Disaster Reduction,
6 United Nations, Geneva, Switzerland, ISBN/ISSN: 9789211320282, 207 pp.
- 7 **IUCN** 2008: Ecosystem-based adaptation: An approach for building resilience and reducing risk for local
8 communities and ecosystems. A submission by IUCN to the Chair of the AWG-LCA with respect to the Shared
9 Vision and Enhanced Action on Adaptation, on behalf of: IUCN, The Nature Conservancy, WWF,
10 Conservation International, BirdLife International, Indigenous Peoples of Africa Co-ordinating Committee,
11 Practical Action, WILD Foundation, Wildlife Conservation Society, Fauna and Flora International and
12 Wetlands International. Gland, Switzerland: IUNC.
- 13 **Izaguirre**, C., F. J. Mendez, M. Menendez, and I. J. Losada. 2011. Global extreme wave height variability based
14 on satellite data, *Geophys. Res. Lett.*, 38, L10607..
- 15 **Jackson**, A.C. and J. McIlvenny, 2011: Coastal squeeze on rocky shores in northern scotland and some possible
16 ecological impacts. *Journal of Experimental Marine Biology and Ecology*, **400(1-2)**, 314-321.
- 17 **Jackson**, M.C., L. Johansen, C. Furlong, A. Colson, and K.F. Sellers 2010: Modelling the effects of climate change
18 on prevalence of malaria in western Africa. *Statistica Neerlandica*, 64, 388-400.
- 19 **Jacob KH**, Gornitz V, Rosenzweig C (2007) Vulnerability of the New York City metropolitan area to coastal
20 hazards, including sea-level rise: inferences for urban coastal risk management and adaptation policies. In:
21 McFadden L, Nicholls RJ, Penning-Roswell E (eds) *Managing coastal vulnerability*, Amsterdam, Elsevier, pp
22 61–88
23 Jax, 2006
- 24 **Jaykus**, L.-A., Woolridge, M., Frank, J.M., Miraglia, M., McQuatters-Gollop, A., Tirado, C. Clarke, R. and Friel,
25 M. 2008: Food and Agriculture Organization of the United Nations (FAO) paper 49pp.
- 26 **Jentoft**, S. 2009. "Future Challenges in Environmental Policy Relative to Integrated Coastal Zone Management."
27 *Integrated Coastal Zone Management* pp. 155-169: Wiley-Blackwell.
- 28 **Jeppesen** E., Sondergaard M., Pedersen A. R., Jurgens K., Strzelczak A., Lauridsen T. L. and Johansson L. S., 2007.
29 Salinity induced regime shift in shallow brackish lagoons. *Ecosystems* 10:47-57.
- 30 **Jha**, A., J. Lamond, R. Bloch, N. Bhattacharya, A. Lopez, N. Papachristodoulou, A. Bird, D. Proverbs, J. Davies,
31 and R. Barker 2011: Five Feet High and Rising – Cities and Flooding in the 21st Century. Policy Research
32 Working Paper 5648, The World Bank, 62 pp.
- 33 **Jha**, A.K. and H. Brecht 2011: Building Urban Resilience in East Asia. An Eye on East Asia and the Pacific. The
34 World Bank, 13 pp. (not peer reviewed).
- 35 **Johnson**, C., S. Ling, J. Ross, S. Shepherd, K. Miller, 2005: Establishment of the long-spined sea urchin
36 (*Centrostephanus rodgersii*) in Tasmania: first assessment of potential threats to fisheries (Tasmanian
37 Aquaculture and Fisheries Institute, Australia.
- 38 Jokieli, P. L., C. L. Hunter, S. Taguchi, L. Watarai, *Coral Reefs* **12**, 177 (1993).
- 39 **Jude**, S., 2008: Investigating the potential role of visualization techniques in participatory coastal
40 management. **36(4)**, 331-349.
- 41 **Kabat**, P., L.O. Fresco, M.J.C. Stive, M.J.C., C.P. Veerman, J.S.L.J. van Alphen, B.W.A.H. Parmet, W. Hazeleger,
42 and C.A. Katsman, C.A. 2009. Dutch coasts in transition. *Nature Geoscience*, Vol. 2, July, pp. 450-452.
- 43 **Kapos**, V., J. Scharlemann, P. Bubb, A. Campbell, A. Chenery, N. Doswald and B. Dickson 2009. Impacts of
44 climate change on biodiversity – A review of the recent scientific literature. In : Review of the Literature on the
45 Links between Biodiversity and Climate Change: Impacts, Adaptation and Mitigation [Campbell, A., V. Kapos,
46 J.P.W. Scharlemann, P. Bubb, A. Chenery, L. Coad, B. Dickson, N. Doswald, M.S.I. Khan, F. Kershaw and M.
47 Rashid, eds.], Secretariat of the Convention on Biological Diversity, Montreal. Technical Series No. 42, p. 7-47.
- 48 **Karaca**, M. and Nicholls, R. J. 2008: Potential Implications of Accelerated Sea-Level Rise for Turkey. *Journal of*
49 *Coastal Research*, 24(2), 288-298.
- 50 **Kebede**, A.S. and Nicholls R.J. 2012: Exposure and vulnerability to climate extremes: population and asset
51 exposure to coastal flooding in Dar es Salaam, Tanzania. *Regional Environmental Change* 12:81–94
- 52 **Keener** V.W., Izuka, S.K., Anthony, S. (2012). Freshwater and Drought on Pacific Islands. In: Climate Change and
53 Pacific Islands: Indicators and Impacts. Report for The 2012 Pacific Islands Regional Climate Assessment.
54 Honolulu, Hawai'i, USA.

- 1 **Kelly, R. P., M. M. Foley, W. S. Fisher, R. A. Feely, B. S. Halpern, G. G. Waldbusser and M. R. Caldwell** 2011.
2 Mitigating local causes of ocean acidification with existing laws. *Science* 332: 1036-1037.
- 3 **Kendall, M. A., M. T. Burrows, A. J. Southward, S. J. Hawkins, *Ibis* 146, 40 (2004).**
- 4 **Kennedy H, Beggins J, Duarte CM, Fourqurean JW, Holmer M, Marbà N, and Middelburg JJ.** 2010. Seagrass
5 sediments as a global carbon sink: Isotopic constraints. *Global Biogeochemical Cycles* 24.
- 6 **Kennedy H. and Björk M.,** 2009: Seagrass meadows. In: Laffoley D. & Grimsditch G. (Eds.). *The management of*
7 *natural coastal carbon sinks*, pp. 23-29. Gland, Switzerland: IUCN.
- 8 **Kennedy, V. S. *Mar. Biol.* 35, 127 (1976).**
- 9 **Kettle N.P.** 2012. Exposing compounding uncertainties in sea level rise assessments. *Journal of Coastal Research*,
10 28: 161-173.
- 11 **Kiessling W. and Simpson C.,** 2011. On the potential for ocean acidification to be a general cause of ancient reef
12 crises. *Global Change Biology* 17:56-67.
- 13 **King, B. R., J. T. Hicks, J. Cornelius, *Emu* 92, 1 (1992).**
- 14 **Kirshen P., S. Merrill, P. Slovinsky and N. Richardson.** 2011. Simplified method for scenario-based risk assessment
15 adaptation planning in the coastal zone. *Climatic Change*, pp: 1-13. Doi: 10.1007/s10584-011-0379-z.
- 16 **Kirwan M. L. and G. R. Guntenspergen,** 2010: The influence of tidal range on the stability of coastal marshland, *J.*
17 *Geophys. Res.*, 115, F02009, doi:10.1029/2009JF001400.
- 18 **Kirwan M.L. and L.K. Blum (2011):** Enhanced decomposition offsets enhanced productivity and soil carbon
19 accumulation in coastal wetlands responding to climate change. *Biogeosciences*, 8, 987–993, 2011
- 20 **Kirwan M.L. and Temmerman S. (2009):** Coastal marsh response to historical and future sea-level acceleration.
21 *Quaternary Science Reviews* 28(17-18), 1801–1808.
- 22 **Kirwan M. L., G. R. Guntenspergen, A. D’Alpaos, J. T. Morris, S. M. Mudd, and S. Temmerman.** 2010: Limits on
23 the adaptability of coastal marshes to rising sea level. *Geophysical Research Letters*. 37(23):L23401.
- 24 **Kittinger J.N. and A.L. Ayers** 2010: Shoreline armoring, risk management, and coastal resilience under rising seas.
25 *Coastal Management*, 38, 634-653.
- 26 **Klein R.J.T., M. Alam, I. Burton, W.W. Dougherty, K. L. Ebi, M. Fernandes, A. Huber-Lee, A.T. Rahman, and C.**
27 **Swartz** 2006: Application of Environmentally Sound Technologies for Adaptation to Climate Change.
28 Technical Paper UNFCCC FCCC/TP/2006/2, 107 pp.
- 29 **Kleypas J. A., Danabasoglu G. and Lough J. M.,** 2008. Potential role of the ocean thermostat in determining
30 regional differences in coral reef bleaching events. *Geophysical Research Letters* 35, ARTN L03613.
31 doi:10.1029/2007GL032257.
- 32 **Knowlton, N., *Proceedings of the National Academy of Sciences of the United States of America* 98, 5419 (2001).**
- 33 **Knowlton, N., J. C. Lang, M. C. Rooney, P. Clifford, *Nature* 294, 251 (1981).**
- 34 **Knutson, T. R., J. L. McBride, J. Chan, K. Emanuel, G. Holland C. Landsea, I. Held, J. P. Kossin, A. K.**
35 **Srivastava, and M. Sugi,** 2010: Tropical Cyclones and Climate Change. *Nature Geoscience*, Review Article, 21
36 February 2010, p. 157-163. DOI: 10.1038/NCEO779.
- 37 **Kolivras K.N.** 2010: Changes in dengue risk potential in Hawaii, USA, due to climate variability and change.
38 *Climate Research*, 42, 1-11.
- 39 **Kolstad, E.W. and K.A. Johansson** 2011: Uncertainties associated with quantifying climate change impacts on
40 human health : a case study for diarrhea. *Environmental Health Perspectives*, 119, 299-305.
- 41 **Komar, P.D., and J.C. Allan,** 2008: Increasing hurricane-generated wave heights along the US East Cost and their
42 climate controls. *Journal of Coastal Research*, 24, 479-488.
- 43 **Kont, A., J. Jaagus, R. Aunap, U. Ratas, and R. Rivis,** 2008: Implications of sea-level rise for Estonia. *Journal of*
44 *Coastal Research*, 24(2), 423-431
- 45 **Krauss, K., D. Cahoon, J. Allen, K. Ewel, J. Lynch, and N. Cormier,** 2010: Surface elevation change and
46 susceptibility of different mangrove zones to sea-level rise on pacific high islands of
47 micronesia. *Ecosystems*, 13(1), 129-143.
- 48 **Kristensen, E., S. Bouillon, T. Dittmar, and C. Marchand.** 2008. Organic carbon dynamics in mangrove ecosystems:
49 A review. *Aquatic Botany* 89 (2):201-219.
- 50 **Kuffner I.B., A.J. Andersson, P.L. Jokiel, K.S. Rodgers, F.T. Mackenzie.** 2008. Decreased abundance of crustose
51 coralline algae due to ocean acidification. *Nature Geoscience* 1, 114 – 117.
- 52 **Langley, J.A., K.L. McKee, D.R. Cahoon, J.A. Cherry, and J.P. Megonigal,** 2009: Elevated CO2 stimulates marsh
53 elevation gain, counterbalancing sea-level rise. *Proceedings of the National Academy of Sciences of the United*
54 *States of America*, 106(15), 6182-6186.

- 1 **Lantuit**, H., Atkinson, D., Overduin, P.P., Grigoriev, M., Rachold, V., Grosse, G., Hubberten, H.W., 2011, Coastal
2 erosion dynamics on the permafrost-dominated Bykovsky Peninsula, north Siberia, 1951-2006, *Polar Research*,
3 30, 7341, DOI: 10.3402/polar.v30i0.7341
- 4 **Larsen**, P.H., S. Goldsmith, O. Smith, M.L. Wilson, K. Strzepek, P. Chinowsky, and B. Saylor 2008: Estimating
5 future costs for Alaska public infrastructure at risk from climate change. *Global Environment Change*, 18, 442-
6 457.
- 7 **Larson**, S. 2010 Understanding barriers to social adaptation: are we targeting the right concerns? *Architectural*
8 *Science Review* 53 (1):51-58.
- 9 **Laruelle** GG, Durr HH, Slomp CP, Borges AV. 2010. Evaluation of sinks and sources of CO₂ in the global coastal
10 ocean using a spatially explicit typology of estuaries and continental shelves. *Geophys. Res. Lett.* 37:L15607
- 11 **Last**, P.R., White, W.T., Gledhill, D. C., Hobday, A.J., Brown, R., Edgar, G.J. and Pecl, G. 2011: Long-term shifts
12 in abundance and distribution of a temperate fish fauna: a response to climate change and fishing practices.
13 *Global Ecol. Biogeogr.* 20, 58–72.
- 14 **Lata** S., and P. Nunn. 2011. Misperceptions of climate-change risk as barriers to climate-change adaptation: a case
15 study from the Rewa Delta, Fiji. *Climatic Change* online first. DOI 10.1007/s10584-011-0062-4
- 16 **Ledoux** L., S. Cornell, T. O'Riordan, R. Harvey, and L. Banyard. 2005. Towards sustainable flood and coastal
17 management: identifying drivers of, and obstacles to, managed realignment. *Land Use Policy* 22 (2):129-144.
- 18 **Lerman** A., Guidry M., Andersson A. J. and Mackenzie F. T., 2011. Coastal ocean last glacial maximum to 2100
19 CO₂-carbonic acid-carbonate system: a modeling approach. *Aquatic Geochemistry* 17:749-773.
- 20 **Levermann**, A., J.L. Bamber, S. Drijfhout, A. Ganopolski, W. Haeberli, Harris, Neil R. P. and Huss, Matthias, K.
21 Krueger, T.M. Lenton, R.W. Lindsay, D. Notz, P. Wadhams, and S. Weber, 2012}: Potential climatic
22 transitions with profound impact on europe review of the current state of six `tipping elements of the climate
23 system'. *Climatic Change*, **110(3-4)**, 845-878.
- 24 **Levinton**, J., M. Doall, D. Ralston, A. Starke, and B. Allam, 2011: Climate change, precipitation and impacts on an
25 estuarine refuge from disease. *Plos One*, **6(4)**.
26 Lewis et al., 2012
- 27 **Li**, J., M.H. Wang and Y.S. Ho 2011. Trends in research on global climate change: A Science Citation Index
28 Expanded-based analysis. *Global and Planetary Change*, 77: 13-20.
- 29 **Li**, K. and G.S. Li, 2011: Vulnerability assessment of storm surges in the coastal area of guangdong
30 province. *Natural Hazards and Earth System Sciences*, **11(7)**, 2003-2010.
- 31 **Lima** F. P. and Wethey D. S., 2012. Three decades of high-resolution coastal sea surface temperatures reveal more
32 than warming. *Nature Communications* 3:1-13.
- 33 **Lima** F. P., Ribeiro P. A., Queiroz, Hawkins S. J. and Santos A. M., 2007. Do distributional shifts of northern and
34 southern species of algae match the warming pattern? *Global Change Biology* 13:2592-2604.
- 35 Limpus, C. J., P. C. Reed, *Australian Wildlife Research* **12**, 523 (1985).
- 36 Limpus, C. J., F. Heidrun, 2006: *Migratory species and climate change: impacts of a changing environment on wild*
37 *animals.*, 34.
- 38 **Ling** S. D., Johnson C. R., Frusher S. D. and Ridgway K. R., 2009. Overfishing reduces resilience of kelp beds to
39 climate-driven catastrophic phase shift. *Proceedings of the National Academy of Science U.S.A.* 106:22341-
40 22345.
- 41 **Linham**, M.M. and R.J. Nicholls, 2010: *Technologies for climate change adaptation - coastal erosion and*
42 *flooding*. UNEP Riso Centre on Energy and Climate and Sustainable Development, Roskilde, Denmark.
- 43 **Lloret** J., Marín A. and Marín-Guirao L., 2008. Is coastal lagoon eutrophication likely to be aggravated by global
44 climate change? *Estuarine, Coastal and Shelf Science* 78:403-412.
- 45 **Lofman** D., M. Petersen, and A. Bower. 2002. Water, energy and environment nexus: The California experience.
46 *International Journal of Water Resources Development* 18(1): p. 73-85.
- 47 **Lonsdale**, K. G., M. J. Gawith, K. Johnstone, R. B. Street, C. C. West, and A. D. Brown. 2010. Attributes of Well-
48 Adapting Organisations. A report prepared by UK Climate Impacts Programme for the Adaptation Sub-
49 Committee: UK Climate Impacts Programme.
- 50 **Lotze** H. K., Lenihan H. S., Bourque B. J., Bradbury R. H., Cooke R. G., Kay M. C., Kidwell S. M., Kirby M. X.,
51 Peterson C. H. and Jackson J. B. C., 2006. Depletion, degradation, and recovery potential of estuaries and
52 coastal seas. *Science* 312:1806-1809.
- 53 **Lough** J. M., 2000. 1997-98: Unprecedented thermal stress to coral reefs? *Geophys. Res. Lett.* 27(23): 3901-3904.

- 1 **Lovelock**, C.E., I.C. Feller, M.F. Adame, R. Reef, H.M. Penrose, L. Wei, and M.C. Ball, 2011: Intense storms and
2 the delivery of materials that relieve nutrient limitations in mangroves of an arid zone estuary. *Functional Plant*
3 *Biology*, **38(6)**, 514-522.
- 4 **Lowe**, J.A. and J.M. Gregory 2010. A sea of uncertainty. *Nature Reports Climate Change*, **4** : 42-43.
- 5 Madsen, A. T., A. S. Murray, T. J. Andersen, M. Pejrup, *Mar. Geol.* **242**, 221 (Aug, 2007).
- 6 **Mangal**, T.D., Paterson, S., and Fenton, A. 2008: Predicting the Impact of Long-Term Temperature Changes on the
7 Epidemiology and Control of Schistosomiasis: A Mechanistic Model. PLoS one, 1438(1) 1-9.
- 8 **Manson**, G.K. and S.M. Solomon (2007). Past and future forcing of Beaufort Sea coastal change. *Atmosphere-*
9 *Ocean*, **45(2)**, 107-122.
- 10 **Manzello** D. P., 2010. Coral growth with thermal stress and ocean acidification: lessons from the eastern tropical
11 Pacific. *Coral Reefs* 29:749-758.
- 12 **Manzello** D. P., Kleypas J. A., Budd D. A., Eakin C. M., Glynn P. W. and Langdon C., 2008. Poorly cemented coral
13 reefs of the eastern tropical Pacific: Possible insights into reef development in a high-CO2 world. Proceedings
14 of the National Academy of Science U.S.A. 105:10450-10455.
- 15 **Marbà**, N. and C.M. Duarte. 2010. Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality.
16 *Global Change Biology* 16: 2366-2375.
- 17 **Marbà**, N., y C.M. Duarte. 1997. Interannual changes in seagrass (*Posidonia oceanica*) growth and environmental
18 change in the Spanish Mediterranean littoral. *Limnology and Oceanography* 42: 800-810.
- 19 **Marfai**, M.A. and L. King 2008: Potential vulnerability implications of coastal inundation due to sea level rise for
20 the coastal zone of Semarang, Indonesia. *Environmental Geology*, **54**, 1235-1245.
- 21 **Mars**, J.C. and Houseknecht, D.W., 2007, Quantitative remote sensing study indicates doubling of coastal erosion
22 rate in past 50 yr along a segment of the Arctic coast of Alaska, *Geology*, v. 35, p. 583-586,
23 doi:10.1130/G23672A.1
- 24 **Martinez**, G., L. Bizikova, D. Blobel and R. Swart. 2011 Emerging Climate Change Coastal Adaptation Strategies
25 and Case Studies Around the World. In *Global Change and Baltic Coastal Zones Coastal Research Library*,
26 2011, Volume 1, Part 4, 249-273, DOI: 10.1007/978-94-007-0400-8_15
- 27 **Massa**, S., Arnaud-Haond, S., Pearson, G., Serrão, E. 2009. Temperature tolerance and survival of intertidal
28 populations of the seagrass *Zostera noltii* (Hornemann) in Southern Europe (Ria Formosa, Portugal).
29 *Hydrobiologia* 619: 195-201.
- 30 Mayot, N., C. F. Boudouresque, A. Leriche, *Comptes Rendus Biologies* **328**, 291 (2005).
- 31 Mazaris, A. D., A. S. Kallimanis, J. Tzanopoulos, S. P. Sgardelis, J. D. Pantis, *Journal of Experimental Marine*
32 *Biology and Ecology* **379**, 23 (2009).
- 33 Mazaris, A. D., G. Matsinos, J. D. Pantis, *Ocean & Coastal Management* **52**, 139 (2009).
- 34 **Mazzotti** S., A. Lambert, M. Van der Kooij, and A. Mainville, 2009: Impact of anthropogenic subsidence on
35 relative sea-level rise in the Fraser River delta. *Geology*, **37**, 771-774.
- 36 **McCulloch** M., S. Fallon, T. Wyndham, E. Hendy, J. Lough, and D. Barnes, 2003: Coral record of increased
37 sediment flux to the inner Great Barrier Reef since European settlement. *Nature*, **421**: 727-730.
- 38 **McFadden** L. 2008. "Exploring the challenges of integrated coastal zone management and reflecting on
39 contributions to 'integration' from geographical thought." *Geographical Journal* 174(4): 299-314.
- 40 **McFadden** L., T. Spencer, and R.J. Nicholls, 2007: Broad-scale modelling of coastal wetlands: What is
41 required? *Hydrobiologia*, **577(1)**, 5-15.
- 42 **McGinnis** M. V. and McGinnis, C. E. 2011: "Adapting to Climate Impacts in California: The Importance of Civic
43 Science in Local Coastal Planning." *Coastal Management* 39(3): 225 - 241.
- 44 **McGranahan** G., D. Balk, and B. Anderson 2007: The rising tide: assessing the risks of climate change and human
45 settlements in low elevation coastal zones. *Environment and Urbanization*, **19**, 17-37.
- 46 **McInnes** K.L., Macadam, I., Hubbert, G.D. and O'Grady, J.G. 2011: An assessment of current and future
47 vulnerability to coastal inundation due to sea level extremes in Victoria, southeast Australia *Int. J. Clim.* DOI:
48 10.1002/joc.3405
- 49 **McKee** K.L., 2011: Biophysical controls on accretion and elevation change in caribbean mangrove
50 ecosystems. *Estuarine, Coastal and Shelf Science*, **91(4)**, 475-483.
- 51 **McKee** K.L., D.R. Cahoon, and I.C. Feller, 2007: Caribbean mangroves adjust to rising sea level through biotic
52 controls on change in soil elevation. *Global Ecology and Biogeography*, **16(5)**, 545-556.
- 53 **McLaughlin** S. and J.A.G. Cooper, 2010: A multi-scale coastal vulnerability index: A tool for coastal
54 managers? **9(3)**, 233-248.

- 1 **McLeod E.**, R. Salm, A. Green and J. Almany 2009. Designing marine protected area networks to address the
2 impacts of climate change. *Frontiers in Ecology and the Environment* 7: 362-370.
- 3 **McLeod, C. D. et al.**, *Biol. Conserv.* **124**, 477 (Aug, 2005).
- 4 **McLeod E.** Hinkel, J. Vafeidis, A.T., Nicholls, R.J., Harvey, N., and Salm, R. 2010a: Sea-level rise vulnerability in
5 the countries of the Coral Triangle Sustainability Science 5, 207-222
- 6 **McLeod E.**, B. Poulter, J. Hinkel, E. Reyes, and R. Salm, 2010b: Sea-level rise impact models and environmental
7 conservation: A review of models and their applications. **53(9)**, 507-517.
- 8 **McMahon, C. R.**, G. C. Hays, *Global Change Biology* **12**, 1330 (2006).
- 9 **McMichael A.J.**, Wilkinson, P., Kovats, R.S., Pattenden, S., Hajat, S., Vajanapoom, N., Niciu, E.M., Mahomed, H.
10 Kingkeow, C., Kosnik, M., O'Neill, M.S., Romieu, I., Ramirez-Aguilar, M., Barreto, M.L, Gouveia, N.,
11 Nikiforov, B. 2008: International study of temperature, heat and urban mortality: the 'ISOTHURM' project
12 *International Journal of Epidemiology* 37:1121-1131
- 13 **McMillan, C.** *Aquat. Bot.* **19**, 369 (1984).
- 14 **Meehl G.A.**, T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M.
15 Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao, 2007: Global Climate
16 Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the*
17 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M.
18 Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press,
19 Cambridge, UK, pp. 747-846.
- 20 **Meissner K. J.**, Lippmann T. and Sen G., A., 2012. Large-scale stress factors affecting coral reefs: open ocean sea
21 surface temperature and surface seawater aragonite saturation over the next 400 years. *Coral Reefs*
- 22 **Melbourne-Thomas J.**, C. R. Johnson, and E. A. Fulton. 2011. Regional-scale scenario analysis for the Meso-
23 American Reef system: Modelling coral reef futures under multiple stressors. *Ecological Modelling* 222
24 (10):1756-1770.
- 25 **Mendelsohn R.** and S. Olmstead, 2009: The economic valuation of environmental amenities and disamenities:
26 Methods and applications. **34**, 325-347.
- 27 **Menendez M.** and P.L. Woodworth, 2010: Changes in extreme high water levels based on a quasi-global tide-gauge
28 dataset. *Journal of Geophysical Research*, **115**, C10011.
- 29 **Menendez M.**, F. J. Mendez, I. J. Losada, and N. E. Graham. 2008. Variability of extreme wave heights in the
30 northeast Pacific Ocean based on buoy measurements, *Geophys. Res. Lett.*, 35, L22607.
- 31 **Mercer J.** 2010. "Disaster risk reduction or climate change adaptation: Are we reinventing the wheel?" *Journal of*
32 *International Development* 22(2): 247-264.
- 33 **Middelburg et al.**, 1997
- 34 **Mieszkwowska N. et al.**, *Hydrobiologia* **555**, 241 (Feb, 2006).
- 35 **Millennium Ecosystem Assessment**, 2005. *Ecosystems and Human Well-being: General Synthesis*. World
36 Resources Institute, Washington, DC.
- 37 **Miller, J., et al.**, *Coral Reefs* **28**, 925 (2009).
- 38 **Milligan J.**, T. O'Riordan, S.A. Nicholson-Cole and A.R. Watkinson. 2009. Nature conservation for future
39 sustainable shorelines: Lessons from seeking to involve the public. *Land Use Policy* 26: 203-213.
- 40 **Milliman J.D.**, and K.L. Farnsworth, 2011: *River Discharge to the Coastal Ocean. A Global Synthesis*. Cambridge
41 University Press, 384pp.
- 42 **Mitchell T.**, M. van Aalst and P.S. Villanueva 2010. Assessing progress on integrating disaster risk reduction and
43 climate change adaptation in development processes. *Strengthening Climate Resilience Discussion Paper 2*,
44 Institute of Development Studies, Brighton, 28 pp.
- 45 **Mitchell T.**, T. Tanner, and E. Wilkinson. 2006. *Overcoming the Barriers: Mainstreaming Climate Change*
46 *Adaptation in Developing Countries: Institute of Development Studies and Tearfund*.
- 47 **Moe, B. et al.**, *Marine Ecology Progress Series* **393**, (2009).
- 48 **Molden D.**, Frenken, K., Barker, R., de Fraiture, C., Mati, B. Svendsen, M., Sadoff, C., Finlayson, M., Atapattu, S.,
49 Giordano, M., Inocencio, A., Lannerstad, M., Manning, N., Molle, F., Smedema, B. and Vallee, D. 2007.
50 Trends in water and agricultural development. In: *Water for food, water for life: A Comprehensive Assessment*
51 *of Water Management in Agriculture* (Ed. Molden, D.) (Earthscan, London, and IWMI, Colombo, pp.57-89)
- 52 **Moller, A. P.**, E. Flensted-Jensen, W. Mardal, *Journal of Animal Ecology* **75**, 657 (2006).
- 53 **Montaggioni L. F.**, 2005. History of Indo-Pacific coral reef systems since the last glaciation: Development patterns
54 and controlling factors. *Earth Science Reviews* 71:1-75.

- 1 Montes-Hugo, M. *et al.*, *Science* **323**, 1470 (Mar, 2009).
- 2 Moore, K. A., J. C. Jarvis, *J. Coast. Res.*, 135 (2008).
- 3 **Moore** K.A., E.C. Shields, D.B. Parrish, and R.J. Orth, 2012: Eelgrass survival in two contrasting systems: Role of
4 turbidity and summer water temperatures. *Marine Ecology-Progress Series*, **448**, 247-258.
- 5 **Moore** K.A., R.L. Wetzel, and R.J. Orth, 1997: Seasonal pulses of turbidity and their relations to eelgrass (*zostera*
6 *marina* L) survival in an estuary. *Journal of Experimental Marine Biology and Ecology*, **215(1)**, 115-134.
- 7 Moore, S. E., H. P. Huntington, *Ecological Applications* **18**, S157 (2008).
- 8 **Moreno** A. and B. Amelung 2009. Climate change and tourist comfort on Europe's beaches in summer: A
9 reassessment. *Coastal Management*, **37**, 550-568
- 10 **Moser** S. C. 2011. Adaptation, mitigation, and their disharmonious discontents. *Climatic Change* online first; DOI
11 10.1007/s10584-011-0106-9.
- 12 **Moser** S. C., and J. A. Ekstrom. 2010. A Framework to Diagnose Barriers to Climate Change Adaptation. *Proc Natl*
13 *Acad Sci* **107** (51):22026-22031.
- 14 **Moser** S. C., and J. Tribbia. 2006/2007. Vulnerability to inundation and climate change impacts in California:
15 Coastal managers' attitudes and perceptions. *Marine Technology Society Journal* **40** (4):35-44.
- 16 **Moser** S. C., R. E. Kasperson, G. Yohe, and J. Agyeman, 2008: Adaptation to climate change in the Northeast
17 United States: Opportunities, processes, constraints. *Mitigation and Adaptation Strategies for Global Change* **13**
18 (5-6):643-659.
- 19 **Moustadraf** J., M. Razack, and M. Sinan 2008: Evaluation of the impacts of climate changes on the coastal Chaouia
20 aquifer, Morocco, using numerical modeling. *Hydrogeology Journal*, **16**, 1411-1426.
- 21 **Mozumder** P., E. Flugman, and T. Randhir, 2011: Adaptation behavior in the face of global climate change: Survey
22 responses from experts and decision makers serving the Florida Keys. *Ocean and Coastal Management* **54**
23 (1):37-44.
- 24 **Mudd** S. M., S. M. Howell, and J. T. Morris, 2009: Impact of dynamic feedbacks between sedimentation, sea-level
25 rise, and biomass production on near-surface marsh stratigraphy and carbon accumulation, *Estuarine Coastal*
26 *Shelf Sci.*, **82**, 377-389, doi:10.1016/j.ecss.2009.01.028.
- 27 **Müller** R., Laepple T., Bartsch I. and Wiencke C., 2009: Impact of oceanic warming on the distribution of seaweeds
28 in polar and cold-temperate waters. *Botanica Marina* **52**:617-638.
- 29 Mumby, P. J., *et al.*, *Mar. Biol.* **139**, 183 (Jul, 2001).
- 30 **Murray** V., G. McBean, M. Bhatt, S. Borsch, T.S. Cheong, W.F. Erian, S. Llosa, F. Nadim, M. Nunez, R. Oyun,
31 and A.G. Suarez, 2012: Case studies. In: *Managing the Risks of Extreme Events and Disasters to Advance*
32 *Climate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D.
33 Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. A Special Report of
34 Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC). Cambridge University
35 Press, Cambridge, UK, and New York, NY, USA, pp. 487-542.
- 36 **Mustafa** D., S. Ahmed, E. Saroch and H. Bell 2011. Pinning down vulnerability: from narratives to numbers.
37 *Disasters* **35**: 62-86.
- 38 **Mustelin** J., Klein, R., Assaid, B., Sitari, T., Khamis, M., Mzee, A. and Haji, T., 2010: Understanding current and
39 future vulnerability in coastal settings: community perceptions and preferences for adaptation in Zanzibar,
40 Tanzania. *Population and Environment* **31(5)**: 371-398.
- 41 **Mustelin** J., M. Khamis, R.G. Klein, A.J. Mzee, T.A. Haji, B. Asseid and T. Sitari, 2011: Coastal forest buffer
42 zones and shoreline change in Zanzibar, Tanzania: practical measures for climate adaptation? In *Experiences of*
43 *Climate Change Adaptation in Africa* *Climate Change Management*, 2011, 133-151, DOI: 10.1007/978-3-642-
44 22315-0_8
- 45 Mydlarz, L. D., C. S. Couch, E. Weil, G. Smith, C. D. Harvell, *Dis. Aquat. Org.* **87**, 67 (Nov, 2009).
- 46 **Nageswara Rao**, K, P. Subrauelu, K.Ch.V. Naga Kumar, G. Demudu, B. Hema Malini, A.S. Rajawat and Ajai, 2010:
47 Impacts of sediment retention by dams on delta shoreline recession: evidences from the Krishna and Godavari
48 deltas, India. *Earth Surface Processes and Landforms*, **35**, 817-827.
- 49 Najjar, R. G. *et al.*, *Estuarine, Coastal and Shelf Science* **86**, 1 (2010).
- 50 Nakane, T., K. Nakaka, H. Bouman, T. Platt, *Estuarine Coastal and Shelf Science* **78**, 796 (2008).
- 51 **Narayan** N., Paul A., Mulitza S. and Schulz M., 2010: Trends in coastal upwelling intensity during the late 20th
52 century. *Ocean Science* **6**:815-823.

- 1 **Narayan, K.** 2006: Climate change impacts on water resources in Guyana. In Climate Variability and Change –
2 Hydrological Impacts [Demuth, S, A. Gustard, E. Planos, F. Scatena, and E. Servat (eds.)]. IAHS publication
3 308, 413-417.
- 4 **Neira C, Grosholz ED, Levin LA, Blake R.** 2006: Mechanisms generating modification of benthos following tidal
5 flat invasion by a *Spartina* hybrid. *Ecol. App.* 16:1391–1404
- 6 **Nevoux, M., H. Weimerskirch, C. Barbraud,** *Oecologia* **162**, 383 (Feb, 2010).
- 7 **Newton K., Cote I. M., Pilling G. M., Jennings S. and Dulvy N. K.,** 2007: Current and future sustainability of island
8 coral reef fisheries. *Current Biology* 17:655-658.
- 9 **Nicholls R.J. and Lowe J.A.,** 2004: Benefits of mitigation of climate change for coastal areas. *Global Environmental*
10 *Change*, 14, 229-244
- 11 **Nicholls, R.** 1995: Coastal mega-cities and climate change. *GeoJournal*, 37, 369-379.
- 12 **Nicholls, R., S. Brown, S. Hanson, and J. Hinkel,** 2010: Economics of Coastal Zone Adaptation to Climate Change
13 (Not Formally Peer-Reviewed), The World Bank.
- 14 **Nicholls, R.J.** 2007: Adaptation Options for Coastal Areas And Infrastructure: An Analysis For 2030. In Adaptation
15 Options for Coastal Areas And Infrastructure: An Analysis For 2030, 35 pp. Bonn: UNFCCC.
- 16 **Nicholls, R.J.** 2010: Impacts of and responses to sea-level rise. Chap. 2. In: Understanding Sea-level rise and
17 variability. Wiley-Blackwell. ISBN 978-4443-3451-7. pp 17-43.
- 18 **Nicholls, R.J. and R.S.J. Tol,** 2006: Impacts and responses to sea-level rise: A global analysis of the SRES scenarios
19 over the twenty-first century. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and*
20 *Engineering Sciences*, **364(1841)**, 1073-1095.
- 21 **Nicholls, R.J., and A. Cazenave,** 2010: Sea-level rise and its impact on coastal zones. *Science*, 328(5985), 1517–
22 1520.
- 23 **Nicholls, R.J., Marinova, N., Lowe, J.A., Brown, S., Vellinga, P., D. Gusmao, D., Hinkel, J., Tol, R.S.J.** 2011: Sea-
24 level rise and its possible impacts given a ‘beyond 4°C world’ in the twenty-first century *Phil. Trans. R. Soc. A*
25 369 (1934), 161–181.
- 26 **Nicholls, R.J., P.P. Wong, V.R. Burkett, J.O. Codignotto, J.E. Hay, R.F. McLean, S. Ragoonaden and C.D.**
27 **Woodroffe,** 2007: Coastal systems and low-lying areas. *Climate Change 2007: Impacts, Adaptation and*
28 *Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental*
29 *Panel on Climate Change, [M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson,*
30 *(eds)], Cambridge University Press, Cambridge, UK, 315-356.*
- 31 **Nicholls, R.J., S. Hanson, C. Herweijer, N. Patmore, S. Hallegatte, J. Corfee-Morlot, J. Château, and R. Muir-Wood**
32 **2008: Ranking Port Cities with High Exposure and Vulnerability to Climate Extremes. OECD Environment**
33 **Working Papers No. 1, OECD Publishing, 62 pp.**
- 34 **Nicholls R., Sally Brown, Susan Hanson and Jochen Hinkel** 2010: Economics of Coastal Zone Adaptation to
35 Climate Change (not formally peer-reviewed). In Economics of Coastal Zone Adaptation to Climate Change
36 (not formally peer-reviewed): The World Bank.
- 37 **Nichols R.J. and Tol R.S.J. (2006):** Impacts and responses to sea-level rise: a global analysis of the SRES scenarios
38 over the twenty-first century. *Philosophical Transactions of the Royal Society A* **364**, 1073-95.
- 39 **Nixon S. W.,** 1982. Nutrient dynamics, primary production and fisheries yields of lagoons. *Oceanologica acta* N°
40 SP:357-371.
- 41 **Nixon, F.C., E.G. Reinhardt, and R. Rothaus,** 2009: Foraminifera and tidal notches: Dating neotectonic events at
42 korphos, greece. *Marine Geology*, **257**
- 43 **NOAA Digital Coast Data Access Viewer :** <http://csc.noaa.gov/dataviewer/?keyword=lidar#>
- 44 **Nowak, D. J., and D. E. Crane.** 2002. Carbon storage and sequestration by urban trees in the USA. *Environmental*
45 *Pollution* 116 (3):381-389.
- 46 **Nowak, D. J., D. E. Crane, and J. C. Stevens.** 2006. Air pollution removal by urban trees and shrubs in the United
47 States. *Urban Forestry and Urban Greening* 4 (3-4):115-123.
- 48 **Nursey-Bray, M. and Shaw, J. R.** 2010. "Australia, Climate Change and the Sea Change." *International Journal of*
49 *Environmental, Cultural, Economic and Social Sustainability* 6(1): 67-80.
- 50 **Oke, P. R., J. H. Middleton,** Nutrient Enrichment off Port Stephens: the Role of the East. *Australian Current*
51 *Continental Shelf Research* **21**, 587-606 (2001).
- 52 **Onozuka, D., Hashizume, M., and Hagihara, A.,**2010: Effects of weather variability on infectious gastroenteritis.
53 *Epidemiol. Infect.* 138: 236–243

- 1 **O'Rourke, D.**, and S. Connolly. 2003. Just oil? The distribution of environmental and social impacts of oil
2 production and consumption. *Annual Review of Environment and Resources* 28 (1):587-617.
- 3 **Orth R. J.**, Carruthers T. J. B., Dennison W. C., Duarte C. M., Fourqurean J. W., Heck Jr K. L., Hughes A. R.,
4 Kendrick G. A., Kenworthy W. J. and Olyarnik S., 2006. A global crisis for seagrass ecosystems. *Bioscience*
5 56:987-996.
- 6 **Orth, R. J.**, K. A. Moore, 1983: Chesapeake Bay: An unprecedented decline in submerged aquatic
7 vegetation. *Science* **222**, 51-53.
- 8 **Orth, R.J.** Carruthers, T.J.B. Dennison, W.C. Duarte, C.M., Fourqurean, J.W. Heck Jr., K.L. Hughes, A.R.,
9 Kendrick, G.A. Kenworthy, K.J., Olyarnik, S., Short, F.T. Waycott, M. and Williams, S.L., 2006: A Global
10 Crisis for Seagrass Ecosystems. *Bioscience* **56 (12)**, 987-996.
- 11 **OzCoasts** website: <http://www.ozcoasts.gov.au/>
- 12 **Ozyurt, G.** and A. Ergin, 2009: Application of sea level rise vulnerability assessment model to selected coastal
13 areas of Turkey. *Journal of Coastal Research*: 248-251.
- 14 **Paaijmans, K.P.**, S. Blanford, A.S. Bell, J.I. Blanford, A.F. Read, and M.B. Thomas 2010: Influence of climate on
15 malaria transmission depends on daily temperature variation. *PNAS*, 107(34), 15135-15139.
- 16 **Pall P.**, Aina T., Stone D. A., Stott P. A., Nozawa T., Hilberts A. G. J., Lohmann D. and Allen M. R., 2011.
17 Anthropogenic greenhouse gas contribution to flood risk in England and Wales in autumn 2000. *Nature*
18 470:380-384.
- 19 **Pandolfi J. M.**, Connolly S. R., Marshall D. J. and Cohen A. L., 2011. Projecting coral reef futures under global
20 warming and ocean acidification. *Science* 333:418-422.
- 21 **Parkinson, R. W.**, 1989:Decelerating Holocene sea-level rise and its influence in southwest Florida coastal
22 evolution: A transgressive/ regressive stratigraphy *Journal of Sedimentary Petrology* 59, 960-972.
- 23 **Parkinson, R. W.**, R. D. Delaune, J. R. White, 1994: Holocene Sea-Level Rise and the Fate of Mangrove Forests
24 within the Wider Caribbean Region. *J. Coast. Res.* **10**, 1077-1086.
- 25 **Parmesan, C.**, G. Yohe, 2003: A globally coherent fingerprint of climate change impacts across natural systems
26 *Nature* 421, 37-42.
- 27 **Parry, M.**, N. Arnell, P. Berry, D. Dodman, S. Fankhauser, C. Hope, S. Kovats, R. Nicholls, D. Satterthwaite, R.
28 Tiffin, and T. Wheeler, 2009: Assessing the costs of adaptation to climate change: A review of the UNFCCC
29 and other recent estimates. *International Institute for Environment and Development and Grantham Institute for*
30 *Climate Change, London, U.K.*
- 31 **Pataki, D. E.**, R. J. Alig, A. S. Fung, N. E. Golubiewski, C. A. Kennedy, E. G. McPherson, D. J. Nowak, R. V.
32 Pouyat, and P. Romero Lankao. 2006. Urban ecosystems and the North American carbon cycle. *Global Change*
33 *Biology* 12 (11):2092-2102.
- 34 **Patt, A. G.**, D. P. van Vuuren, F. Berkhout, A. Aaheim, A. F. Hof, M. Isaac and R. Mechler 2010. Adaptation in
35 integrated assessment modeling: where do we stand? *Climatic Change* 99: 383-402.
- 36 **Pauly D.** and Yáñez-Arancibia A., 1994: Fisheries in coastal lagoons. In: Kjerfve B. (Ed.), *Coastal lagoon*
37 *processes*, pp. 377-399. Amsterdam: Elsevier.
- 38 **Pearce, T.**, Smit, B., Duerden, F., Ford, J.D., Goose, A., Kataoyak, F., 2010: Inuit vulnerability and adaptive
39 capacity to climate change in Ulukhaktok, Northwest Territories, Canada. *Polar Record* 46 (237): 157-177.
- 40 **Pearson, G. A.**, A. Lago-Leston, C. Mota, 2009: Frayed at the edges: Selective pressure and adaptive response to
41 abiotic stressors are mismatched in low diversity edge populations. *J. Ecol.* **97**, 450-462.
- 42 **Peck L. S.**, Clark M. S., Morley S. A., Massey A. and Rossetti H., 2009a. Animal temperature limits and ecological
43 relevance: effects of size, activity and rates of change. *Functional Ecology* 23:248-256.
- 44 **Peck L. S.**, Massey A., Thorne M. A. S. and Clark M. S., 2009b. Lack of acclimation in *Ophionotus victoriae*: brittle
45 stars are not fish. *Polar Biology* 32:399-402.
- 46 **Peirano, A.**, Damasso, V., Montefalcone, M., Morri, C., Bianchi, C.N., 2005: Growth of the seagrass *Posidonia*
47 *oceanica* (L.) Delile in Liguria (NW Italy, Mediterranean Sea): relations with climate, alien species invasion and
48 anthropogenic impacts. *Marine Pollution Bulletin* **50**, 817-822.
- 49 **Pemberton, D.**, R. Gales, 2004: Australian fur seals (*Arctocephalus pusillus doriferous*) in Tasmania: population
50 size and status. *Wildlife Research* **31**, 301-309.
- 51 **Peperzak, L.**, 2003: Climate change and harmful algal blooms in the North Sea *Acta Oecologica-International*
52 *Journal of Ecology* **24**, S139-S144.
- 53 **Peperzak, L.**, 2005: Future increase in harmful algal blooms in the North Sea due to climate change *Water Science*
54 *and Technology* **51**, 31-36.

- 1 **Pérez, Á.A.**, B.H. Fernández and R.C. Gatti, eds., 2010: Building resilience to climate change: Ecosystem-based
2 adaptation and lessons from the field. IUCN, Gland, Ecosystem Management Series No. 9, pp. 160-165.
- 3 **Perry, A. L.**, P. J. Low, J. R. Ellis, J. D. Reynolds, 2005: Change and distribution shifts in marine fishes. *Science*
4 **308**, 1912-1915.
- 5 **Piao S. L.**, Friedlingstein P., Ciais P., de Noblet-Ducoudre N., Labat D. and Zaehle S., 2007: Changes in climate and
6 land use have a larger direct impact than rising CO₂ on global river runoff trends. Proceedings of the National
7 Academy of Science U.S.A. 104:15242-15247.
- 8 Pike, D. A. Do green turtles modify their nesting seasons in response to environmental temperatures? *Chelonian*
9 *Conservation and Biology* **8**, 43-47 (May, 2009).
- 10 Pike, D. A., J. C. Stiner, Sea turtle species vary in their susceptibility to tropical cyclones *Oecologia* **153**, 471-478
11 (2007).
- 12 Piquet, A. M. T., H. Bolhuis, A. T. Davidson, P. G. Thomson, A. G. J. Buma, Diversity and dynamics of Antarctic
13 marine microbial eukaryotes under manipulated environmental UV radiation *FEMS Microbiology Ecology* **66**,
14 352-366 (2008).
- 15 **Plant, N.**, Stockdon, H., Sallenger, A., Turco, M., East, J., Taylor, A., and Shaffer, W., 2010, Forecasting hurricane
16 impact on coastal topography, EOS, Transactions of the American Geophysical Union, Vol 91, No 7, P.65–72
- 17 **Plestan, M.**, Ponsoero, A. & Yésou, P. 2009. Forte abondance du Puffin des Baléares *Puffinus mauretanicus* en
18 Bretagne (hiver 2007-2008). *Ornithos* 16: 209-213.
- 19 **Polack, E.** 2010: Integrating climate change into regional disaster risk management at the Mekong River
20 Commission. Strengthening Climate Resilience Discussion Paper 4, Institute of Development Studies, Brighton,
21 36 pp.
- 22 **Polasky, S.** and K. Segerson 2009: Integrating Ecology and Economics in the Study of Ecosystem Services: Some
23 Lessons Learned. *Annual Review of Resource Economics* 1: 409-434.
- 24 **Pollack, J.B.**, T.A. Palmer, and P.A. Montagna, 2011: Long-term trends in the response of benthic macrofauna to
25 climate variability in the lavaca-colorado estuary, texas. *Marine Ecology-Progress Series*, **436**, 67-80.
- 26 **Poloczanska E. S.**, Smith S., Fauconnet L., Healy J., Tibbetts I. R., Burrows M. T. and Richardson A. J., 2011.
27 Little change in the distribution of rocky shore faunal communities on the Australian east coast after 50 years of
28 rapid warming. *Journal of Experimental Marine Biology and Ecology* 400:145-154.
- 29 Polovina, J. J., W. R. Haight, R. B. Moffitt, F. A. Parrish, The role of benthic habitat, oceanography and fishing on
30 the population dynamics of the spiny lobster *Panulirus marginatus* (Decapoda, Palinuridae) in the Hawaiian
31 Archipelago, *Crustaceana* **68**, 203-212 (1995).
- 32 Polovina, J.J., G.T Mitchum, N.E Graham, M.P Craig, E.E Demartini, E.N Flint. Physical and biological
33 consequences of a climate event in the central North Pacific. *Fisheries Oceanography* **3**, 15-21 (1994).
- 34 Porter, J. W., O. W. Meier, Quantification of loss and change in Floridian reef coral populations. *American*
35 *Zoologist* **32**, 625-640 (1992).
- 36 **Porzio L.**, Buia M. C. and Hall-Spencer J. M., 2011: Effects of ocean acidification on macroalgal communities.
37 *Journal of Experimental Marine Biology and Ecology* 400:278-287.
- 38 Prada, C., E. Weil, P. M. Yoshioka, Octocoral bleaching during unusual thermal stress. *Coral Reefs* **29**, 41-45 (Mar,
39 2010).
- 40 **Precht W. F.** and Aronson R. B., 2004: Climate flickers and range shifts of reef corals. *Frontiers in Ecology and the*
41 *Environment* 2:307-314.
- 42 Preen, A. R., W. J. L. Long, R. G. Coles, Flood and cyclone related loss and partial recovery of more than 1000 km²
43 of seagrass in Hervey Bay, Queensland. Australia. *Aquat. Bot.* **52**, 3-17 (1995).
44 *Proc. Nat. Acad. Sci.* **102**, 2826 (2005).
- 45 **Provoost P.**, van Heuven S., Soetaert K., Laane R. W. P. M. and Middelburg J. J., 2010: Seasonal and long-term
46 changes in pH in the Dutch coastal zone. *Biogeosciences* 7:3869-3878.
- 47 **Pruszk, Z.**, and E. Zawadzka, 2008: Potential implications of sea-level rise for Poland. *Journal of Coastal*
48 *Research*, 24(2), 410-422
- 49 **Przeslawski R.**, Ah Yong S., Byrne M., Wörheide G. and Hutchings P., 2008: Beyond corals and fish: the effects of
50 climate change on noncoral benthic invertebrates of tropical reefs. *Global Change Biology* 14:2773-2795.
- 51 **Purcell, K.M.**, P.L. Klerks and P.L. Leberg 2010: Adaptation to sea level rise: does local adaptation influence the
52 demography of coastal fish populations. *Jour of Fish Biology*, 77 : 1209-1218.
- 53 Queensland, Australia *Mar. Freshw. Res.* **47**, 763-771 (1996).

- 1 **Rabalais**, N. N., Turner, R. E., Diaz, R. J., and Justic, D., 2009: Climate change and eutrophication of coastal
2 waters, *ICES J. Mar. Sci.*, 1528-1537.
- 3 **Rahman**, M.H., T. Lund, and I. Bryceson 2011: Salinity impacts on agro-biodiversity in three coastal, rural villages
4 of Bangladesh. *Ocean and Coastal Management*, 54, 455-468.
- 5 **Rahmstorf**, S. 2007: A Semi-Empirical Approach to Projecting Future Sea-Level Rise. *Science* 315 (5810):368-
6 370.
- 7 **Rahmstorf**, S. 2010: A new view on sea level rise. *Nature Reports Climate Change*, 4: 44-45
- 8 **Raihan**, M.S, M.J. Huq, N.G. Alsted and M.H. Andreassen 2010: Understanding climate change from below,
9 addressing barriers from above: Practical experience and learning from a community-based adaptation project
10 in Bangladesh. *ActionAid Bangladesh*, 98 pp.
- 11 **Ramasamy**, R. and S.N. Surendran 2011: Possible impact of rising sea levels on vector-borne infectious diseases.
12 *BMC Infectious Diseases*, 11:18. doi:10.1186/1471-2334-11-18.
- 13 **Ranasinghe**, R, Callaghan, D., and Stive, M., 2012, Estimating coastal recession due to sea level rise: beyond the
14 Bruun rule, *Climatic Change*, **110(3-4)**, 561-574.
- 15 **Ranasinghe**, R. and M.J.F. Stive, 2009: Rising seas and retreating coastlines. *Climatic Change*, **97(3)**, 465-468.
- 16 **Rappaport**, E.N., 2000: Loss of life in the United States associated with recent Atlantic tropical cyclones. *Bulletin*
17 *of the American meteorological Society*, **81(9)**, 2065-2073.
- 18 **Rasheed** M. A. and Unsworth R. K. F., 2011: Long-term climate-associated dynamics of a tropical seagrass
19 meadow: implications for the future. *Marine Ecology Progress Series* 422:93-103.
- 20 Ratcliffe, N., S. Schmitt, A. Mayo, J. Tratalos, A. Drewitt, *Seabird* **21**, 55 (2008). Reusch, T. B. H., A. Ehlers, A.
21 Hammerli, B. Worm, Ecosystem recovery after climatic extremes enhanced by genotypic diversity
- 22 **Rau** GH, McLeod E, Hoegh-Guldberg O., 2012: The need for new conservation strategies and policies in a high
23 CO2 world. *Nature Climate Change*.
- 24 **Raudsepp-Hearne**, C., G. D. Peterson, M. Tengo, E. M. Bennett, T. Holland, K. Benessaiah, G. K. MacDonald and
25 L. Pfeifer 2010. Untangling the Environmentalist's Paradox: Why Is Human Well-being Increasing as
26 Ecosystem Services Degrade? *Bioscience* **60(8)**, 576-589.
- 27 **Redfield**, A. C. 1972 Development of a New England salt marsh. *Ecological Monographs*, **42(2)**, 201-237.
- 28 **Reeve**, D.E., 1998: Coastal flood risk assessment. *Journal of Waterway, Port, Coastal, and Ocean Engineering* **124**,
29 219.
- 30 **Reguero**, B., Mendez, F.J., I.J. Losada 2012: Variability of multivariate wave climate in Latin America and the
31 Caribbean. *Global and Planetary Change*. (under review).
- 32 **Reid**, H., M. Alam, R. Berger, T. Cannon, S. Huq, and Angela Milligan (eds.). 2009: Special Issue: Community-
33 based adaptation to climate change. *Participatory Learning and Action* 60.
- 34 **Reusch**, T.B.H., A. Ehlers, A. Hämmerli, and B. Worm, 2005: Ecosystem recovery after climatic extremes
35 enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences* 102: 2826-2831.
- 36 **Revell**, D.L., R. Battalio, B. Spear, P. Ruggiero, and J. Vandever, 2011: A methodology for predicting future coastal
37 hazards due to sea-level rise on the California coast. *Climatic Change*, **109(SUPPL. 1)**, 251-276.
- 38 **Revenga**, C., Nackoney, J., Hoshino, E., Kura, Y., Maidens, J., 2003: AS 12 Irrawaddy. *Watersheds of the World*,
39 Water Resources Institute.
- 40 **Reynaud** S., Leclercq N., Romaine-Lioud S., Ferrier-Pagès C., Jaubert J. and Gattuso J.-P., 2003: Interacting effects
41 of CO2 partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Global*
42 *Change Biology* 9:1660-1668.
- 43 Rhein M. and *et al.*, in prep. Observations: Ocean. In: XXX (Eds.), IPCC WGI Fifth Assessment Report.
- 44 Richardson, A. J., D. S. Schoeman, Climate Impact on Plankton Ecosystems in the Northeast Atlantic. *Science* 305,
45 1609-1612 (2004)
- 46 **Ridgeway**, K.R. 2007: Long-term trend and decadal variability of the southward penetration of the East Australian
47 Current. *Geophys. Res. Lett.* 34, L13613, doi:10.1029/2007GL030393
- 48 **Riebesell** U., 2008: Acid test for marine biodiversity. *Nature* 454:46-47.
- 49 **Rivadeneira** M. M. and Fernández M., 2005: Shifts in southern endpoints of distribution in rocky intertidal species
50 along the south-eastern Pacific coast. *Journal of Biogeography* 32:203-209.
- 51 **Rodolfo-Metalpa** R., Houlbrèque F., Tambutté É., Boisson F., Baggini C., Patti F. P., Jeffree R., Fine M., Foggo A.,
52 Gattuso J.-P. and Hall-Spencer J. M., 2011: Coral and mollusc resistance to ocean acidification adversely
53 affected by warming. *Nature Climate Change* 1:308-312.

- 1 Rodriguez, R. W., R. M. T. Webb, D. M. Bush, Another Look at the Impact of Hurricane Hugo on the Shelf and
2 Coastal Resources of Puerto Rico, U.S.A. *J. Coast. Res.* **10**, 278-296 (Spr, 1994).
- 3 Rogers, K., K. M. Wilton, N. Saintilan, Vegetation change and surface elevation dynamics in estuarine wetlands of
4 southeast Australia *Estuarine Coastal and Shelf Science* **66**, 559-569 (2006).
- 5 **Romieu**, E., Welle, T., Schneiderbauer, S., Pelling, M. and Vinchon, C. 2010: "Vulnerability assessment within
6 climate change and natural hazard contexts: revealing gaps and synergies through coastal applications."
7 *Sustainability Science* 5(2): 159-170.
- 8 Root, T.L., J.T. Price, K.R. Hall, S.H. Schneider, C. Rosenzweig, and J.A. Pounds, 2003: Fingerprints of global
9 warming on wild animals and plants. *Nature*, **421**, 57-60,
- 10 Rose, G. A., Capelin (*Mallotus villosus*) distribution and climate: a sea "canary" for ecosystem change *Ices Journal*
11 *of Marine Science* **62**, 1524-1530 (2005).
- 12 **Rosenzweig**, C., Solecki, W.D., Blake, R., Bowman, M., Faris, C., Gornitz, V., Horton, R., Jacob, K., LeBlanc, A.,
13 Lichenko, R., Linkin, M., Major, D., O'Grady, L.P., Sussman, E., Yohe, G., Zimmerman, R., 2011: Developing
14 coastal adaptation to climate change in the New York City infrastructure-shed: process, approach, tools, and
15 strategies. *Climatic Change* **106(1)**, 93-127.
- 16 **Rozema** J, Dorel F, Janissen R, Lenssen G, Broekman R, et al. 1991: Effect of elevated atmospheric CO2 on
17 growth, photosynthesis and water relations of salt marsh grass species. *Aquat. Bot.* 39:45–55
- 18 **Ruddiman** W. F., 2007. The early anthropogenic hypothesis: challenges and responses. *Reviews in Geophysics* 45,
19 RG4001. doi:10.1029/2006RG000207.
- 20 Ryan et al., 2011
- 21 Sahid, 2009
- 22 **Saintilan**, N., R. J. Williams, 1999: Mangrove transgression into saltmarsh environments in south-east Australia.
23 *Global Ecology and Biogeography* **8 (2)**, 117-124.
- 24 **Sakabe**, R. and J.M. Lyle, 2010: The influence of tidal cycles and freshwater inflow on the distribution and
25 movement of an estuarine resident fish *acanthopagrus butcheri*. *Journal of Fish Biology*, **77(3)**, 643-660.
- 26 **Sales Jr**, R. F. M. 2009: Vulnerability and adaptation of coastal communities to climate variability and sea-level
27 rise: Their implications for integrated coastal management in Cavite City, Philippines. *Ocean and Coastal*
28 *Management* 52 (7):395-404.
- 29 **Salisbury** J., Green M., Hunt C. and Campbell J., 2008. Coastal acidification by rivers: a new threat to shellfish?
30 *Eos, Transactions, American Geophysical Union* 89:513.
- 31 **Sallenger**, A., R. Morton, C. Fletcher, E. R. Thieler, and P. Howd, 2000, Discussion of 'Sea level rise shown to drive
32 coastal erosion' by Leatherman et al. (2000): EOS, Trans. American Geophysical Union, Vol. 81, No. 38, p.
33 436.
- 34 **Sanchez-Arcilla**, A., J.A. Jimenez, H.I. Valdemoro, and V. Gracia, 1998: Implications of climatic change on
35 Spanish Mediterranean low-lying coasts: The Ebro Delta case. *Journal of Coastal Research*, 24, 306–316.
- 36 **Saroar**, M., and J. K. Routray, 2010: Adaptation in situ or retreat? a multivariate approach to explore the factors that
37 guide the peoples' preference against the impacts of sea level rise in coastal Bangladesh. *Local Environment:*
38 *The International Journal of Justice and Sustainability* 15 (7):663 - 686.
- 39 **Schanz**, A. and H. Asmus, 2003: Impact of hydrodynamics on development and morphology of intertidal seagrasses
40 in the wadden sea. *Marine Ecology-Progress Series*, **261**, 123-134.
- 41 **Schleupner**, C. 2008: Evaluation of coastal squeeze and its consequences for the Caribbean island Martinique.
42 *Ocean and Coastal Management*, 51, 383-390.
- 43 **Schofield**, G. Bishop, C.M., Katselidis, K.A., Dimopoulos, P., Pantis, J.D., Hays, G.C., 2009: Microhabitat
44 selection by sea turtles in a dynamic thermal marine environment *Journal of Animal Ecology* **78**, 14-21.
- 45 **Schuerch**, M., J. Rapaglia, V. Liebetrau, A. Vafeidis, and K. Reise, 2012: Salt marsh accretion and storm tide
46 variation: An example from a barrier island in the north sea. *Estuaries and Coasts*, **35(2)**, 486-500.
- 47 **Schwierz**, C., Köllner-Heck, P., Zenklusen Mutter, E., Bresch, D.N., Vidale, P.-L., Wild, M. and C. Schär, 2010:
48 Modelling European winter wind storm losses in current and future climate. *Climate Change*, **101**, 485-514.
- 49 **Scott**, D., B. Amelung, S. Becken, J.P. Ceron, G. Dubois, S. Gossling, P. Peeters, and M.C. Simpson, 2008. Climate
50 Change and Tourism: Responding to Global Challenges. United Nations World Tourism Organization, Madrid,
51 Spain, 256 pp UNWTO, 2008: Climate change and tourism: Responding to Global Challenges. 269 pp.
- 52 **Secretariat of the Pacific Regional Environment Programme (SPREP)**: [http://www.sprep.org/Climate-](http://www.sprep.org/Climate-Change/climate-change-overview)
53 [Change/climate-change-overview](http://www.sprep.org/Climate-Change/climate-change-overview)
- 54 Seddon, S., A. C. Cheshire, *Mar. Ecol.-Prog. Ser.* **220**, 119 (2001).

- 1 Seddon, S., R. M. Connolly, K. S. Edyvane, Large-scale seagrass dieback in northern Spencer Gulf, South Australia.
2 *Aquat. Bot.* **66**, 297-310 (Apr, 2000).
- 3 **Selig** E. R., K.S. Casey and J.F. Bruno 2012: Temperature-driven coral decline: the role of marine protected areas.
4 *Global Change Biology*, 18: 1561-1570.
- 5 **Seneviratne**, S.I., N. Nicholls, D. Easterling, C.M. Goodess, S. Kanae, J. Kossin, Y. Luo, J. Marengo, K. McInnes,
6 M. Rahimi, M. Reichstein, A. Sorteberg, C. Vera, and X. Zhang, 2012: Changes in climate extremes and their
7 impacts on the natural physical environment. In: *Managing the Risks of Extreme Events and Disasters to*
8 *Advance Climate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi,
9 M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. A Special
10 Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC). Cambridge
11 University Press, Cambridge, UK, and New York, NY, USA, pp. 109-230.
- 12 **Shaffer** G., Olsen S. M. and Pedersen J. O. P., 2009: Long-term ocean oxygen depletion in response to carbon
13 dioxide emissions from fossil fuels. *Nature Geoscience* 2:105-109.
- 14 **Shahid**, S. 2009: Probable impacts of climate change on public health in Bangladesh. *Asia Pacific Journal of Public*
15 *Health*, 22, 310-319.
- 16 **Shaw**, R., J. M. Pulhin, and J. J. Pereira eds. 2010: *Climate Change Adaptation and Disaster Risk Reduction: Issues*
17 *and Challenges*. Bingley, UK: Emerald Group Publishing.
- 18 **Sheik Mujabar**, P. and N. Chandrasekar (In Press): Coastal erosion hazard and vulnerability assessment for
19 southern coastal Tamil Nadu of India by using remote sensing and GIS. *Natural Hazards*.
- 20 **Sheppard**, S. R. J., A. Shaw, D. Flanders, S. Burch, A. Wiek, J. Carmichael, J. Robinson and S. Cohen 2011. Future
21 visioning of local climate change: A framework for community engagement and planning with scenarios and
22 visualisation. *Futures* **43(4)**, 400-412.
- 23 **Shipman** B. and Stojanovic, T. 2007: "Facts, Fictions, and Failures of Integrated Coastal Zone Management in
24 Europe." *Coastal Management* 35(2): 375 - 398.
- 25 **Short**, F.T. and H.A. Neckles, 1999: The effects of global climate change on seagrasses. *Aquatic Botany*, **63(3-4)**,
26 169-196.
- 27 **Short**, F.T., B. Polidoro, S.R. Livingstone, K.E. Carpenter, S. Bandeira, J.S. Bujang, H.P. Calumpong, T.J.B.
28 Carruthers, R.G. Coles, W.C. Dennison, P.L.A. Erftemeijer, M.D. Fortes, A.S. Freeman, T.G. Jagtap, A.H.M.
29 Kamal, G.A. Kendrick, W.J. Kenworthy, Y.A. La Nafie, I.M. Nasution, R.J. Orth, A. Prathep, J.C. Sanciangco,
30 B. van Tussenbroek, S.G. Vergara, M. Waycott, and J.C. Zieman, 2011: A global crisis for seagrass
31 ecosystems. *Biological Conservation*, **144(7)**, 1961-1971.
- 32 **Sietz**, D., M. Boschütz, and R. J. T. Klein 2011: Mainstreaming climate adaptation into development assistance:
33 rationale, institutional barriers and opportunities in Mozambique. *Environmental Science and Policy* 14 (4):493-
34 502.
- 35 **Silva**, R., C. Coelho, et al. (2007): "Dynamic Numerical Simulation of Medium-term Coastal Evolution of the West
36 Coast of Portugal." *Journal of Coastal Research*: 263-267.
- 37 **Silverman** J., Lazar B., Cao L., Caldeira K. and Erez J., 2009: Coral reefs may start dissolving when atmospheric
38 CO2 doubles. *Geophysical Research Letters* 36, L05606. doi:10.1029/2008GL036282.
- 39 **Simeoni**, U. and C. Corbau, 2009: A review of the Delta Po evolution (Italy) related to climatic changes and human
40 impacts. *Geomorphology*, 107, 64–71.
- 41 **Simpson**, M.C., D. Scott, M. Harrison, R. Slim, N. Silver, E. O’Keeffe, S. Harrison et al 2011: Quantification and
42 Magnitude of Losses and Damages Resulting from the Impacts of Climate Change: Modelling the
43 Transformational Impacts and Costs of Sea Level Rise in the Caribbean. (Key Points and Summary for Policy
44 Makers Document). United Nations Development Programme (UNDP), Barbados, West Indies, 30 pp.
- 45 **Sims**, D. W., V. J. Wearmouth, M. J. Genner, A. J. Southward, S. J. Hawkins, *Journal of Animal Ecology* **73**, 333
46 (Mar, 2004).
- 47 **Smart**, J., J. A. Gill, Climate change and the potential impact on breeding waders in the UK. *Wader Study Group*
48 *Bulletin* **100**, 80-85 (2003).
- 49 **Smith** S. V., 2005a: Length of the global coastal zone. In: Crossland C. J., Kremer H. H., Lindeboom H. J., Marshall
50 Crossland J. I. and Le Tissier M. D. A. (Eds.), *Coastal fluxes in the anthropocene*, pp. 3. Berlin: Springer-
51 Verlag.
- 52 **Smith** S., Buddemeier R., Wulff F., Swaney D., Camacho-Ibar V., David L., Dupra V., Kleypas J., San D.-M.,
53 Maria, McLaughlin C. and Sandhei P., 2005b. C, N, P fluxes in the coastal zone. In: Crossland C. J., Kremer H.

- 1 H., Lindeboom H. J., Marshall Crossland J. I. and Le Tissier M. D. A. (Eds.), Coastal fluxes in the
2 anthropocene, pp. 95-143. Berlin: Springer-Verlag.
- 3 **Smith, K.** 2011. We are seven billion. *Nature Climate Change*, 1 : 331-335.
- 4 **Smithers, B. V., D. R. Peck, A. K. Krockenberger, B. C. Congdon,** Elevated sea-surface temperature reduced
5 provisioning and reproductive failure of Wedge-tailed shearwaters (*Puffinus pacificus*) in the Southern Great
6 Barrier Reef. *Mar. Freshw. Res.* **54**, 973-977 (2003).
- 7 **Snoussi, M., T. Ouchani, A. Khouakhi, and I. Niang-Diop,** 2009: Impacts of sea-level rise on the moroccan coastal
8 zone: Quantifying coastal erosion and flooding in the tangier bay. *Computational Geosciences*, **14(4)**, 503-508.
- 9 **Sokolow S.,** 2009. Effects of a changing climate on the dynamics of coral infectious disease: a review of the
10 evidence. *Diseases of Aquatic Organisms* 87:5-18.
- 11 **Spalding E. and M. Hester** 2007: Interactive effects of hydrology and salinity on oligohaline plant species
12 productivity: Implications of relative sea-level rise *Estuaries Coasts* 30:214-225
- 13 **Steneck R. S. Graham, M.H., Bourque, B.J., Corbett, D., Erlandson5, J.M., Estes, J.A. and Tegner, M.J.** Kelp forest
14 ecosystems: biodiversity, stability, resilience and future. *Environ. Conserv.* **29**, 436 (Dec, 2002).
- 15 **Stenseth N. C., Mysterud A., Ottersen G., Hurrell J. W., Chan K. S. and Lima M.,** 2002. Ecological effects of
16 climate fluctuations. *Science* 297:1292-1296.
- 17 **Sterr, H.,** 2008: Assessment of vulnerability and adaptation to sea-level rise for the coastal zone of Germany.
18 *Journal of Coastal Research*, 24(2), 380-393
- 19 **Steyl, G. and Dennis, I.,** 2010: Review of coastal-area aquifers in Africa, *Hydrogeology Journal* 18: 217–225
- 20 **Stinchcomb G. E., Messner T. C., Driese S. G., Nordt L. C. and Stewart R. M.,** 2011. Pre-colonial (AD 1100-1600)
21 sedimentation related to prehistoric maize agriculture and climate change in eastern North America. *Geology*
22 39:363-366.
- 23 **Stive, M.J.C., L.O. Fresco, P. Kabat, B.W.A.H. Parmet, and C.P. Veerman** 2011: How the Dutch plan to stay dry
24 over the next century. *Proceedings of Institution of Civil Engineers*. 164, pp. 114-121.
- 25 **Stojanovic, T.A. and R.C. Ballinger** 2009: Integrated Coastal Management: A comparative analysis of four UK
26 initiatives. *Applied Geography*, 29 : 49-62.
- 27 **Stokes, J., and A. Horvath.** 2006: Life Cycle Energy Assessment of Alternative Water Supply Systems (9 pp). *The*
28 *International Journal of Life Cycle Assessment* 11 (5):335-343.
- 29 **Storbjörk, S.** 2010: "It Takes More to Get a Ship to Change Course": Barriers for Organizational Learning and
30 Local Climate Adaptation in Sweden. *Journal of Environmental Policy and Planning* 12 (3):235 - 254.
- 31 **Storbjörk, S., and J. Hedrén** 2011: Institutional capacity-building for targeting sea-level rise in the climate
32 adaptation of Swedish coastal zone management. *Lessons from Coastby. Ocean and Coastal Management* 54
33 (3):265-273
- 34 **Storlazzi C. D., Elias E., Field M. E. and Presto M. K.,** 2011: Numerical modeling of the impact of sea-level rise on
35 fringing coral reef hydrodynamics and sediment transport. *Coral Reefs* 30:83-96.
- 36 **Stratten, L., O'Neill, M.S., Kruk, M.E., Bell, M. L.** 2008: The persistent problem of malaria: Addressing the
37 fundamental causes of a global killer. *Social Science and Medicine* 67 (2008) 854–862.
- 38 **Straw, P., N. Saintilan, G. C. Boere, C. A. Galbraith, D. A. Stroud,** *Waterbirds around the world: a global overview*
39 *of the conservation, management and research of the world's waterbird flyways. International conference on*
40 *waterbirds held in Edinburgh in April 2004.*, 717 (2006).
- 41 **Sverdrup H. U., Johnson M. W. and Fleming R. H.,** 1942: The oceans, their physics, chemistry, and general
42 biology. 1087 p. New York: Prentice-Hall.
- 43 **Syvitski, J.P.M., A.J. Kettner, A. Correggiari, and B.W. Nelson,** 2005: Distributary channels and their impact on
44 sediment dispersal. *Marine Geology*, 222–223, 75–94.
- 45 **Syvitski, J.P.M. and A. Kettner,** 2011: Sediment flux and the Anthropocene. *Philosophical Transactions of Royal*
46 *Society A*, 369, 957-975
- 47 **Syvitski, J.P.M.,** 2008: Deltas at risk. *Sustainability Science*, 3, 23–32.
- 48 **Syvitski, J.P.M., A.J. Kettner, I. Overeem, E.W.H. Hutton, M.T. Hannon, G.R. Brakenridge, J. Day, C. Vörösmarty,**
49 **Y. Saito, L. Giosan, and R.J. Nicholls,** 2009: Sinking deltas due to human activities. *Nature Geoscience*, 2, 681–
50 686.
- 51 **Tabet, L. and L. Fanning** 2012: Integrated coastal zone management under authoritarian rule: An evaluation
52 framework of coastal governance in Egypt. *Ocean and Coastal Management*, 61: 1-9.
- 53 **Tamura, T., K. Horaguchi, Y. Saito, L.N. Van, M. Tateishi, K.O.T., THi, F., Nanayama and K. Watanabe** (2010):
54 Monsoon-influenced variations in morphology and sediment of mesotidal beach on the Mekong River delta

- 1 coasta. *Geomorphology*, 116(1-2). 11-23.
- 2 **Tebaldi**, C., Strauss, B., and Zervas, C., 2012, Modeling sea level rise impacts on storm surges along US coasts,
3 *Environ. Res. Lett.*, doi:10.1088/1748-9326/7/1/014032
- 4 **Teneva** L., Karnauskas M., Logan C. A., Bianucci L., Currie J. C. and Kleypas J. A., 2011: Predicting coral
5 bleaching hotspots: the role of regional variability in thermal stress and potential adaptation rates. *Coral Reefs*
6 31:1-12.
- 7 **Terry**, J.P. and A.C. Falkland 2010: Responses of atoll freshwater lenses to storm-surge overwash in the Northern
8 Cook Islands. *Hydrology Journal*, 18, 749-759
- 9 Thinesh, T., G. Mathews, J. K. P. Edward, Coral disease prevalence in Mandapam group of islands, Gulf of Mannar,
10 Southeastern India *Indian J. Mar. Sci.* **38**, 444-450 (Dec, 2009).
- 11 Thomas, L. P., D. R. Moore, R. C. Work, Effects of hurricane Donna on the turtle grass beds of Biscayne Bay,
12 Florida. *Bull Mar Sci Gulf and Caribbean* **11**, 191-197 (1961).
- 13 **Thomsen** J. and Melzner F., 2010: Moderate seawater acidification does not elicit long-term metabolic depression
14 in the blue mussel *Mytilus edulis*. *Marine Biology* 157:2667-2676.
- 15 **Thomsen** J., Gutowska M. A., Saphörster J., Heinemann A., Trübenbach K., Fietzke J., Hiebenthal C., Eisenhauer
16 A., Körtzinger A., Wahl M. and Melzner F., 2010. Calcifying invertebrates succeed in a naturally CO2 enriched
17 coastal habitat but are threatened by high levels of future acidification. *Biogeosciences* 7:3879-3891.
- 18 Thresher, R. E., G. P. Harris, J. S. Gunn, L. A. Clementson, Planktonic production pulses and episodic settlement of
19 a temperate marine fish. *Nature* **341**, 641-642 (1989).
- 20 **Tobey**, J., P. Rubinoff, D. Robadue, G. Ricci, R. Volk, J. Furlow, and G. Anderson. 2010: Practicing Coastal
21 Adaptation to Climate Change: Lessons from Integrated Coastal Management. *Coastal Management* 38 (3):317-
22 335.
- 23 **Tol**, R.S.J., 2007: The double trade-off between adaptation and mitigation for sea level rise: An application of
24 FUND. *Mitigation and Adaptation Strategies for Global Change*, **12(5)**, 741-753.
- 25 **Tol**, R.S.J., R.J.T. Klein, and R.J. Nicholls, 2008: Towards successful adaptation to sea-level rise along europe's
26 coasts. *Journal of Coastal Research* **24(2)**, 432-442.
- 27 **Trenhaile**, A.S., 2011: Predicting the response of hard and soft rock coasts to changes in sea level and wave
28 height. *Climatic Change*, **109(3-4)**, 599-615.
- 29 **Tribbia**, J. and S.C. Moser, 2008: More than information: What coastal managers need to plan for climate
30 change. *Environmental Science and Policy* **11(4)**, 315-328.
- 31 **Turner**, R. K., D. Burgess, D. Hadley, E. Coombes, and N. Jackson, 2007: A cost-benefit appraisal of coastal
32 managed realignment policy. *Global Environmental Change* 17 397-407.
- 33 **UNCTAD** 2008: Maritime Transport and the Climate Change Challenge. UNCTAD Secretariat Note
34 TD/B/C.I/MEM.1/2. 9 Dec 2008, 17 pp.
- 35 UNEP 2009
- 36 **UNFCCC** 2010a: Potential costs and benefits of adaptation options: A review of existing literature. Technical paper.
37 In Potential costs and benefits of adaptation options: A review of existing literature. Technical paper. Geneva:
38 United Nations Framework Convention on Climate Change
- 39 **UNFCCC** 2010b: Synthesis report on efforts undertaken to assess the costs and benefits of adaptation options, and
40 views on lessons learned, good practices, gaps and needs. In Synthesis report on efforts undertaken to assess the
41 costs and benefits of adaptation options, and views on lessons learned, good practices, gaps and needs. Geneva:
42 United Nations Framework Convention on Climate Change.
- 43 **UNISDR**, 2011: Global Assessment Report on Disaster Risk Reduction: Revealing Risk, Redefining Development.
44 United Nations International Strategy for Disaster Reduction Secretariat, Geneva, Switzerland, Information
45 Press, Oxford, UK, 178 pp.
- 46 **US Department of Energy** (2006): Energy Demands on Water Resources. Report to Congress on the
47 Interdependency of Energy and Water. U.S. Department of Energy, Sandia National Laboratory. Available at:
48 http://www.sandia.gov/energy-water/congress_report.htm
- 49 **Van Aalst**, M. K., Cannon, T. and Burton, I. 2008: "Community level adaptation to climate change: The potential
50 role of participatory community risk assessment." *Global Environmental Change* 18(1): 165-179.
- 51 **Van Baars**, S. and I.M. Van Kempen, 2009: The Causes and Mechanisms of Historical Dike Failures in the
52 Netherlands. E-Water Official Publication of the European Water Association, Hennem, Germany, 14 pp.
- 53 **Van der Wal**, D. and K. Pye, 2004: Patterns, rates and possible causes of saltmarsh erosion in the greater thames
54 area (UK). *Geomorphology*, **61(3-4)**, 373-391.

- 1 **Van Kleef**, E., H. Bambrick, and S. Hales 2010: The geographic distribution of dengue fever and the potential
2 influence of global climate change. *TropIKA.net* <http://journal.tropika.net>.
- 3 **Van Koningsveld**, M., Mulder, J. P. M., Stive, M. J. F., VanDerValk, L. and VanDerWeck, A. W. 2008: "Living
4 with Sea-Level Rise and Climate Change: A Case Study of the Netherlands." *Journal of Coastal Research*: 367-
5 379.
- 6 VanWoesik, R., L. M. DeVantier, J. S. Glazebrook, Effects of Cyclone 'Joy' on nearshore coral communities of the
7 Great Barrier Reef *Mar. Ecol.-Prog. Ser.* **128**, 261-270 (Nov, 1995).
- 8 **Vaquer-Sunyer** R. and Duarte C. M., 2011: Temperature effects on oxygen thresholds for hypoxia in marine
9 benthic organisms. *Global Change Biology* 17:1788-1797.
- 10 Venrick, E. L., J. A. McGowan, D. R. Cayan, T. L. Hayward, Climate and Chlorophyll a: Long-Term Trends in the
11 Central North Pacific Ocean *Science* **238**, 70-72 (1987).
- 12 **Vermaat** J., W. Salomons, L. Bouwer, K. Turner, R. Nicholls, and R. Klein, 2005: Climate change and coastal
13 management on Europe's coast. In *Managing European Coasts*, Eds. R. Allan, U. Förstner and W. Salomons
14 199-226: Springer Berlin Heidelberg.
- 15 **Vignola**, R., B. Locatelli, C. Martinez, and P. Imbach. 2009. Ecosystem-based adaptation to climate change: what
16 role for policy-makers, society and scientists? *Mitigation and Adaptation Strategies for Global Change* 14
17 (8):691-696.
- 18 **Vinchon**, C., S. Aubie, Y. Balouin, L. Closset, M. Garcin, D. Idier, and C. Mallet, 2009: Anticipate response of
19 climate change on coastal risks at regional scale in Aquitaine and Languedoc Roussillon (France). *Ocean and*
20 *Coastal Management*, 52(1), 47-56
- 21 **Waldbusser** G. G., Bergschneider H. and Green M. A., 2010. Size-dependent pH effect on calcification in post-
22 larval hard clam *Mercenaria* spp. *Marine Ecology Progress Series* 417:171-182.
- 23 Walker, D. I., R. I. T. Prince, Distribution and biogeography of seagrass species on the northwest coast of Australia
24 *Aquat. Bot.* **29**, 19-32 (1987).
- 25 Walker, D. I., The effect of sea temperature on seagrasses and algae on the Western Australian coastline *Journal of*
26 *the Royal Society of Western Australia* **74**, 71 (1991).
- 27 **Wang**, X. L. L., V. R. Swail, F. W. Zwiers, X. B. Zhang, and Y. Feng, 2009: Detection of external influence on
28 trends of atmospheric storminess and northern oceans wave heights. *Climate Dynamics*, **32**, 189-203.
- 29 **Ward**, P.J., M.A. Marfai, F. Yulianto, D.R. Hizbaron, and J.C.J.H. Aerts 2011: Coastal inundation and damage
30 exposure estimation: a case study for Jakarta. *Natural Hazards*, 56, 899-916.
- 31 **Warren**, R. 2011: The role of interactions in a world implementing adaptation and mitigation solutions to climate
32 change. *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences*
33 369: 217-241.
- 34 **Wassmann**, R., S.V.K. Jagadish, K. Sumfleth, H. Pathak, G.Howell, A. Ismail, R. Serraj, E. Redona, R.K. Singh,
35 and S. Heur 2009: Regional vulnerability of climate change impacts on Asian rice production and scope for
36 adaptation. *Advances in Agronomy*, 102, 91-133.
- 37 Watanuki, Y., M. Ito, T. Deguchi, S. Minobe, Climate-forced seasonal mismatch between the hatching of rhinoceros
38 auklets and the availability of anchovy *Mar. Ecol.-Prog. Ser.* **393**, 259-271 (2009).
- 39 **Waycott**, M., C.M. Duarte, T.J.B. Carruthers, R.J. Orth, W.C. Dennison, S. Olyarnik, A. Calladine, J.W.
40 Fourqurean, J.K.L. Heck, A.R. Hughes, G.A. Kendrick, W.J. Kenworthy, F.T. Short, and S.L. Williams, 2009:
41 Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National*
42 *Academy of Sciences of the United States of America*, **106(30)**, 12377-12381.
- 43 Webb and Kench, 2010
- 44 **Webster** I. T. and Harris G. P., 2004: Anthropogenic impacts on the ecosystems of coastal lagoons: modelling
45 fundamental biogeochemical processes and management implications. *Marine and Freshwater Research* 55:67-
46 78.
- 47 Weishampel, J. F., D. A. Bagley, L. M. Ehrhart, Earlier nesting by loggerhead sea turtles following sea surface
48 warming. *Global Change Biology* **10**, 1424-1427 (2004).
- 49 Welsford, D. C., J. M. Lyle, Redbait (*emmelichthys nitidus*): a synopsis of fishery and biological data. *Tasmanian*
50 *Aquaculture and Fisheries Institute Technical Report Series* **20**, i (2003).
- 51 **Werner**, A.D. 2010: A review of seawater intrusion and its management in Australia. *Hydrogeology Journal* 18:
52 281-285
- 53 **White**, I. and Falkland, T, 2010: Management of freshwater lenses on small Pacific islands *Hydrogeology Journal*
54 18: 227-246

- 1 **White, I., T. Falkland, P. Perez, A. Dray, T. Metutera, E. Metia, and M. Overmars 2007:** Challenges in freshwater
2 management in low coral atolls. *Jour of Cleaner Production*, 15, 1522-1528.
- 3 Whitfield and Elliot, 2012
- 4 **Whiting, G. J., and J. P. Chanton, 2001:** Greenhouse carbon balance of wetlands: methane emission versus carbon
5 sequestration. *Tellus B* 53 (5):521-528.
- 6 **Whittock, A.,** The impact of sea level rise on a major mediterranean loggerhead sea turtle nesting site: Zakynthos
7 Island, Greece. *Testudo* 7, 49 (2009).
- 8 **Wilby, R.L. 2007:** A review of climate change on the built environment. *Built Environment*. 33: 31-45.
- 9 **Wild C., Hoegh-Guldberg O., Naumann M. S., Florencia Colombo-Pallotta M., Ateweberhan M., Fitt W. K.,**
10 **Iglesias-Prieto R., Palmer C., Bythell J. C., Ortiz J.-C., Loya Y. & van Woesik R., 2011.** Climate change
11 impedes scleractinian corals as primary reef ecosystem engineers. *Marine and Freshwater Research* 62:205-215.
- 12 **Wisser D., Fekete B. M., Vorosmarty C. J. and Schumann A. H., 2010:** Reconstructing 20th century global
13 hydrography: a contribution to the Global Terrestrial Network- Hydrology (GTN-H). *Hydrology and Earth*
14 *System Sciences* 14:1-24.
- 15 **Woodroffe, C. D.,** The impact of sea-level rise on mangrove shorelines *Prog. Phys. Geogr.* 14, 483-520 (Dec, 1990).
- 16 **Woodroffe, C.D., 2008:** Reef-island topography and the vulnerability of atolls to sea-level rise. *Global and*
17 *Planetary Change*, 62(1-2), 77-96.
- 18 **Woodroffe, C.D., Nicholls, R.J., Saito, Y., Chen, Z., Goodbred, S.L., 2006:** Landscape variability and the response
19 of Asian megadeltas to environmental change. In Harvey, N. (ed.), *Global Change and Integrated Coastal*
20 *Management: the Asia-Pacific Region. Coastal Systems and Continental Margins, Vol. 10.* Springer, pp. 277–
21 314.
- 22 **Woodward, R.T. and Y. Wui, 2001:** The economic value of wetland services: A meta-analysis. *Ecological*
23 *Economics*, 37(2), 257-270.
- 24 **Woollings, T., Gregory, J. M., Pinto, J. G., Reyers, M. and Brayshaw, D. J., 2012,** Response of the North Atlantic
25 storm track to climate change shaped by ocean-atmosphere coupling, *Nature Geoscience*, DOI:
26 10.1038/NNGEO1438
- 27 **Wootton J. T., Pfister C. A. and Forester J. D., 2008:** Dynamic patterns and ecological impacts of declining ocean
28 pH in a high-resolution multi-year dataset. *Proceedings of the National Academy of Science U.S.A.* 105:18848-
29 18853.
- 30 **WRI, World Resource Institute (2007).** Bangladesh: Cyclone Preparedness Program.
31 http://maindb.unfccc.int/public/adaptation/adaptation_casestudy.pl?id_project=37
- 32 **WTO (World Tourism Organization) 2007:** Climate Change and Tourism – Responding to Global Challenges:
33 Summary. Madrid, 24 pp.
- 34 **Wu, H.Y., Zou, D.H., Gao, K.S., 2008.** Impacts of increased atmospheric CO2 concentration on photosynthesis and
35 growth of micro- and macro-algae. *Science in China Series C: Life Sciences*. 51: 1144-1150
- 36 **Wynn, R. B., S. A. Josey, A. P. Martin, D. G. Johns, P. Yesou,** Climate-driven range expansion of a critically
37 endangered top predator in northeast Atlantic waters *Biology Letters* 3, 529-532 (Oct, 2007).
- 38 **Xia, J., R.A. Falconer, B. Lin, and G. Tan, 2011 }** Estimation of future coastal flood risk in the severn estuary due to
39 a barrage. *Journal of Flood Risk Management*, 4(3), 247-259.
- 40 **Yamane, L., S. E. Gilman,** Opposite responses by an intertidal predator to increasing aquatic and aerial temperatures
41 *Marine Ecology Progress Series* 393, 27-36 (2009).
- 42 **Yamano H., Sugihara K. and Nomura K., 2011:** Rapid poleward range expansion of tropical reef corals in response
43 to rising sea surface temperatures. *Geophysical Research Letters* 38, L04501. doi:10.1029/2010GL046474.
- 44 **Yang, S.L., M. Li, S.B. Dai, Z. Liu, J. Zhang, and P.X. Ding, 2006:** Drastic decrease in sediment supply from the
45 Yangtze River and its challenge to coastal wetland management. *Geophysical Research Letters*, 33, L06408.
- 46 **Yang, S.L., Milliman, J.D., Li, P., and Xu, K., 2011:** 50,000 dams later: Erosion of the Yangtze River and its delta.
47 *Global and Planetary Change*, 75, 14-20.
- 48 **Yang, X. Y., R. X. Huang, and D. X. Wang, 2007:** Decadal changes of wind stress over the Southern Ocean
49 associated with Antarctic ozone depletion. *Journal of Climate*, 20, 3395-3410.
- 50 **Yasuhara, K., S. Murakami, N. Mimura, H. Komine, and J. Recio 2007:** Influence of global warming on coastal
51 infrastructural instability. *Sustainability Science*, 2, 13-25.
- 52 **Yin, J, Yin, Z, Wang, J, Xu S. 2012:** National assessment of coastal vulnerability to sea-level rise for the Chinese
53 coast. *Journal Coastal Conservation* 16(1):123–133

- 1 **Yin**, K. D., P. J. Harrison, R. H. Goldblatt, R. J. Beamish, 1996: Spring bloom in the central Strait of Georgia:
2 interactions of river discharge, winds and grazing. *Mar. Ecol.-Prog. Ser.* **138**, 255-263.
- 3 **Yohe** G., Knee K. and Kirshen P. 2011. On the economics of coastal adaptation solutions in an uncertain world,
4 *Climatic Change*, 106 : 71-92.
- 5 **Yoo**, G., J.H. Hwang, and C. Choi, 2011}: Development and application of a methodology for vulnerability
6 assessment of climate change in coastal cities. *Ocean and Coastal Management*, **54(7)**, 524-534.
- 7 **Young**, I.R., Zieger, S., Babanin, A. V. 2011: Global trends in wind speed and wave height. VOL 332 Science, Vol.
8 332, 451-455.
- 9 **Zacher**, K., R. Rautenberger, D. Hanelt, A. Wulff, C. Wiencke, 2009: The abiotic environment of polar marine
10 benthic algae. *Botanica Marina* **52**, 483-490.
- 11 **Zampatti**, B.P., C.M. Bice, and P.R. Jennings, 2010: Temporal variability in fish assemblage structure and
12 recruitment in a freshwater-deprived estuary: The coorong, australia. *Marine and Freshwater Research*, **61(11)**,
13 1298-1312.
- 14 **Zeitlin** H.L., I. Meliane, S. Davidson, T. Sandwith and J. Hoekstra 2012: Ecosystem-based adaptation in marine and
15 coastal ecosystems. *Environmental Sciences*, 25: 1-10.
- 16 **Zhang** J., Gilbert D., Gooday A. J., Levin L., Naqvi S. W. A., Middelburg J. J., Scranton M., Ekau W., Peña A.,
17 Dewitte B., Oguz T., Monteiro P. M. S., Urban E., Rabalais N. N., Ittekkot V., Kemp W. M., Ulloa O., Elmgren
18 R., Escobar-Briones E. and Van der Plas A. K., 2010. Natural and human-induced hypoxia and consequences
19 for coastal areas: synthesis and future development. *Biogeosciences* 7:1443-1467.
- 20 **Zhang**, Y., Bi, P., Hiller, J.E., Sun, Y., and Ryan, P., 2007: Climate variations and bacillary dysentery in northern
21 and southern cities of China *Journal of Infection* 55, 194e200
- 22 **Zimmerman**, R. and Faris, C., 2010: Infrastructure impacts and adaptation challenges. In: Rosenzweig, C., Solecki,
23 W. (Eds) *New York City Panel on Climate Change, 2010: Climate change adaptation in new your city: building*
24 *a risk management response*. Prepared for use by the New York City Climate Change Adaptation Task Force.
25 *Annals of the New York Academy of Science* 2010. New York, NY, pp 63-85.
- 26 **Zimmerman**, R.C., R.D. Smith, and R.S. Alberte, 1989: Thermal acclimation and whole plant carbon balance
27 in *zostera marina* L. (eelgrass) in San Francisco Bay. *Journal of Experimental Marine Biology and*
28 *Ecology*, **130**, 93-109.
- 29 **Zue** Y., B. Huang, Z. -Z. Hu, A. Kumar, C. Wen, D. Behringer, and S. Nadiga, 2010: An assessment of oceanic
30 variability in the NCEP climate forecast system reanalysis. *Climate Dynamics*, 10.1007/s00382-010-0954-4,1-
31 29.
- 32

Table 5-1: Survey of literature on the impacts of anthropogenic climate change on marine ecosystems (adapted and updated from Hoegh-Guldberg and Bruno, 2010). [To be revised and confined to coastal ecosystems.]

Climate system change	Organism/ ecosystem	Expected impact	Observed change	References
Increasing temperature	Seagrass	Seasonal and permanent loss of seagrass biomass with increased frequency and intensity of extreme temperatures	Increased temperatures results in a reduction in the above-ground biomass of seagrass and the disruption of the photosystem. Mass die-offs and ecosystem loss in areas exposed to prolonged extreme temperatures	Borum et al., 2005; Campbell et al 2006; Greve et al, 2003; Mayot et al, 2005; Moore and Jarvis, 2008; Najjar et al, 2010; Orth et al, 2006; Seddon and Cheshire; 2001; Seddon et al, 2000; Short and Neckles, 1999.
		Shift in community structure	Warm-water species proliferate, dominating communities in areas of low-level warming	Boudouresque et al, 2009; Ehlers et al, 2008; Francour et al, 1994; McMillan, 1984; Peirano et al, 2005; Walker, 1991
	Mangroves	Changes in species distribution and loss of habitat	Increased salinity due to higher evaporation leads to mortality and redistribution of species and reduced species richness due to variable salinity tolerance levels. Prolonged periods of extreme salinity may result in the formation of salt pan systems	Ball, 1998; Ball and Pidsley, 1995; Bertness and Pennings, 2000
		Rocky shores	Poleward shift in species ranges	The range and abundance of warm-water species are increasing, whilst those of coldwater species are diminishing
	Kelp communities		Zonation patterns influenced by both air and sea temperatures	Reduced recruitment of fucoids and intertidal invertebrates in the littoral zone due to rising temperatures causing desiccation of propagules and suppressing growth leaving new recruits more susceptible to grazers
			Decline of kelp ecosystems with rising sea surface	Range and distribution of kelps is diminishing with rising temperatures due to requirements of sporophytes. Species

	temperature	living close to their physiological limits will be likely to recede to higher latitudes and cooler waters.	Deysher and Dean, 1986; Steneck et al, 2002
Phyto-plankton	Changes in distribution and frequency of harmful algal blooms	Increased frequency of bloom events associated with increasing sea surface temperatures.	Peperzak, 2003; Peperzak, 2005
	Altered growth rates, species dependent	Some species growth rates increased with temperature	Fu et al, 2007; Fu et al, 2008
	Poleward shift in species ranges	Warm water species are increasing their distribution towards the poles as cold water warms	Beaugrand and Reid, 2008; Edwards, 2004;
	Altered abundance	A greater increase in abundance in cooler waters experiencing warming compared to warmer waters experiencing warming	Richardson and Schoeman, 2004
	Earlier appearance	Phytoplankton appearing earlier in summer in temperate regions	Edwards and Richardson, 2004
Zooplankton	Poleward shift in species ranges	A shift in community assemblages and biogeographical range, extending polewards with increasing sea surface temperatures	Beaugrand et al, 2002;Beaugrand et al, 2009; Parmesan and Yohe, 2003; Root et al, 2003.
	Alteration of phenology	Zooplankton communities appear earlier with warming sea surface temperatures	Edwards and Richardson, 2004; Parmesan and Yohe, 2003
	Altered abundance	Increase in abundance with warming water	Richardson and Schoeman, 2004; Aoyama et al, 2008
Coral reefs	Increased frequency and severity of coral bleaching with changing sea surface temperature	Severe bleaching events occurring globally with associated coral mortality	Hoegh-Guldberg, 1999; Knowlton, 2001; Miller et al, 2009; Mumby et al, 2001, Prada et al, 2010
	Increased occurrence of diseases	Frequency and severity of coral diseases increasing	Croquer and Weil, 2009; Mydlarz et al, 2009; Sokolow, 2009; Thinesh et al, 2009; Baker et al, 2008;

	Loss of coral reef species due to coral bleaching and mortality	Loss of coral reef fish, crustacea and other invertebrate diversity and abundance with loss of live coral habitat due to rising temperatures	Baker et al, 2008; Bruce and Biol, 1976; Castro, 1978; Castro, 1988; Feary et al, 2007
Seabirds	Poleward shift in species ranges and a shift in abundance toward species tolerant of warmer waters	Seabirds of Western Australia are becoming more abundant and extending their range polewards with changes in prey distribution with rising sea surface temperatures	Bancroft et al, 2004; Dunlop et al, 2001, Dunlop and Wooller, 1986; Smithers et al, 2003; Wynn et al, 2007
	Birds migrating earlier in temperate and subtropical regions	Alteration of breeding date with changing temperature, favouring early breeding and altered selection patterns	Moe et al, 2009; Moller et al, 2006
	Altered breeding seasons, affecting nesting and laying times	Extended breeding seasons in the temperate and tropical regions with earlier nesting and laying times	Dunlop and Wooller, 1986; Chambers, 2004, Nevoux et al, 2010
	Breeding success affected by climate change and prey Availability	Temperature and associated changes in prey availability and match-mismatch of breeding affect population success	Bustnes et al, 2010; Plestan et al, 2009; Watanuki et al, 2009
	Alteration of coastal habitats affect nesting bird populations	Penguin populations benefit from less snow and ice allowing better nesting and more abundant prey species improving breeding success	Huang et al, 2009
Marine turtles	Poleward shift in species foraging ranges	Temperature change has implications on migratory patterns, forcing a poleward shift in populations	Chaloupka et al, 2008; McMahon and Hays, 2006
	Change in the sex ratios	Changes in temperatures affect the sex ratio with rising temperatures favouring female populations	Booth and Freeman, 2006; Fuentes et al, 2010; Godley et al, 2001
	Changes in breeding	Warmer foraging and nesting grounds affect the timing of breeding, clutch number and nesting season length	Mazaris et al, 2009; Pike, 2009; Schofield et al, 2009; Weishampel et al, 2004
Marine mammals	Change in distribution range of Cetacea	Poleward migration of species causing a reduction in the range of cold water species and extension of warm water species resulting in changes in community structure	Azzellino et al, 2008; Gambaiani et al, 2009; MacLeod et al, 2005
Polar Ice	Ice thinning and loss results	Prolonged periods of ice loss, or thin ice affects the growth	Montes-Hugo et al, 2009; Zacker et

	Habitats	in greater UV penetration to the marine system	and distribution of benthic and pelagic microalgae and cyanobacteria altering productivity	al, 2009
		Changes to seasonal ice loss patterns	Changes to the seasonal ice break events alters the marine eukaryotic communities and system function	Piquet et al, 2008
		Loss of ice will change the distribution of ice-dependent macrofauna	Changes in migration patterns, adaptation to changing habitats and possible declining in population depending on level of dependency	Moore and Huntington, 2008
	Demersal and pelagic fish	Species range alters with warming	Poleward shift in species ranges and a shift in abundance toward species tolerant of warmer waters	Beare et al, 2004, Byrkjedal et al, 2004, Perry et al, 2005; Rose, 2005; Welsford and Lyle, 2003; Hiddink and Hofstede, 2008; Last et al, 2011; Ling et al, 2009
		Migration dates altered with warming	Earlier dates of mean migration and spawning in temperate and subtropical species	Sims et al, 2004
Wind strength change	Phyto-plankton and zooplankton	Alteration of productivity with wind-driven mixing of surface waters	Increased productivity where wind mixing is enhanced and a reduction where wind strength is declined	Montes-Hugo et al, 2009; Buranapratheprat et al, 2008; Harris et al, 1991; Polovina et al, 1994
		Changes in community structure with surface mixing	Alteration of the surface stratification with wind-driven upwelling can cause alterations in community structure and bloom formation	Nakane et al, 2008; Yin et al, 1996
	Coastal fish	Abundance of fish linked to wind strength	Increased wind-driven upwelling and mixing results in greater recruitment due to areas of higher productivity	Harris et al, 1991; Polovina et al, 1994; Criales et al, 2002; Thresher et al, 1989
	Seabirds	Alteration of breeding success with changing wind intensity and patterns	Prolonged periods of strong winds causes a reduction in the breeding success of seabirds	Devney et al, 2009; King et al, 1992

Alteration of currents	Seagrass	Changes in distribution of species with changing currents	Loss of cold-water species and appearance of topical species further poleward correlated with changes in warm water currents	Walker and Prince, 1987
	Mangroves	Breakdown in control of latitudinal distribution through propagule current translocation	Changes to currents responsible for propagule distribution results in the redistribution of mangroves	Delange and Delange, 1994
		Changes in local distribution patterns with changing sediment transport patterns	Changes in current-driven sediment distribution affects growth rates and success of plant	Ellison and Farnsworth, 1996
	Kelp communities	Local extinction of cold-water species with changes in currents and/or the appearance of warm-water species	Alteration of larval supply changes the distribution of species and success in altered thermal conditions	Johnson et al, 2005; Ling et al, 2009
	Rocky shores	Poleward shift of warm water species	Tropical species appearing in temperate latitudes due to changes in distribution of larvae	Griffiths, 2003
	Phyto-plankton and zooplankton	Change in distribution and occurrence of plankton communities with an extension polewards of warm-water species	Warm nutrient rich waters resulting form changes in current trajectories results in plankton bloom events	Blackburn and Cresswell, 1993; Oke and Middleton, 2001
Decline in mixed layer depth/ increasing stratification	Seabirds	Increased mortality and reduced reproductive success	Reductions in surface water prey availability due to strengthened stratification and reduced mixed layer leads to mortality and reduced reproductive success	Richardson and Schoeman, 2004; Smithers et al, 2003; Wynn et al, 2007
	Pelagic fish	Abundance and distribution	Stratification and associated plankton community changes	Richardson and Schoeman, 2004;

		changes due to thermal stratification of upper ocean	alters food supply for fishes, altering community structure and distribution	Hsieh et al, 2009
	Phyto-plankton and zooplankton	Changes in distribution and abundance due to altered stratification zones	Vertical stratification resulting from changes in sea surface temperatures strengthens existing thermoclines in warmer stratified waters and encourage the development of their formation in cooler turbulent waters creating suitable habitat for zooplankton	Richardson and Schoeman, 2004; Hsieh et al, 2009
		Decline in phytoplankton abundance	As the mixed surface layer diminishes phytoplankton productivity decreases	Polovina et al, 1994; Polovina et al, 1995; Venrick et al, 1987
Increasing intensity of storms/ greater inundation events from shifting rainfall	Seagrass	Physical destruction of seagrass beds	Storm-driven currents scour the benthos uprooting large areas of seagrass and removing from the site	Orth et al, 2006; Preen et al, 1995; Rodrigues et al, 1994; Thomas et al, 1961
		Changes in sedimentation regimes cause mortality	Sediment deposition caused by storm activity and increased rainfall runoff smothers seagrass	Preen et al, 1995; Rodrigues et al, 1994;
		Change in community composition as water clarity is changed	Alteration to light conditions due to reduced water quality resulting from increased sediment load results in a change of community shifting towards species adapted to low light levels	Rodrigues et al, 1994; Hale et al, 2004; Orth and Moore, 1983
	Mangroves	Change in community abundance associated with increased rainfall events	Mangrove community distribution increase due to altered salinity, nutrient and sediment loading	Gilman et al, 2008; Harty, 2004; Rogers et al, 2006; Saintilan and Williams, 1999
		Reproductive success and growth influenced by storm activity	Prolonged periods of flooding may cause the mortality and impeded propagation of juvenile plants	Gilman et al, 2008
	Rocky shore	Increased wave energy alters community structure	Storm-driven wave damage change in species zonation patterns	Helmuth et al 2006; Barry et al, 1995
Increased storm frequency affects community structure and function group		Furoid species will be lost and associated invertebrates, allowing those species that can withstand high energy environments, such as mussels and barnacles, to dominate	Kendall et al, 2004	

	prevalence		
	Increased freshwater inputs alters zonation	Changes in species zonation driven by changes in salinity due to extreme rain events	Garza and Robles, 2010
Kelp communities	Change in community structure	Switch from canopy forming macroalgae to predominantly turf-algae due to physical wave damage and increased eutrophication from land run-off	Cole et al, 2001; Gorgula and Connell, 2004; Graham, 1997; Graham et al, 1997; Steneck et al, 2002
Coral reefs	Mass mortality due to physical damage	Extreme storm events cause physical destruction and mortality of corals with increased frequency preventing recovery leaving reef susceptible to less intense events	Gardner et al, 2005; Alvarez-Filip and Gil, 2006; Alvarez-Filip et al, 2009; Connell, 1997; Guillemot et al, 2010; Hughes and Connell, 1999; Knowlton et al, 1981; Porter and Meier, 1992
	Mass bleaching and mortality due to associated large freshwater flood events	Extended periods of extreme freshwater input from land run-off causes mass bleaching and potential mortality	Alongi and McKinnon, 2005; Alongi and Robertson, 1995; Goreau, 1964; Jokiel et al, 1993; VanWoesik et al, 1995
	Changes in reef community structure and composition	Differentiation in sensitivity to freshwater and mechanical stress, and differences in recover rates causes a shift in community composition and reduced diversity	Jokiel et al, 1993; VanWoesik et al, 1995; Adjeroud et al, 2009; Bythell et al, 2000; DeVantier et al, 2006
Phyto-plankton and zooplankton	Nutrient enrichment of surface waters due to terrestrial run-off	Increased nutrient state causes a change in community structure and dynamics causing a shift from heterotrophy to autotrophy	Carlsson et al, 1995; De Carlo et al, 2007; Goffart et al, 2002; Guadayol et al, 2009; Hoover et al, 2006
	Storm-forced upwelling of nutrient rich waters	Nutrient rich water promotes phytoplankton growth	Acker et al, 2009
Marine turtles and mammals	Increased mortality and reduced breeding success	Severe storm events cause mortality of terrestrial-obligate mammals and turtles including loss of turtle clutches	Dodd and Dreslik, 2008; Edmiston et al, 2008; Limpus and Reed, 1985; Pemberton and Gales, 2004; Pike and Stiner, 2007

	Seabirds	Increased feeding	Increased plankton abundance drives foraging success and breeding population dynamics	Devney et al, 2009
		Restriction and alteration of foraging and migration	Storm events prevent birds from travelling usual routes and cause changes in flight patterns	Blomqvist and Peterz, 1984
Rising sea levels	Seagrass	Loss of seagrass habitat	Rising sea levels results in increased light attenuation forcing seagrass migration landwards to areas of shallower water	Orth et al, 2006; Abal and Dennison, 1996
		Reduction in growth rate and changes in community structure due to lower light levels	Change in community structure with species with lower light demands dominating deeper zones	Short and Neckles, 1999
	Mangrove	Loss of mangrove habitat	Increased frequency and severity of extreme sea levels may results in mortality where migration is impeded	Blasco et al, 1996; Ellison, 1993; Ellison and Stoddart, 1991; Woodroffe, 1990
		Changes in habitat distribution	Landward migration in response to slow sea-level rise allowing the maintenance of relative height	Gilman et al, 2008; Ellison, 1993; Alongi, 2008; Gilman et al, 2006; Madsen et al, 2007; Parkinson, 1989 Parkinson et al, 1994
	Seabirds	Loss of nesting and breeding habitat	Inundation of nesting habitats in low lying habitat areas by water will cause a reduction the potential habitat for populations	Galbraith et al, 2005; Ratcliffe et al, 2008; Smart and Gill, 2003; Straw et al, 2006;
	Marine turtles and mammals	Loss of nesting and breeding habitat	Inundation of turtle nesting habitats in low lying areas by water will cause a reduction the potential habitat for populations	Fish et al, 2005; Limpus and Heidrun, 2006; Mazaris et al, 2009a, b; Whittock, 2009
Coral reefs	Mortality and redistribution of communities	Distribution of corals will shift so as to maintain their relative sea-level while corals living at their physiological light limit will die if rate of sea-level change exceeds growth rate	Hoegh_Guldberg, 1999; Graus and Macintyre, 1998	

Table 5-2: Top ten nations with the largest populations and the highest proportions of population in low-lying coastal areas (Countries with fewer than 100,000 inhabitants are not included; 15 SIDs (Small Island Developing States) with total of 423,000 inhabitants are also excluded). Source: Bollman *et al.*, 2010.

Top ten nations classified by population in low-lying coastal regions			Top ten nations classified by proportion of population in low-lying coastal areas		
Nation	Population in low-lying coastal regions (10 ³)	% of population in low-lying coastal regions	Nation	Population in low-lying coastal regions (10 ³)	% of population in low-lying coastal regions
1. China	127,038	10 %	1. Maldives	291	100 %
2. India	63,341	6 %	2. Bahamas	267	88 %
3. Bangladesh	53,111	39 %	3. Bahrain	501	78 %
4. Indonesia	41,807	20 %	4. Suriname	325	78 %
5. Vietnam	41,439	53 %	5. Netherlands	9590	60 %
6. Japan	30,827	24 %	6. Macao	264	59 %
7. Egypt	24,411	36 %	7. Guyana	419	55 %
8. USA	23,279	8 %	8. Vietnam	41,439	53 %
9. Thailand	15,689	25 %	9. Djibouti	250	40 %
10. Philippines	15,122	20 %	10. Bangladesh	53,111	39 %

Table 5-3: Impacts of sea level rise, coastal floods, and storms on critical coastal infrastructure by sector.

Table 3 Impacts of sea level rise, coastal floods, and storms on critical coastal infrastructure by sector

Communications	Energy	Transportation	Water and waste
Higher average sea level			
<ul style="list-style-type: none"> Increased salt water encroachment and damage to low-lying communications infrastructure not built to withstand saltwater exposure Increased rates of coastal erosion and/or permanent inundation of low-lying areas, causing increased maintenance costs and shortened replacement cycles Tower destruction or loss of function 	<ul style="list-style-type: none"> Increased rates of coastal erosion and/or permanent inundation of low-lying areas, threatening coastal power plants Increased equipment damage from corrosive effects of salt water encroachment resulting in higher maintenance costs and shorter replacement cycles 	<ul style="list-style-type: none"> Increased salt water encroachment and damage to infrastructure not built to withstand saltwater exposure Increased rates of coastal erosion and/or permanent inundation of low-lying areas, resulting in increased maintenance costs and shorter replacement cycles Decreased clearance levels under bridges 	<ul style="list-style-type: none"> Increased salt water encroachment and damage to water and waste infrastructure not built to withstand saltwater exposure Increased release of pollution and contaminant runoff from sewer systems, treatment plants, brownfields and waste-storage facilities Permanent inundation of low-lying areas, wetlands, piers, and marine transfer stations Increased salt water infiltration into distribution systems transfer stations
More frequent and intense coastal flooding			
<ul style="list-style-type: none"> Increased need for emergency management actions with high demand on communications infrastructure Increased damage to communications equipment and infrastructure in low-lying areas 	<ul style="list-style-type: none"> Increased need for emergency management actions Exacerbated flooding of low-lying power plants and equipment, as well as structural damage to infrastructure due to wave action Increased use of energy to control floodwaters Increased number and duration of local outages due to flooded and corroded equipment 	<ul style="list-style-type: none"> Increased need for emergency management actions Exacerbated flooding of streets, subways, tunnel and bridge entrances, as well as structural damage to Decreased levels of service from infrastructure due to wave action flooded roadways; increased hours of delay from congestion during street-flooding episodes Increased energy use for pumping 	<ul style="list-style-type: none"> Increased need for emergency management actions Exacerbated street, basement and sewer flooding, leading to structural damage to infrastructure Episodic inundation of low-lying areas, wetlands, piers, and marine transfer stations

Sources: Horton and Rosenzweig (2010), Zimmerman and Faris (2010)

Climate Change (2011) 106-93-127

107

Table 5-4: Summary of Coastal Vulnerability Assessments conducted at national or regional scale. Studies that either costed (Adapt\$) or considered adaptation options (Adapt) are indicated. In studies that have considered adaptation: WOA=without adaptation; WA=with adaptation. CVI=Coastal Vulnerability Index; GIS=Geographic Information System, DIVA=Dynamic Interactive Vulnerability Assessment model. RSLR=relative sea level rise.

Location	Physical impacts considered	SLR scenario	Socio-economic scenario	Physical and ecosystem vulnerability	Human system vulnerability	Method and tools	Reference
Africa							
Africa	Erosion, flooding	0.64-1.26 m SLR by 2100, plus local subsidence /uplift	IMAGE model scenario		WOA: 16–27 million people flooded and US\$5-9 billion annual damage costs WA: Adaptation cuts people flooded by two orders of magnitude and damage costs by half, adaptation deficit US\$300 billion and annual costs US\$2-6 billion	DIVA (Adapt\$)	Hinkel et al., 2011
Ghana (east coast)	Erosion, submergence	1-5m SLR		Possible erosion of existing coastal buffer zones that separate open coast from coastal lagoons and inundation of coastal plains		GIS	Boateng, 2012b
Tanzania (Dar-es-Salaam)	Flood exposure	0.13-0.66 m by 2070, plus local subsidence /uplift, plus to 1-in-100 event	A1 with rapid urbanization		210,000 people and US\$10 billion exposed to 100-year coastal flood by 2070	GIS	Kebede and Nicholls, 2012
Asia							
China				Sediment supply to major delta regions has reduced by 50% since early 1980's and is driving coastal erosion			Cai et al., 2009
China	Flooding, erosion	Current rate of RSLR (average 6mm/yr)		Vulnerability to flooding and erosion of the 18,000 km coastline, is ranked very high-3%; high-29%; moderate- 58% and low-10%.		CVI	Yin et al., 2012
East Timor	Flooding,	0.31-0.54	A2 and B1	Wetland area reduction under	Expected number of people flooded	DIVA	McLeod

	erosion wetland loss, salt water intrusion	m by 2100, plus local subsidence / uplift		0.54m SLR is greatest for the Solomon Islands (68%), the Philippines (51%) and East Timor (50%). Composition of wetland areas in 2100 change to include more mangroves and less unvegetated wetland areas in 2100 compared to 2010. Increased salinity intrusion up major rivers 14-27% under A2 Increased salinity intrusion into land areas 7-12% under A2	annually relative national population (pop):	(Adapt\$)	et al., 2010
Indonesia					WOA 0.54% pop A2; WA 0.01% pop A2		
Malaysia					WOA 1.19% pop A2; WA 0.00% pop A2		
Papua New Guinea					WOA 0.46% pop A2; WA 0.00% pop A2		
Philippines					WOA 0.07% pop A2; WA 0.01% pop A2		
Solomon Islands					WOA 0.80% pop A2; WA 0.00% pop A2 WOA 0.27% pop A2; WA 0.00% pop A2		
India (Udupi coast)	Erosion, submergence	1-10m SLR		Erosion risk: of the 95 km of coastline, 59% assessed as very high; 7% high, 4% moderate, 30% low erosion risk. 372-42 km ² exposed to submergence from 1m SLR	Qualitative assessment of affected areas include residential and recreational areas, agricultural land and the natural ecosystem	CVI	Dwarakish et al., 2009
Vietnam	Submergence exposure	1-5m			Exposure is largest in Red River and Mekong deltas. Options identified to prolong the use of these areas into the future	GIS	Boateng, 2012a
Australasia							
Australia	Flood exposure, Erosion	1.1 m, combined with either 1-in-100 year flood event or mean high tide level		Exposure to tropical cyclone related hazards across northern half of continent, health of Great Barrier Reef will affect coastal resilience in NE. Southern coastline with extensive cliffs, large bays, estuaries, gulfs and flats vulnerable to SLR inundation and cliff instability if wave climate changes. Greater erosion along populous eastern coast due to SLR and storm changes.	Potential inundation of, 157,000–247,600 of the 711,000 existing residential buildings close to water. Nearly 39,000 buildings at risk from erosion due to SLR. Indigenous communities, including island based deemed particularly vulnerable due to their remoteness and location on low elevation land.	GIS coastal geomorphology model and Bruun Rule GIS DEM infrastructure database	Department of Climate Change, 2009
Australia (Victoria)	Flood exposure	0.8-1.4 m SLR by 2100, plus 1-in-		Across 9 coastal settlements considered, area exposed to 1-in-100 flood ranges from 153 to 408 km ² for 0-1.4 m SLR	Across 9 coastal settlements considered, land parcels exposed to 1-in-100 flood ranges from 2,362 to 47,102 for 0 to 1.4 m SLR	Hydrodynamic modelling, GIS	McInnes et al., 2011

		100 year flood event					
Europe							
Croatia	Submergence exposure	0.2-0.86 m SLR		A long narrow steep coastline with one major reclaimed alluvial plain will become increasingly vulnerable to inundation particularly for larger SLR projections. Erosion of pocket beaches will increase.	Sewage systems, agriculture and maritime transport and tourism affected by greater flood frequency of low-lying land and berths and piers.	Qualitative assessment	Baric et al., 2008
Denmark-Copenhagen	Flood exposure	0.0-1.25 m SLR			Copenhagen not highly vulnerable to coastal flooding due to existing flood protection. WOA: direct costs of 1-in-100 year event increase from €3-4.8 billion with 0.5m SLR	GIS (Adapt)	Hallegatte et al., 2011
Estonia	Submergence exposure	1 m SLR adjusted for regional uplift (i.e. 69–73 cm RSLR)		Observed beach erosion has resulted from increased storminess in the eastern Baltic Sea, combined with decline in winter sea-ice cover. Future land loss will impact major bird breeding grounds	Possible productivity benefit from longer growing season. Major towns are not threatened due to location inland and mitigating effects of uplift. Sandy beaches and emerging coastal tourism at risk.	Qualitative assessment	Kont et al., 2008
France	Flood exposure, erosion	0.88 m SLR + 5.8m surge (Atlantic) and 2.7 m surge (Mediterranean)		Atlantic coast with extensive dune systems is assessed as more resilient to rising sea levels over the coming century compared with Mediterranean coast where narrow dune systems are highly urbanized		GIS	Vinchon et al., 2009
Germany	Submergence exposure	1 m SLR		There is a high level of reliance on hard coastal protection against extreme sea level hazards which will increase ecological vulnerability over time	300,000 people exposed in the coastal cities and communities. Erosion and flooding risks US\$300 billion (based on 1995 values) of assets	GIS DEM, land-use, socioeconomic data	Sterr, 2008
Great	Erosion,	0.8 – 1m		Large parts of the coasts are	At the national scale, economic losses due	Qualitati	de la

Britain	submergence	SLR		presently sediment starved and eroding and this will continue	to erosion are expected to remain considerably smaller than flood losses	ve assessment	Vega-Leinert and Nicholls, 2008
Norway	Submergence	0.5 - 1m SLR		Nationally, low susceptibility to accelerated sea-level rise due to mainly steep and resistant coastlines	Extensive infrastructure on northern and western coastlines likely to be negatively affected by sea-level rise, and adaptation costs could be significant	Qualitative assessment	Aunan and Romstad, 2008
Poland	Submergence	0.3 – 1m SLR		Lagoons, river deltas and estuaries in the far east and west were considered most vulnerable		Qualitative assessment	Pruszek and Zawadzka, 2008
Portugal	Submergence	0.14-0.57 m by 2100 SLR		Estuaries and coastal lagoons are assessed as most vulnerable and already sediment starved coastal beaches will continue to erode		Qualitative assessment	Ferreira et al., 2008
Turkey	Submergence	1 m SLR			Without adaptation, impacts could cost 6% of current GNP. Adaptation/protection could cost 10% of current GNP.	Qualitative (Adapt)	Karaca and Nicholls, 2008
European Union	Erosion, flooding, salinity intrusion	0.35-0.45 m by 2100 SLR	A2 and B1		WOA: 0.2-0.8 million people flooded and US\$17 billion annual damage costs WA: adaptation cuts exposure by a factor of 100 and annual damage costs by a factor of 10.	DIVA (Adapt\$)	Hinkel et al., 2010
N. America							
NW territories (Canada)	Shoreline stability and population exposure	Temperature, wind patterns		Loss of sea ice and shorter winter season and more variable weather patterns lead to changes in coastal sea ice coverage. Effects on species health and numbers.	More hazardous travel conditions for traditional hunting practices. Loss of traditional knowledge, skills and values.	Qualitative (Adapt)	Pearce et al., 2010

Table 5-5: Global assessments of costs of sea-level rise.

Reference	Physical impacts	SLR scenario	Socio-economic scenario	Impact indicators	Without adaptation	With adaptation
Anthoff et al., 2010	Submergence, wetland loss	0.5-2.0m SLR by 2100	A1, A2, B1, B2	Net present value of total cost for 2005-2100 (pure rate of time preference 1%)	US\$ 800-3300 billion in 2100	US\$200-2200 billion in 2100
Nicholls et al., 2011	Submergence, erosion	0.5-2.0 m SLR by 2100	A1FI	Cumulative number of people displaced due to land loss to submergence and erosion	72-187 million people during 21 st century	0.04-0.3 million people during 21 st century
				Annual adaptation cost	N/a	US\$ 25-270 billion/yr
Hinkel et al., 2012	Flooding	0.6-1.3m by 2100	UN medium fertility	Annual expected number of people flooded	170-260 million people/yr in 2100	Two orders of magnitude smaller than w/o adaptation
				Annual total cost (including dike upgrade, dike maintenance and residual damage cost)	US\$ 160-300 billion/yr in 2100	US\$ 30-90 billion/yr in 2100
				Annual total cost relative to GDP	0.05% -0.09% of global GDP in 2100	0.01%-0.03% of global GDP in 2100

Table 5-6: Approaches to integrative, adaptive coastal management.

Characteristics	Traditional Coastal Zone Management	Integrated Coastal Zone Management	Disaster Risk Reduction	Ecosystem-based Adaptation	Community Based Adaptation
Focus/purpose	Balancing multiple goals; economic development typically dominant	Sustainable multi-purpose, coastal development, accounting for synergies, trade-offs	Hazards, risks, disasters main focus; increasing attention to development	Ecosystem preservation/restoration to protect against CC impacts; make ecosystems more resistant/resilient to CC	Integration of poverty reduction, development and other coastal goals (pro-poor adaptation)
Institutional arrangements	Multi-scalar, separate institutions	Multi-scalar; integration across "silos"	Multi-scalar (different levels emphasized)	Emphasis on local to regional level	Emphasis on local level
Stakeholder engagement	limited	central	varies, central at local level	varies	central
Other traits to compare??					
Other traits to compare??					
Sample applications and critical analyses of approaches (since AR4)	Hansen, 2011; Hallegatte, 2009; Tribbia and Moser, 2008; Van Koningsveld <i>et al.</i> , 2008	Nursey-Bray and Shaw, 2010; Jentoft, 2009; Dawson <i>et al.</i> , 2009; Sales, 2009; McFadden, 2008; Shipman and Stojanovic, 2007; Stojanovic and Ballinger, 2009; Falaleeva <i>et al.</i> , 2011	Romieu <i>et al.</i> , 2010; Mercer, 2010; Mitchell <i>et al.</i> , 2010; Polack, 2010; Gero <i>et al.</i> , 2011; Kirshen <i>et al.</i> , 2011; Halpern <i>et al.</i> , 2008b	Espinosa-Romero <i>et al.</i> , 2011; McGinnis and McGinnis, 2011; Pérez <i>et al.</i> , 2010; Anthony <i>et al.</i> , 2009; Alongi, 2008	van Aalst <i>et al.</i> , 2008; Dumaru, 2010; Mustelin <i>et al.</i> , 2010; Raihan <i>et al.</i> , 2010; Milligan <i>et al.</i> , 2009

Table 5-7: Common barriers to coastal adaptation.

Location	Common Barriers to Coastal Adaptation Identified	Reference
Australia	<ul style="list-style-type: none"> • Polarized views in the community regarding the risk of sea level rise • Among the vocal portion of population that does not recognize threat from sea-level rise, expectations that <ul style="list-style-type: none"> ○ governments or insurance will compensate landholders for loss of property due to sea-level rise ○ governments will fund hard protection against rising seas ○ land owners will be allowed to build defences to protect their property ○ Private property rights should not be revoked under threat from sea-level rise • Concerns about fairness about retreat scheme 	Ryan <i>et al.</i> , 2011
US, Alaska	<ul style="list-style-type: none"> • Currently no government agencies with the mandate or authority to address climate-induced relocation • Lack of financial resources locally or from federal sources to pay for relocation from eroding coastal locations • Assimilation into Western society undermines language, culture, and ties to the land and sea and seriously challenges the resilience of Inuit culture (loss of social institutions of support, traditional ecological knowledge etc.) 	Adger <i>et al.</i> , 2011
Fiji, Rewa Delta	<ul style="list-style-type: none"> • Lack of awareness of climate change/sea-level rise risks • Lack of understanding of climate change (e.g., confusion with variability, natural cycles) • Short-term planning perspectives • Gap between official climate policy position and actual actions • Spiritual beliefs • Traditional governance structures (e.g., departmental divisions, top-down, consultative approach, non-democratic, hierarchical, exclusive) 	Lata and Nunn, 2011
US, Florida Keys	<ul style="list-style-type: none"> • Limited information resulting in lack of awareness • Lack a formal institutional framework necessary to shape and execute adaptation measures (network for monitoring key indicators, coordination mechanism across scales of governance, interagency collaboration) • Insufficient budget for the development of adaptation policies • Lack of direction and leadership • Lack of perceived importance to public officials • Lack of assistance from state and federal agencies • Lack of public demand to take action • Lack of a legal mandate to account for climate change impacts • Lack of perceived solutions • Opposition from stakeholder groups 	Mozumber <i>et al.</i> , 2011
Sweden	<ul style="list-style-type: none"> • Lack of clear institutional frameworks at the national and regional levels (lack of formal, coherent policy from higher level) • Disconnect between technical and strategic planning work related to coastal erosion • Weak vertical administrative interplay (local, regional national) • New proactive integrative policy approach not embraced by those outside the inner circle of erosion managers • Inability to reach general acceptability and organizational mainstreaming of climate concerns • “One-man show” (strong leader in one agency with cemented role and responsibilities) hinders cross-sectoral ownership, learning, and common frames of reference • Professional integrity and inter-departmental rivalry in the way of more 	Storbjörk and Hedrén, 2011; Storbjörk, 2010

	<p>integrated and learning-oriented approaches</p> <ul style="list-style-type: none"> • Time and effort required to change departmental priorities • Differences in professional interests and priorities, administrative cultures and goals • Tensions between different interests, values and priorities (trade-offs) • Short-term planning perspective 	
Bangladesh	<ul style="list-style-type: none"> • Lack of familiarity with the term “sea-level rise” (but clear familiarity with more immediate impacts of SLR) • Perception that SLR and its impacts are not an immediate threat to livelihood • Preference for retreat option decreased with <ul style="list-style-type: none"> ○ greater length of attachment with coastal environment ○ lower wealth and social standing (lack of mobility) ○ lower climate familiarity and resiliency (lack of education, job mobility) ○ stronger local social network ○ lower frequency of current coping and adaptive behavior (threshold of acceptability, fear) ○ higher exposure potential ○ greater tacit knowledge of SLR impacts (sense of manageability, less fatalism) ○ greater access to weather information through radio (increasing precautionary actions) ○ better access to shelters ○ age 	Saroar and Routray, 2010
Pacific Atolls	<ul style="list-style-type: none"> • Limited adaptation options (due to small land area, high population densities, limited economic resources, economic marginalization due to isolation, and generally low levels of human resource development) • Climate change is still a foreign, unfamiliar concept for many • Language barriers (climate change information predominantly in English) • Climate change impacts perceived as occurring in distant places • Weakening of traditional cultural exchange mechanisms (based on reciprocity) • Loss of traditional ecological knowledge with modernization/ • Westernization of culture in some islands • International emigration is perceived as giving up 	Barnett and Campbell, 2010; Adger, <i>et al.</i> 2011
US, Northeast	<ul style="list-style-type: none"> • Regulations restricting fishermen’s ability to switch fisheries when stocks of one species are low • Traditional values and independence-mindedness of fishermen limit willingness to change jobs • Limited extent of higher education and professional training limit job mobility • Limited economic alternatives for fishermen in island communities • Past development and land use decisions in coastal areas restrict perceived and economical options • Expectations of protection and government assistance based on historical experience • Ingrained socioeconomic interests in the status quo 	Moser <i>et al.</i> 2008
US, California	<ul style="list-style-type: none"> • Monetary constraints locally and lack of funding from state and federal sources • Insufficient staff resources and time • Currently pressing issues all-consuming • Lack of legal mandate • Lack of perceived importance • Lack of perceived solution options 	Tribbia and Moser, 2008; Moser and Tribbia, 2006/2007

	<ul style="list-style-type: none"> • Lack of public awareness and demand • Lack of technical assistance from state or federal agencies • Limited social acceptability • Pressure to maintain status quo and stakeholder opposition • Lack of relevant information or science too uncertain • Lack of analytic capacity to use climate change information for decision-making • Lack of boundary organizations connecting climate change science with coastal management 	
United Kingdom	<ul style="list-style-type: none"> • Lack of adequate financial compensation to landowners • Need to provide compensatory habitats under the Habitats Regulations • Lack of public support (esp. locally) • Lack of political acceptance for the loss of existing defence line and lack of support from public opinion • Insufficient consultation • Potential high cost of managed realignment • Potential loss of terrestrial and freshwater habitats • Managed realignment is ineffective if carried out on a piecemeal basis • Lack of access to or information about suitable funding • Insufficient robustness of flood and coastal defences • Difficulty of recreating an environmentally diverse habitat 	Ledoux <i>et al.</i> , 2005
Netherlands	<ul style="list-style-type: none"> • The costs and benefits of the adaptation options can not be estimated with accuracy • For the majority of the options knowledge gaps exist, • data are missing or their reliability is insufficient • methodological difficulties and insufficient quantitative data to run social cost-benefit analysis 	De Bruin <i>et al.</i> , 2009

Note: For studies that produced quantitative results the top three constraints are presented in **bold**.

Table 5-8: Positive synergies and tradeoffs between selected coastal adaptation and mitigation options.

Measure or Option	Positive Implications for Mitigation	Positive Implications for Adaptation	References <small>[REQUEST TO OTHER CHAPTER CO-AUTHORS AND REFS FOR NEW REFS. ALSO: WE MAY BE ABLE TO ADD CONFIDENCE LANGUAGE TO THESE, IF DESIRED/ABLE TO DO SO.]</small>
Coastal seagrass and tidal marsh restoration	Increased carbon storage ^a	Storm buffer, species habitat, fish nursery	Whiting and Chanton, 2001; Turner <i>et al.</i> , 2005; Chmura, 2011; Kennedy and Björk, 2011
Mangrove restoration species	Carbon storage ^a	Habitat and species protection, flood control, soil preservation	Alongi, 2002; Kristensen <i>et al.</i> , 2008; Bouillon <i>et al.</i> , 2011
Reduction/cessation of off-shore oil production	Reduction in liquid fuel-related GHG emissions	Reduced risk of oil spills, reduction of stresses on marine/coastal eco-systems; variable socio-economic impacts on human communities and public health (and thus on vulnerability)	O'Rourke and Connolly, 2003
Increased urban tree cover	Increased carbon storage, shading resulting in lower cooling energy demand	Increased shading, lesser urban heat island, better air quality	Nowak and Crane, 2002; Nowak <i>et al.</i> , 2006; Pataki <i>et al.</i> , 2006; Chen <i>et al.</i> , 2011
Adaptation Measure or Option	Potential Negative Implications for Mitigation		
Desalinization, increased water reuse, groundwater banking and pumping, and inter-basin water transfers (if fossil fuel-based)	Higher ongoing energy consumption to fuel water pumping, storage and transfer processes, increase in GHG emissions		US DOE, 2006; Stokes and Horvath, 2006; Lofman <i>et al.</i> , 2002
Relocation of infrastructure and development out of coastal floodplains	Increase in one-time GHG emissions due to rebuilding of structures; possible increase in sprawl and ongoing transportation-related emissions		Biesbroek <i>et al.</i> , 2009
Building of large dams or massive coastal protection structures	Increased (one-time) energy use and GHG emissions related to construction (cement)		Boden <i>et al.</i> , 2011
Mitigation Measure or Option	Potential Negative Implications for Adaptation		
Reforestation or forest conservation	Negative consequences for rural livelihoods (thus potentially increased vulnerability) if forest ownership and management are not held by local community		Chhatre and Agrawal, 2009
More compact coastal urban design	Potential increase in urban heat island, increased development in floodplains (if present)		Giridharan <i>et al.</i> , 2007

Offshore renewable energy development	Potentially additional drivers on near- and offshore coastal and marine ecosystems and species	Gill, 2005; Boehlert and Gill, 2010
Rapid switch to low-or no-GHG energy sources	Higher energy prices may slow economic development and disproportionately affect low-income populations, increasing their vulnerability or reducing the resources available for adaptation	Tol, 2007

Source: Adapted from Moser (2011) and references cited in Table;

Notes: ^a – DeLaune *et al.* (2011) suggested this benefit may be smaller than previously thought given the losses of sequestered carbon in soils that erode during coastal storms.




Figure 5-1: Coastal zone.




Figure 5-2: The intersection of inland and storm surge flooding. Red shows May 5, 2008 MODIS mapping of the tropical cyclone Nargis storm surge along the Ayeyarwady Delta and to the east, Myanmar. The blue areas to the north were flooded by the river in prior years. (From Brakenridge *et al.*, 2012, submitted in 2011).




Figure 5-3: Time series of modelled gross primary production (A; Gypens *et al.*, 2009) and measured pH_T (B) at a fixed station in the southern North Sea (Borges, 2011). PH is expressed on the total scale. Shown is the regression before and after 1987 (solid lines) and the change in pH expected from increased atmospheric CO₂ alone (broken line).




Figure 5-4: Confidence in Detection and Attribution of observed impacts for coastal systems. Values will be inserted at right positions post FOD, and iterated across chapters to ensure consistency. [Combined one for all coasts still to be developed.]



Figure 5-5: The same coral community before and after a bleaching event in February 2002 at 5 m depth, Halfway Island, Great Barrier Reef. Coral cover at the time of bleaching was 95% almost all of which was severely bleached, resulting in mortality of 20.9% (Elvidge *et al.* 2004). C and D: three CO₂ seeps in Milne Bay Province, Papua New Guinea show that prolonged exposure to high CO₂ is related to fundamental changes in coral reef structures (Fabricius *et al.*, 2011). Coral communities at three high CO₂ (median pH_T 7.7, 7.7 and 8.0), compared with three control sites (median pH_T 8.02), are characterised by significantly reduced coral diversity (-39%), severely reduced structural complexity (-67%), low densities of young corals (-66%) and few crustose coralline algae (-85%). Reef development ceases at pH values below 7.7. Photo credit: R. Berkelmans (A and B) and K. Fabricius (C and D).




Figure 5-6: Percent of reef locations (1°x1° latitude/longitude cells which have coral reefs) that experience no bleaching (green), at least one mild bleaching event (reddish-brown), or at least one severe bleaching event (purple) for each decade. Observed bleaching events are summarized from the ReefBase dataset (Kleypas *et al.* 2008). In the observations, some of the “no bleaching” cells may have experienced bleaching but it was either not observed or not reported. Modeled bleaching events are averages of data from four ensemble runs of the Community Climate System Model version 3 using the standard degree heating month formula (Teneva *et al.*, 2011). The labels of values ≤ 1% are not shown.




Figure 5-7: Paradigm shift in adaptation to rising sea levels in the Netherlands: Source: Stive *et al.*, 2011.